

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|---|-----------|---|
| (51) International Patent Classification ⁶ : A61K 38/00, 39/00, 39/02, C07H 19/00, 21/02, C07K 2/00, 4/04, 14/20, C12P 19/34 | A1 | (11) International Publication Number: WO 97/09994 (43) International Publication Date: 20 March 1997 (20.03.97) |
| (21) International Application Number: PCT/US96/14819 (22) International Filing Date: 16 September 1996 (16.09.96) (30) Priority Data: 08/529,055 15 September 1995 (15.09.95) US (71) Applicant: UAB RESEARCH FOUNDATION [US/US]; UAB Station, Birmingham, AL 35294 (US). (72) Inventors: BRILES, David, E.; 760 Linwood Road, Birmingham, AL 23222 (US). McDANIEL, Larry, S.; 103 Canterbury Place, Ridgeland, MS 39157 (US). SWIATLO, Edwin; 2848 Vestavia Forest Drive, Birmingham, AL 25216 (US). YOTHER, Janet; 2208 Heatherbrook Road, Birmingham, AL 35242 (US). CRAIN, Marilyn, J.; 760 Linwood Road, Birmingham, AL 23222 (US). HOLLINGSHEAD, Susan; 1008 32nd Street South, Birmingham, AL 35205 (US). TART, Rebecca; 7540 Old Fair Ground Road, Benson, NC 27504 (US). BROOKS-WALTER, Alexis; 2136 Brookview Drive, Birmingham, AL 35226 (US). (74) Agents: FROMMER, William, S. et al.; Curtis, Morris & Safford, P.C., 530 Fifth Avenue, New York, NY 10036 (US). | | (81) Designated States: AU, CA, FI, IL, JP, NO, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> |
| (54) Title: PNEUMOCOCCAL GENES, PORTIONS THEREOF, EXPRESSION PRODUCTS THEREFROM, AND USES OF SUCH GENES, PORTIONS AND PRODUCTS (57) Abstract The present invention relates to pneumococcal genes, portions thereof, expression products therefrom and uses of such genes, portions and products; especially to genes of <i>Streptococcus pneumoniae</i> , e.g., the gene encoding pneumococcal surface protein A (PspA), i.e., the <i>pspA</i> gene, the gene encoding pneumococcal surface protein A-like proteins, such as <i>pspA</i> -like genes, e.g., the gene encoding pneumococcal surface protein C (PspC), i.e., the <i>pspC</i> gene, portions of such genes, expression products therefrom, and the uses of such genes, portions thereof and expression products therefrom. | | |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | |
|----|--------------------------|----|--|----|--------------------------|
| AM | Armenia | GB | United Kingdom | MW | Malawi |
| AT | Austria | GE | Georgia | MX | Mexico |
| AU | Australia | GN | Guinea | NE | Niger |
| BB | Barbados | GR | Greece | NL | Netherlands |
| BE | Belgium | HU | Hungary | NO | Norway |
| BF | Burkina Faso | IE | Ireland | NZ | New Zealand |
| BG | Bulgaria | IT | Italy | PL | Poland |
| BJ | Benin | JP | Japan | PT | Portugal |
| BR | Brazil | KE | Kenya | RO | Romania |
| BY | Belarus | KG | Kyrgyzstan | RU | Russian Federation |
| CA | Canada | KP | Democratic People's Republic of Korea | SD | Sudan |
| CF | Central African Republic | KR | Republic of Korea | SE | Sweden |
| CG | Congo | KZ | Kazakhstan | SG | Singapore |
| CH | Switzerland | LI | Liechtenstein | SI | Slovenia |
| CI | Côte d'Ivoire | LK | Sri Lanka | SK | Slovakia |
| CM | Cameroon | LR | Liberia | SN | Senegal |
| CN | China | LT | Lithuania | SZ | Swaziland |
| CS | Czechoslovakia | LU | Luxembourg | TD | Chad |
| CZ | Czech Republic | LV | Latvia | TG | Togo |
| DE | Germany | MC | Monaco | TJ | Tajikistan |
| DK | Denmark | MD | Republic of Moldova | TT | Trinidad and Tobago |
| EE | Estonia | MG | Madagascar | UA | Ukraine |
| ES | Spain | ML | Mali | UG | Uganda |
| FI | Finland | MN | Mongolia | US | United States of America |
| FR | France | MR | Mauritania | UZ | Uzbekistan |
| GA | Gabon | | | VN | Viet Nam |

**PNEUMOCOCCAL GENES, PORTIONS THEREOF,
EXPRESSION PRODUCTS THEREFROM,
AND USES OF SUCH GENES, PORTIONS AND PRODUCTS**

RELATED APPLICATIONS

This application is a continuation-in-part ("CIP"): of application Serial Nos. 08,529,055, filed September 15, 1995, 08/226,844, filed May 29, 1992, 08/093,907, filed May 29, 1992, 07/884,918, filed July 5, 1994 (corresponding to PCT/US93/05191); of application Serial No. 08/482,981, filed June 7, 1995; of application Serial No. 08/458,399, filed June 2, 1995; of application Serial No. 08/446,201, filed May 19, 1995 (as a CIP of USSN 08/246,636); of application Serial No. 08/246,636, filed May 20, 1994 (as a CIP of USSN 08/048,896, filed April 20, 1993 as a CIP of USSN 07/835,698, filed February 12, 1992 as a CIP of USSN 07/656,773); of application Serial 08/319,795, filed October 7, 1994 (as a CIP of USSN 08/246,636); of application Serial No. 08/072,070, filed June 3, 1993; of application Serial No. 07/656,773, filed February 15, 1991 (USSN 656,773 and 835,698 corresponding to Int'l application WO 92/1448); and, each of these applications, as well as each application, document or reference cited in these applications, is hereby incorporated herein by reference. Documents or references are also cited in the following text, either in a Reference List appended to certain Examples, or before the claims, or in the text itself; and, each of these documents or references is hereby expressly incorporated herein by reference.

FIELD OF THE INVENTION

This invention relates to pneumococcal genes, portions thereof, expression products therefrom and uses of such genes, portions and products; especially to genes of *Streptococcus pneumoniae*, e.g., the gene encoding pneumococcal surface protein A (PspA) (said gene being "pspA"), pspA-like genes, pneumococcal surface protein C (PspC) (said gene being "pspC"), portions of such genes, expression products therefrom, and the uses of such genes, portions thereof and expression products therefrom. Such uses include uses of the genes and portions thereof for obtaining expression products by recombinant techniques, as well as for detecting the presence of *Streptococcus pneumoniae* or strains thereof by detecting DNA thereof by hybridization or amplification (e.g., PCR) and hybridization techniques (e.g., obtaining DNA-containing sample, contacting same with genes or fragment under PCR, amplification and/or hybridization conditions, and detecting presence of or isolating hybrid or amplified product). The expression product uses include use in preparing antigenic, immunological or vaccine compositions, for eliciting antibodies, an immunological response (other than or additional to antibodies) or a protective response (including antibody or other immunological response by administering composition to a suitable host); or, the expression product can be for use in detecting the presence of *Streptococcus pneumoniae* by detecting antibodies to *Streptococcus pneumoniae* protein(s) or

antibodies to a portion thereof in a host, e.g., by obtaining an antibody-containing sample from a relevant host, contacting the sample with expression product and detecting binding (for instance by having the product labeled); and, the antibodies generated by the aforementioned compositions are useful in diagnostic or detection kits or assays. Thus, the invention relates to varied compositions of matter and methods for use thereof.

BACKGROUND OF THE INVENTION

Streptococcus pneumoniae is an important cause of otitis media, meningitis, bacteremia and pneumonia. Despite the use of antibiotics and vaccines, the prevalence of pneumococcal infections has declined little over the last twenty-five years.

It is generally accepted that immunity to *Streptococcus pneumoniae* can be mediated by specific antibodies against the polysaccharide capsule of the pneumococcus. However, neonates and young children fail to make an immune response against polysaccharide antigens and can have repeated infections involving the same capsular serotype.

One approach to immunizing infants against a number of encapsulated bacteria is to conjugate the capsular polysaccharide antigens to protein to make them immunogenic. This approach has been successful, for example, with *Haemophilus influenzae b* (see U.S. Patent no. 4,496,538 to Gordon and U.S. Patent no. 4,673,574 to Anderson). However, there are over eighty known capsular

serotypes of *S. pneumoniae* of which twenty-three account for most of the disease. For a pneumococcal polysaccharide-protein conjugate to be successful, the capsular types responsible for most pneumococcal infections would have to be made adequately immunogenic. This approach may be difficult, because the twenty-three polysaccharides included in the presently-available vaccine are not all adequately immunogenic, even in adults.

An alternative approach for protecting children, and also the elderly, from pneumococcal infection would be to identify protein antigens that could elicit protective immune responses. Such proteins may serve as a vaccine by themselves, may be used in conjunction with successful polysaccharide-protein conjugates, or as carriers for polysaccharides.

McDaniel et al. (I), *J. Exp. Med.* 160:386-397, 1984, relates to the production of hybridoma antibodies that recognize cell surface polypeptide(s) on *S. pneumoniae* and protection of mice from infection with certain strains of encapsulated pneumococci by such antibodies. This surface protein antigen has been termed "pneumococcal surface protein A" or PspA for short.

McDaniel et al. (II), *Microbial Pathogenesis* 1:519-531, 1986, relates to studies on the characterization of the PspA. Considerable diversity in the PspA molecule in different strains was found, as were differences in the epitopes recognized by different antibodies.

McDaniel et al. (III), J. Exp. Med. 165:381-394, 1987, relates to immunization of X-linked immunodeficient (XID) mice with non-encapsulated pneumococci expressing PspA, but not isogenic pneumococci lacking PspA, protects mice from subsequent fatal infection with pneumococci.

McDaniel et al. (IV), Infect. Immun., 59:222-228, 1991, relates to immunization of mice with a recombinant full length fragment of PspA that is able to elicit protection against pneumococcal strains of capsular types 6A and 3.

Crain et al, Infect.Immun., 56:3293-3299, 1990, relates to a rabbit antiserum that detects PspA in 100% (n = 95) of clinical and laboratory isolates of strains of *S. pneumoniae*. When reacted with seven monoclonal antibodies to PspA, fifty-seven *S. pneumoniae* isolates exhibited thirty-one different patterns of reactivity.

The PspA protein type is independent of capsular type. It would seem that genetic mutation or exchange in the environment has allowed for the development of a large pool of strains which are highly diverse with respect to capsule, PspA, and possibly other molecules with variable structures. Variability of PspA's from different strains also is evident in their molecular weights, which range from 67 to 99 kD. The observed differences are stably inherited and are not the result of protein degradation.

Immunization with a partially purified PspA from a recombinant λ gt11 clone, elicited protection against challenge with several *S. pneumoniae* strains representing different capsular and PspA types, as described in McDaniel et al. (IV), Infect. Immun. 59:222-228, 1991. Although clones expressing PspA were constructed according to that paper, the product was insoluble and isolation from cell fragments following lysis was not possible.

While the protein is variable in structure between different pneumococcal strains, numerous cross-reactions exist between all PspA's, suggesting that sufficient common epitopes may be present to allow a single PspA or at least a small number of PspA's to elicit protection against a large number of *S. pneumoniae* strains.

In addition to the published literature specifically referred to above, the inventors, in conjunction with co-workers, have published further details concerning PspA's, as follows:

1. Abstracts of 89th Annual Meeting of the American Society for Microbiology, p. 125, item D-257, May 1989;
2. Abstracts of 90th Annual Meeting of the American Society for Microbiology, p. 98, item D-106, May 1990;
3. Abstracts of 3rd International ASM Conference on Streptococcal Genetics, p. 11, item 12, June 1990;
4. Talkington et al, Infect. Immun. 59:1285-1289, 1991;

5. Yother et al (I), J. Bacteriol. 174:601-609, 1992;
and
6. Yother et al (II), J. Bacteriol. 174:610-618,
1992.
7. McDaniel et al (V), Microbiol. Pathogenesis,
13:261-268.

It would be useful to provide PspA or fragments thereof in compositions, including PspA's or fragments from varying strains in such compositions, to provide antigenic, immunological or vaccine compositions; and, it is even further useful to show that the various strains can be grouped or typed, thereby providing a basis for cross-reactivities of PspA's or fragments thereof, and thus providing a means for determining which strains to represent in such compositions (as well as how to test for, detect or diagnose one strain from another).

Further, it would be advantageous to provide a *pspA* - like gene or a *pspC* gene in certain strains, as well as primers (oligonucleotides) for identification of such a gene, as well as of conserved regions in that gene and in *pspA*; for instance, for detecting, determining, isolating, or diagnosing strains of *S. pneumonia*. These uses and advantages, it is believed, have not heretofore been provided in the art.

OBJECTS AND SUMMARY OF THE INVENTION

The invention provides an isolated amino acid molecule comprising residues 1 to 115, 1 to 260, 192 to 588, 192 to 299, or residues 192 to 260 of pneumococcal surface protein A of *Streptococcus pneumoniae*.

The invention further provides an isolated DNA molecule comprising a fragment of a pneumococcal surface protein A gene of *Streptococcus pneumoniae* encoding the isolated amino acid molecule.

The invention also provides PCR primers or hybridization probes comprising the isolated DNA molecule.

The invention additionally provides an antigenic, vaccine or immunological composition comprising the amino acid molecule.

The invention includes an isolated DNA molecule comprising nucleotides 1 to 26, 1967 to 1990, 161 to 187, 1093 to 1117, or 1312 to 1331 or 1333 to 1355 of a pneumococcal surface protein A gene of *Streptococcus pneumoniae*. The DNA molecule can be used as a PCR primer or hybridization probe; and therefore the invention comprehends a PCR primer or hybridization probe comprising the isolated DNA molecule.

The invention also includes an isolated DNA molecule comprising a fragment having homology with a portion of a pneumococcal surface protein A gene of *Streptococcus pneumoniae*. The DNA preferably is the following (which include the portion

having homology and restriction sites, and selection of other restriction sites or sequences for such DNA is within the ambit of the skilled artisan from this disclosure):

```
CCGGATCCAGCTCCTGCACCAAAAAC;  
GCGCGTCGACGGCTTAAACCCATTACCATTTGG;  
CCGGATCCTGAGCCAGAGCAGTTGGCTG;  
CCGGATCCGCTCAAAGAGATTGATGAGTCTG;  
GCGGATCCCGTAGCCAGTCAGTCTAAAGCTG;  
CTGAGTCGACTGGAGTTTCTGGAGCTGGAGC;  
CCGGATCCAGCTCCAGCTCCAGAAACTCCAG;  
GCGGATCCTTGACCAATATTTACGGAGGAGGC;  
GTTTTTGGTGCAGGAGCTGG;  
GCTATGGGCTACAGGTTG;  
CCACCTGTAGCCATAGC;  
CCGCATCCAGCGTGCCTATCTTAGGGGCTGGTT; and  
GCAAGCTTATGATATAGAAATTTGTAAC
```

(thus, the invention broadly comprehends DNA homologous to portions of *pspA*; preferably further including restriction sequences).

These DNA molecules can be used as PCR primers or probes; and thus, the invention comprehends a primer or probe comprising any one or more of these molecules.

The invention further still provides PCR probe(s) which distinguishes between *pspA* and *pspA*-like nucleotide sequence, as well as PCR probe(s) which hybridizes to both *pspA* and *pspA*-like nucleotide sequences.

Additionally, the invention includes a PspA extract prepared by a process comprising: growing pneumococci in a first medium containing choline chloride, eluting live pneumococci with a choline chloride containing salt solution, and growing the pneumococci in a second medium containing an alkanolamine and

substantially no choline; as well as a PspA extract prepared by that process and further comprising purifying PspA by isolation on a choline-Sepharose affinity column. These processes are also included in the invention.

An immunological composition comprising thses extracts is comprehended by the invention, as well as an immunological composition comprising the full length PspA.

A method for enhancing the immunogenicity of a PspA-containing immunological composition comprising, in said composition, the C-terminal portion of PspA, is additionally comprehended, as well.

An immunological composition comprising at least two PspAs. The latter immunological composition can have the PspAs from different groups or families; the groups or families can be based on RFLP or sequence studies (see, e.g., Fig. 13).

Further, the invention provides an isolated amino acid molecule comprising pneumococcal surface protein C, PspC, of *Streptococcus pneumoniae* having an alpha-helical, proline rich and repeat regions, an isolated DNA molecule comprising a pneumococcal surface protein C gene encoding the aforementioned PspC, and primers and hybrization probes consisting essentially of the isolated DNA molecule.

Still further, an isolated amino acid molecule comprising pneumococcal surface protein C, PspC, of *Streptococcus pneumoniae* is provided, having an alpha-helical, proline rich and

repeat regions, having substantial homology with a protection eliciting region of PspA, and an isolated DNA molecule comprising a pneumococcal surface protein C gene encoding the aforementioned PspC, and primers and hybridization probes consisting essentially of the isolated DNA molecule are provided by the present invention.

Additionally, the present invention provides immunological compositions comprising PspC.

These and other embodiments are disclosed or are obvious from the following detailed description.

BRIEF DESCRIPTION OF THE FIGURES

Figures 1A and 1B show: Evaluation of digested plasmid constructs. Fig. 1A: 1% agarose gel electrophoresis of plasmids isolated from transformed *E. coli* BL21(DE3) strains stained with ethidium bromide. Lane 1: 1 kb DNA ladder (sizes noted in kb), lane 2: pRCT125; lane 3: pRC105, lane 4: DBL5 pspA insert, lane 5: pRCT113, lane 6: BG9739 pspA insert, lane 7: pRCT117, and lane 8: L81905 pspA insert. Fig. 1B: Corresponding Southern blot of gel in Fig. 1A probed with full-length Rx1 pspA and hybridization detected as described in Example 1. The arrow indicates the 1.2 kb pspA digested inserts from plasmid constructs and the PCR-amplified pspA fragments from the pneumococcal donor strains used in cloning.

Figure 2 shows: Evaluation of strain RCT105 cell fractions containing truncated DBL5 PspA. Proteins from *E. coli*

cell fractions were resolved by 10% SDS-PAGE, transferred to NC, and probed with MAb XiR278. Lane 1: molecular weight markers (noted in kDa), lane 2: full-length, native DBL5 PspA, lane 3: uninduced cells, lanes 4-6: induced cells; 1 hr, 2 hr, and 3 hr of IPTG induction respectively, lane 7: periplasmic proteins, lane 8: cytoplasmic proteins, and lane 9: insoluble cell wall/membrane material.

Figure 3 shows: SDS-PAGE of R36A PspA (80 ng) column isolated from CDM-ET and an equal volume of an equivalent WG44.1 prep. Identical gels are shown stained with Bio-Rad silver kit (A) or immunoblotted with PspA MAb XiR278(B). The PspA isolated from R36A shows the characteristic monomer (84 kDa) and dimer bands.

Figure 4 shows: Cell lysates of pneumococcal isolates MC27 and MC28 were subjected to SDS-PAGE and transferred to nitrocellulose for Western blotting with seven MAb to PspA. 7D2 detected a protein of 82 kDa in each isolate and XiR278 and 2A4 detected a protein of 190 kDa in each isolate. MAb Xi64, Xi126, 1A4 and SR4W4 were not reactive. Strains MC25 and MC26 yielded identical results.

Figure 5 (Figs. 5A and 5B) shows: Southern blot of *Hind* III digest of MC25-MC28 chromosomal DNA developed at a stringency greater than 95 percent. A digest of Rx1 DNA was used as a comparison. The blot was probed with LSMpspA13/2, a full length Rx1 probe (Fig. 5) and LSMpspA12/6 a 5' probe of Rx1 pspA

(Fig. 5). The same concentration of Rx1 DNA was used in both panels, but the concentrations of MC25-MC28 DNA in Fig. 5B were half that used in Fig. 5A to avoid detection of partial digests.

Figure 6 shows: RFLP of amplified *pspA*. *PspA* from MC25 was amplified by PCR using 5' and 3' primers for *pspA* (LSM13 and LSM, respectively). The amplified DNA was digested with individual restriction endonucleases prior to electrophoresis and staining with ethidium bromide. Lane 1 *BclI*, Lane 2 *BAMHI*, Lane 3 *BstNI*, Lane 4 *PstI*, Lane 5 *SacI*, Lane 6 *EcoRI*, Lane 7 *SmaI*, Lane 8 *KpnI*.

Figure 7 shows: A depiction of *PspA* showing the relative location and orientation of the oligonucleotides.

Figure 8 shows: Derivatives of the *S. pneumoniae* D39-Rx1 family.

Figures 9 to 10 show: Electrophoresis of *pspA* or amplified *pspA* product with *HhaI* (Fig. 9), *Sau3AI* (Fig. 10).

Figure 11 shows: RFLP pattern of two isolates from six families.

Figure 12 shows: RFLP pattern of two isolates from six families (using products from amplification with SKH2 and LSM13).

Figure 13 shows: Sequence primarily in the N-terminal half of *PspA*.

Figure 14 shows: Cell lysates of pneumococcal isolates MC27 and MC28, subjected to SDS-PAGE and Western blotting with seven MAbs to *PspA*; 7D2 detected a protein of 82 kDa in each

isolate, and Xi278 and 2A4 detected a protein of 190 kDa in each isolate; MAbs Xi64, Xi126, 1A4 and SR4W4 were not reactive; strains MC25 and MC26 yielded identical results (not shown).

Figure 15A and 15B show: a Southern blot of *Hind* III digest of MC25-28 chromosomal DNA, using a digest of Rx1 DNA as a comparison; the blot was probed with LSMpspA13/2, a full length Rx1 probe (A), and LSMpspA12/6, a 5' probe of Rx1 *pspA* (B); the same concentration of Rx1 DNA was used in both panels, but the concentrations of MC25-28 DNA in B were half that used in A to avoid detection of partial digests.

Figures 15C and 15D show: the nucleotide sequences of primers LSM13, LSM2, LSM12 and LSM6, and that of probes LSMpspA13/2 and LSMpspA12/6.

Figure 16 shows: RFLP of amplified *pspA*, wherein PspA from MC25 was amplified by PCR using 5' and 3' primers for *pspA* (LSM13 and LSM2, respectively); the amplified DNA was digested with individual restriction endonucleases prior to electrophoresis and staining with ethidium bromide; *Bcl* I was used in lane 1; *Bam*H I was used in lane 2; *Bst*N I was used in lane 3; *Pst* I was used in lane 4; *Sac* I was used in lane 5; *Eco*R I was used in lane 6; *Sma* I was used in lane 7; and *Kpn* I was used in lane 8.

Figure 17 shows: position and orientation of oligonucleotides relative to domains encoded by *pspA*; numbers along the bottom of the Figure represent amino acids in the

0

mature PspA polypeptide from strain Rx1, and arrows represent the relative position (not to scale) and orientation of oligonucleotides.

Figure 18 shows: a restriction map of the pZero vector.

Figure 19 shows: the nucleotide sequences of SKH2, LSM13, N192 and C588.

Figure 20 shows: a comparison of the structural motifs of PspA and PspC; PspA has a smaller alpha-helical region, and does not contain the direct repeats within the alpha-helix (indicated by the dashed lines); the alpha-helical regions which are homologous between PspA and PspC are indicated by the dashed lines); the alpha-helical regions which are homologous between PspA and PspC are indicated by the striped pattern; and PCR primers are indicated by the arrows.

Figure 21 shows: the amino acid and nucleotide sequence of PspC, wherein the putative -10 and -35 regions are underlined, and the ribosomal binding site is in lower case.

Figure 22 shows: the Bestfit analysis of PspA and PspC; percent identity is 69% and percent similarity is 77%; amino acids of PspA are on the bottom line (1-588) and amino acids of PspC are on the top line (249-891), and a dashed line indicated identity.

Figure 23 shows: the coiled coil motif of the alpha-helix of PspC; amino acids that are not in the coiled coil motif are in the right column.

Figure 24 shows: a matrix plot comparison of the repeat regions of the alpha-helical region of PspC.

Figure 25 shows: the sequence of the alpha helical and proline regions of LXS532 (PspC.D39).

Figure 26 shows: a comparison of nucleotides of *pspA.Rx1* to *pspC.D39*.

Figure 27 shows: a BESTFIT analysis of *pspC.EF6797* and *pspC.D39*.

Figure 28 shows: the amino acid comparison of PspC of EF6797 and D39.

Figure 29 shows: the amino acid comparison of PspC.D39 and PspA.Rx1.

DETAILED DESCRIPTION

Knowledge of and familiarity with the applications incorporated herein by reference is assumed; and, those applications disclose the sequence of *pspA* as well as certain portions thereof, and PspA and compositions containing PspA.

As discussed above and in the following Examples, the invention relates to truncated PspA, e.g., PspA C-terminal to position 192 such as a.a. 192-588 ("BC100") 192-299 and 192-260 of PspA eliciting cross-protection, as well as to DNA encoding such truncated PspA (which amplify the coding for these amino acid regions homologous to most PspAs).

The invention further relates to a *pspA*-like gene, or a *pspC* gene and portions thereof (e.g., probes, primers) which can

hybridize thereto and/or amplify that gene, as well as to DNA molecules which hybridize to *pspA*, so that one can, by hybridization assay and/or amplification, ascertain the presence of a particular pneumococcal strain; and, the invention provides that a PspC can be produced by the *pspA*-like or *pspC* sequence (which PspC can be used like PspA).

Indeed, the invention further relates to oligonucleotide probes and/or primers which react with *pspA* and/or *pspC* of many, if not all, strains, so as to permit identification, detection or diagnosis of any pneumococcal strain, as well as to expression products of such probes and/or primers, which can provide cross-reactive epitopes of interest.

The repeat region of *pspA* and/or *pspC* is highly conserved such that the present invention provides oligonucleotide probes or primers to this region reactive with most, if not all strains, thereby providing diagnostic assays and a means for identifying epitopes of interest.

The invention demonstrates that the *pspC* gene is homologous to the *pspA* gene in the leader sequence, first portion of the proline-rich region and in the repeat region; but, these genes differ in the second portion of their proline-rich regions and at the very 3' end of the gene encoding the 17 amino acid tail of PspA. The product of the *pspC* gene is expected to lack a C-terminal tail, suggesting different anchoring than PspA. Drug interference with functions such as surface binding of the coding

for repeat regions of *pspA* and the *pspC* genes, or with the repeat regions of the expression products, is therefore a target for intervention of pneumococcal infection.

Further still, the invention provides evidence of additional *pspA* homologous sequences, in addition to *pspA* and the *pspC* sequence. The invention, as mentioned above, includes oligonucleotide probes or primers which distinguish between *pspA* and the *pspC* sequence, e.g., LSM1 and LSM2, useful for diagnostic detecting, or isolating purposes; and LSM1 and LSM10 or LSM1 and LSM7 which amplify a portion of the *pspC* gene, particularly the portion of that gene which encodes an antigenic, immunological or protective protein.

The invention further relates to a method for the isolation of native PspA by growth of pneumococci medium containing high concentrations of (about 0.9% to about 1.4%, preferably 1.2%) choline chloride, elation of live pneumococci with a salt solution containing choline chloride, e.g., about 1% about 3%, preferably 2% choline chloride, and growth of pneumococci in medium in which the choline in the medium has been almost or substantially completely replaced with a lower alkanolamine, e.g., C₁-C₆, preferably C₂ alkanolamine, i.e., preferably C₂ alkanolamine, i.e., preferably ethanolamine (e.g., 0.0000005% to 0.0000015%, preferably 0.000001% choline chloride plus 0.02% to 0.04% alkanolamine (ethanolamine), preferably 0.03%). PspA from such pneumococci is then preferably isolated

from a choline-sepharose affinity column, thereby providing highly purified PspA. Such isolated and/or purified PspA is highly immunogenic and is useful in antigenic, immunological or vaccine composition.

Indeed, the growth media of the pneumococci grown in the presence of the alkanolamine (rather than choline) contains PspA and is itself highly immunogenic and therefore useful as an antigenic, immunological or vaccine composition; and, is rather inexpensive to produce. Per microgram of PspA, the PspA in the alkanolamine medium is much more protective than PspA isolated by other means, e.g., from extracts. Perhaps, without wishing to necessarily be bound by any one particular theory, there is a synergistic effect upon PspA by the other components present prior to isolation, or simply PspA is more protective (more antigenic) prior to isolation and/or purification (implying a possibility of some loss of activity from the step of isolation and/or purification).

The invention further relates to the N-terminal 115 amino acids of PspA, which is useful for compositions comprising an epitope of interest, immunological or vaccine compositions, as well as the DNA coding therefor, which is useful in preparing these N-terminal amino acids by recombination, or for use as probes and/or primers for hybridization and/or amplification for identification, detection or diagnosis purposes.

The invention further demonstrates that there is a grouping among the *pspA* RFLP families. This provides a method of identifying families of different PspAs based on RFLP pattern of *pspAs*, as well as a means for obtaining diversity of PspAs in an antigenic, immunological or vaccine composition; and, a method of characterizing clonotypes of PspA based on RFLP patterns of PspA. And, the invention thus provides oligonucleotides which permit amplification of most, e.g., a majority, if not all of *S. pneumoniae* and thereby permit RFLP analysis of a majority, if not all, *S. pneumoniae*.

The invention also provides PspC, having an approximate molecular weight of 105 kD, with an estimated pI of 6.09, and comprising an alpha-helical region, followed by a proline-rich domain and repeat region. A major cross-protective region of PspA comprises the C-terminal third of the alpha-helical region (between residues 192 and 260 of PspA), which region accounts for the binding of 4 of 5 cross-protective MAb, and PspA fragments comprising this region can elicit cross-protective immunity in mice. Homology between PspC and PspA begins at amino acid 148 of PspA, thus including the region from 192 to 299, and including the entire PspC sequence C-terminal of amino acid 486. Due to the substantial sequence homology between PspA and PspC in a region comprising the epitopes of interest, known to be protection eliciting, PspC is likely to comprise epitopes of interest similar to those found in PspA. Antibodies specific for

this region of PspA, i.e., between amino acids 148 and 299, should cross-react with PspC, and thus afford protection by reacting with PspC and PspA. Similarly, immunization with PspC would be expected to elicit antibodies cross-protective against PspA.

An epitope of interest is an antigen or immunogen or immunologically active fragment thereof from a pathogen or toxin of veterinary or human interest.

The present invention provides an immunogenic, immunological or vaccine composition containing the pneumococcal epitope of interest, and a pharmaceutically acceptable carrier or diluent. An immunological composition containing the pneumococcal epitope of interest, elicits an immunological response - local or systemic. The response can, but need not be, protective. An immunogenic composition containing the pneumococcal epitope of interest, likewise elicits a local or systemic immunological response which can, but need not be, protective. A vaccine composition elicits a local or systemic protective response. Accordingly, the terms "immunological composition" and "immunogenic composition" include a "vaccine composition" (as the two former terms can be protective compositions).

The invention therefore also provides a method of inducing an immunological response in a host mammal comprising administering to the host an immunogenic, immunological or

vaccine composition comprising the pneumococcal epitope of interest, and a pharmaceutically acceptable carrier or diluent.

The DNA encoding the pneumococcal epitope of interest can be DNA which codes for full length PspA, PspC, or fragments thereof. A sequence which codes for a fragment of PspA or PspC can encode that portion of PspA or PspC which contains an epitope of interest, such as a protection-eliciting epitope of the protein.

Regions of PspA and PspC have been identified from the Rx1 strain of *S. pneumoniae* which not only contain protection-eliciting epitopes, but are also sufficiently cross-reactive with other PspAs from other *S. pneumoniae* strains so as to be suitable candidates for the region of PspA to be incorporated into a vaccine, immunological or immunogenic composition. Epitopic regions of PspA include residues 1 to 115, 1 to 314, 192 to 260 and 192 to 588. DNA encoding fragments of PspA can comprise DNA which codes for the aforementioned epitopic regions of PspA; or it can comprise DNA encoding overlapping fragments of PspA, e.g., fragment 192 to 588 includes 192 to 260, and fragment 1 to 314 includes 1 to 115 and 192 to 260.

As to epitopes of interest, one skilled in the art can determine an epitope of immunodominant region of a peptide or polypeptide and ergo the coding DNA therefor from the knowledge of the amino acid and corresponding DNA sequences of the peptide or polypeptide, as well as from the nature of particular amino

acids (e.g., size, charge, etc.) and the codon dictionary, without undue experimentation.

A general method for determining which portions of a protein to use in an immunological composition focuses on the size and sequence of the antigen of interest. "In general, large proteins, because they have more potential determinants are better antigens than small ones. The more foreign an antigen, that is the less similar to self configurations which induce tolerance, the more effective it is in provoking an immune response." Ivan Roitt, Essential Immunology, 1988.

As to size, the skilled artisan can maximize the size of the protein encoded by the DNA sequence to be inserted into the viral vector (keeping in mind the packaging limitations of the vector). To minimize the DNA inserted while maximizing the size of the protein expressed, the DNA sequence can exclude introns (regions of a gene which are transcribed but which are subsequently excised from the primary RNA transcript).

At a minimum, the DNA sequence can code for a peptide at least 8 or 9 amino acids long. This is the minimum length that a peptide needs to be in order to stimulate a CD4+ T cell response (which recognizes virus infected cells or cancerous cells). A minimum peptide length of 13 to 25 amino acids is useful to stimulate a CD8+ T cell response (which recognizes special antigen presenting cells which have engulfed the pathogen). See Kendrew, *supra*. However, as these are minimum

lengths, these peptides are likely to generate an immunological response, i.e., an antibody or T cell response; but, for a protective response (as from a vaccine composition), a longer peptide is preferred.

With respect to the sequence, the DNA sequence preferably encodes at least regions of the peptide that generate an antibody response or a T cell response. One method to determine T and B cell epitopes involves epitope mapping. The protein of interest "is fragmented into overlapping peptides with proteolytic enzymes. The individual peptides are then tested for their ability to bind to an antibody elicited by the native protein or to induce T cell or B cell activation. This approach has been particularly useful in mapping T-cell epitopes since the T cell recognizes short linear peptides complexed with MHC molecules. The method is less effective for determining B-cell epitopes" since B cell epitopes are often not linear amino acid sequence but rather result from the tertiary structure of the folded three dimensional protein. Janis Kuby, Immunology, (1992) pp. 79-80.

Another method for determining an epitope of interest is to choose the regions of the protein that are hydrophilic. Hydrophilic residues are often on the surface of the protein and therefore often the regions of the protein which are accessible to the antibody. Janis Kuby, Immunology, (1992) P. 81.

Yet another method for determining an epitope of interest is to perform an X-ray crystallographic analysis of the antigen (full length)-antibody complex. Janis Kuby, Immunology, (1992) p. 80.

Still another method for choosing an epitope of interest which can generate a T cell response is to identify from the protein sequence potential HLA anchor binding motifs which are peptide sequences which are known to be likely to bind to the MHC molecule.

The peptide which is a putative epitope, to generate a T cell response, should be presented in a MHC complex. The peptide preferably contains appropriate anchor motifs for binding to the MHC molecules, and should bind with high enough affinity to generate an immune response. Factors which can be considered are: the HLA type of the patient (vertebrate, animal or human) expected to be immunized, the sequence of the protein, the presence of appropriate anchor motifs and the occurrence of the peptide sequence in other vital cells.

An immune response is generated, in general, as follows: T cells recognize proteins only when the protein has been cleaved into smaller peptides and is presented in a complex called the "major histocompatibility complex MHC" located on another cell's surface. There are two classes of MHC complexes - class I and class II, and each class is made up of many different

alleles. Different patients have different types of MHC complex alleles; they are said to have a 'different HLA type'.

Class I MHC complexes are found on virtually every cell and present peptides from proteins produced inside the cell. Thus, Class I MHC complexes are useful for killing cells which when infected by viruses or which have become cancerous and as the result of expression of an oncogene. T cells which have a protein called CD4 on their surface, bind to the MHC class I cells and secrete lymphokines. The lymphokines stimulate a response; cells arrive and kill the viral infected cell.

Class II MHC complexes are found only on antigen-presenting cells and are used to present peptides from circulating pathogens which have been endocytosed by the antigen-presenting cells. T cells which have a protein called CD8 bind to the MHC class II cells and kill the cell by exocytosis of lytic granules.

Some guidelines in determining whether a protein is an epitopes of interest which will stimulate a T cell response, include: Peptide length - the peptide should be at least 8 or 9 amino acids long to fit into the MHC class I complex and at least 13-25 amino acids long to fit into a class II MHC complex. This length is a minimum for the peptide to bind to the MHC complex. It is preferred for the peptides to be longer than these lengths because cells may cut the expressed peptides. The peptide should contain an appropriate anchor motif which will enable it to bind

to the various class I or class II molecules with high enough specificity to generate an immune response (See Bocchia, M. et al, Specific Binding of Leukemia Oncogene Fusion Protein Peptides to HLA Class I Molecules, Blood 85:2680-2684; Englehard, VH, Structure of peptides associated with class I and class II MHC molecules Ann. Rev. Immunol. 12:181 (1994)). This can be done, without undue experimentation, by comparing the sequence of the protein of interest with published structures of peptides associated with the MHC molecules. Protein epitopes recognized by T cell receptors are peptides generated by enzymatic degradation of the protein molecule and are presented on the cell surface in association with class I or class II MHC molecules.

Further, the skilled artisan can ascertain an epitope of interest by comparing the protein sequence with sequences listed in the protein data base. Regions of the protein which share little or no homology are better choices for being an epitope of that protein and are therefore useful in a vaccine or immunological composition. Regions which share great homology with widely found sequences present in vital cells should be avoided.

Even further, another method is simply to generate or express portions of a protein of interest, generate monoclonal antibodies to those portions of the protein of interest, and then ascertain whether those antibodies inhibit growth *in vitro* of the pathogen from which the protein was derived. The

skilled artisan can use the other guidelines set forth in this disclosure and in the art for generating or expressing portions of a protein of interest for analysis as to whether antibodies thereto inhibit growth in vitro. For example, the skilled artisan can generate portions of a protein of interest by: selecting 8 to 9 or 13 to 25 amino acid length portions of the protein, selecting hydrophilic regions, selecting portions shown to bind from X-ray data of the antigen (full length)-antibody complex, selecting regions which differ in sequence from other proteins, selecting potential HLA anchor binding motifs, or any combination of these methods or other methods known in the art.

Epitopes recognized by antibodies are expressed on the surface of a protein. To determine the regions of a protein most likely to stimulate an antibody response one skilled in the art can preferably perform an epitope map, using the general methods described above, or other mapping methods known in the art.

As can be seen from the foregoing, without undue experimentation, from this disclosure and the knowledge in the art, the skilled artisan can ascertain the amino acid and corresponding DNA sequence of an epitope of interest for obtaining a T cell, B cell and/or antibody response. In addition, reference is made to Gefter et al., U.S. Patent No. 5,019,384, issued May 28, 1991, and the documents it cites, incorporated herein by reference (Note especially the "Relevant Literature" section of this patent, and column 13 of this patent

which discloses that: "A large number of epitopes have been defined for a wide variety of organisms of interest. Of particular interest are those epitopes to which neutralizing antibodies are directed. Disclosures of such epitopes are in many of the references cited in the Relevant Literature section.")

Further, the invention demonstrates that more than one serologically complementary PspA molecule can be in an antigenic, immunological or vaccine composition, so as to elicit better response, e.g., protection, for instance, against a variety of strains of pneumococci; and, the invention provides a system of selecting PspAs for a multivalent composition which includes cross-protection evaluation so as to provide a maximally efficacious composition.

The determination of the amount of antigen, e.g., PspA or truncated portion thereof and optional adjuvant in the inventive compositions and the preparation of those compositions can be in accordance with standard techniques well known to those skilled in the pharmaceutical or veterinary arts. In particular, the amount of antigen and adjuvant in the inventive compositions and the dosages administered are determined by techniques well known to those skilled in the medical or veterinary arts taking into consideration such factors as the particular antigen, the adjuvant (if present), the age, sex, weight, species and condition of the particular patient, and the route of

administration. For instance, dosages of particular PspA antigens for suitable hosts in which an immunological response is desired, can be readily ascertained by those skilled in the art from this disclosure (see, e.g., the Examples), as is the amount of any adjuvant typically administered therewith. Thus, the skilled artisan can readily determine the amount of antigen and optional adjuvant in compositions and to be administered in methods of the invention. Typically, an adjuvant is commonly used as 0.001 to 50 wt% solution in phosphate buffered saline, and the antigen is present on the order of micrograms to milligrams, such as about 0.0001 to about 5 wt%, preferably about 0.0001 to about 1 wt%, most preferably about 0.0001 to about 0.05 wt% (see, e.g., Examples below or in applications cited herein).

Typically, however, the antigen is present in an amount on the order of micrograms to milligrams, or, about 0.001 to about 20 wt%, preferably about 0.01 to about 10 wt%, and most preferably about 0.05 to about 5 wt% (see, e.g., Examples below).

Of course, for any composition to be administered to an animal or human, including the components thereof, and for any particular method of administration, it is preferred to determine therefor: toxicity, such as by determining the lethal dose (LD) and LD₅₀ in a suitable animal model e.g., rodent such as mouse; and, the dosage of the composition(s), concentration of components therein and timing of administering the composition(s), which elicit a suitable immunological response,

such as by titrations of sera and analysis thereof for antibodies or antigens, e.g., by ELISA and/or RFFIT analysis. Such determinations do not require undue experimentation from the knowledge of the skilled artisan, this disclosure and the documents cited herein. And, the time for sequential administrations can be ascertained without undue experimentation.

Examples of compositions of the invention include liquid preparations for orifice, e.g., oral, nasal, anal, vaginal, peroral, intragastric, mucosal (e.g., perlingual, alveolar, gingival, olfactory or respiratory mucosa) etc., administration such as suspensions, syrups or elixirs; and, preparations for parenteral, subcutaneous, intradermal, intramuscular or intravenous administration (e.g., injectable administration), such as sterile suspensions or emulsions. Such compositions may be in admixture with a suitable carrier, diluent, or excipient such as sterile water, physiological saline, glucose or the like. The compositions can also be lyophilized. The compositions can contain auxiliary substances such as wetting or emulsifying agents, pH buffering agents, gelling or viscosity enhancing additives, preservatives, flavoring agents, colors, and the like, depending upon the route of administration and the preparation desired. Standard texts, such as "REMINGTON'S PHARMACEUTICAL SCIENCE", 17th edition, 1985, incorporated herein by reference, may be consulted to prepare suitable preparations, without undue experimentation.

Compositions of the invention, are conveniently provided as liquid preparations, e.g., isotonic aqueous solutions, suspensions, emulsions or viscous compositions which may be buffered to a selected pH. If digestive tract absorption is preferred, compositions of the invention can be in the "solid" form of pills, tablets, capsules, caplets and the like, including "solid" preparations which are time-released or which have a liquid filling, e.g., gelatin covered liquid, whereby the gelatin is dissolved in the stomach for delivery to the gut. If nasal or respiratory (mucosal) administration is desired, compositions may be in a form and dispensed by a squeeze spray dispenser, pump dispenser or aerosol dispenser. Aerosols are usually under pressure by means of a hydrocarbon. Pump dispensers can preferably dispense a metered dose or, a dose having a particular particle size.

Compositions of the invention can contain pharmaceutically acceptable flavors and/or colors for rendering them more appealing, especially if they are administered orally. The viscous compositions may be in the form of gels, lotions, ointments, creams and the like and will typically contain a sufficient amount of a thickening agent so that the viscosity is from about 2500 to 6500 cps, although more viscous compositions, even up to 10,000 cps may be employed. Viscous compositions have a viscosity preferably of 2500 to 5000 cps, since above that range they become more difficult to administer. However, above

that range, the compositions can approach solid or gelatin forms which are then easily administered as a swallowed pill for oral ingestion.

Liquid preparations are normally easier to prepare than gels, other viscous compositions, and solid compositions. Additionally, liquid compositions are somewhat more convenient to administer, especially by injection or orally, to animals, children, particularly small children, and others who may have difficulty swallowing a pill, tablet, capsule or the like, or in multi-dose situations. Viscous compositions, on the other hand, can be formulated within the appropriate viscosity range to provide longer contact periods with mucosa, such as the lining of the stomach or nasal mucosa.

Obviously, the choice of suitable carriers and other additives will depend on the exact route of administration and the nature of the particular dosage form, e.g., liquid dosage form [e.g., whether the composition is to be formulated into a solution, a suspension, gel or another liquid form], or solid dosage form [e.g., whether the composition is to be formulated into a pill, tablet, capsule, caplet, time release form or liquid-filled form].

Solutions, suspensions and gels, normally contain a major amount of water (preferably purified water) in addition to the antigen, lipoprotein and optional adjuvant. Minor amounts of other ingredients such as pH adjusters (e.g., a base such as

NaOH), emulsifiers or dispersing agents, buffering agents, preservatives, wetting agents, jelling agents, (e.g., methylcellulose), colors and/or flavors may also be present. The compositions can be isotonic, i.e., it can have the same osmotic pressure as blood and lacrimal fluid.

The desired isotonicity of the compositions of this invention may be accomplished using sodium chloride, or other pharmaceutically acceptable agents such as dextrose, boric acid, sodium tartrate, propylene glycol or other inorganic or organic solutes. Sodium chloride is preferred particularly for buffers containing sodium ions.

Viscosity of the compositions may be maintained at the selected level using a pharmaceutically acceptable thickening agent. Methylcellulose is preferred because it is readily and economically available and is easy to work with. Other suitable thickening agents include, for example, xanthan gum, carboxymethyl cellulose, hydroxypropyl cellulose, carbomer, and the like. The preferred concentration of the thickener will depend upon the agent selected. The important point is to use an amount which will achieve the selected viscosity. Viscous compositions are normally prepared from solutions by the addition of such thickening agents.

A pharmaceutically acceptable preservative can be employed to increase the shelf-life of the compositions. Benzyl alcohol may be suitable, although a variety of preservatives

including, for example, parabens, thimerosal, chlorobutanol, or benzalkonium chloride may also be employed. A suitable concentration of the preservative will be from 0.02% to 2% based on the total weight although there may be appreciable variation depending upon the agent selected.

Those skilled in the art will recognize that the components of the compositions must be selected to be chemically inert with respect to the PspA antigen and optional adjuvant. This will present no problem to those skilled in chemical and pharmaceutical principles, or problems can be readily avoided by reference to standard texts or by simple experiments (not involving undue experimentation), from this disclosure and the documents cited herein.

The immunologically effective compositions of this invention are prepared by mixing the ingredients following generally accepted procedures. For example the selected components may be simply mixed in a blender, or other standard device to produce a concentrated mixture which may then be adjusted to the final concentration and viscosity by the addition of water or thickening agent and possibly a buffer to control pH or an additional solute to control tonicity. Generally the pH may be from about 3 to 7.5. Compositions can be administered in dosages and by techniques well known to those skilled in the medical and veterinary arts taking into consideration such factors as the age, sex, weight, and condition of the particular

patient or animal, and the composition form used for administration (e.g., solid vs. liquid). Dosages for humans or other mammals can be determined without undue experimentation by the skilled artisan, from this disclosure, the documents cited herein, the Examples below (e.g., from the Examples involving mice).

Suitable regimes for initial administration and booster doses or for sequential administrations also are variable, may include an initial administration followed by subsequent administrations; but nonetheless, may be ascertained by the skilled artisan, from this disclosure, the documents cited herein, and the Examples below.

PCR techniques for amplifying sample DNA for diagnostic detection or assay methods are known from the art cited herein and the documents cited herein (see Examples), as are hybridization techniques for such methods. And, without undue experimentation, the skilled artisan can use gene products and antibodies therefrom in diagnostic, detection or assay methods by procedures known in the art.

The following Examples are provided for illustration and are not to be considered a limitation of the invention.

EXAMPLES

EXAMPLE 1 - Truncated *Streptococcus pneumoniae* PspA Molecules Elicit Cross-Protective Immunity Against Pneumococcal Challenge

Since the isolation of *S. pneumoniae* from human saliva in 1881 and its subsequent connection with lobar pneumonia two years later, human disease resulting from pneumococcal infection has been associated with a significant degree of morbidity and mortality. A recent survey of urgently needed vaccines in the developing and developed world places an improved pneumococcal vaccine among the top three vaccine priorities of industrialized countries. The currently licensed vaccine is a 23-valent composition of pneumococcal capsular polysaccharides that is only about 60% effective in the elderly and due to poor efficacy is not recommended for use in children below two years of age. Furthermore the growing frequency of multi-drug resistant strains of *S. pneumoniae* being isolated accentuates the need for a more effective vaccine to prevent pneumococcal infections.

The immunogenic nature of proteins makes them prime targets for new vaccine strategies. Pneumococcal molecules being investigated as potential protein vaccine candidates include pneumolysis, neuraminidase, autolysin and PspA. All of these proteins are capable of eliciting immunity in mice resulting in extension of life and protection against death with challenge doses near the LD₅₀. PspA is unique among these macromolecules

in that it can elicit antibodies in animals that protect against inoculums 100-fold greater than the LD₅₀.

PspA is a surface-exposed protein with an apparent molecular weight of 67-99 kDa that is expressed by all clinically relevant *S. pneumoniae* strains examined to date. Though PspAs from different pneumococcal strains are serologically variable, many PspA antibodies exhibit cross-reactivities with PspAs from unrelated strains. Upon active immunization with PspA, mice generate PspA antibodies that protect against subsequent challenge with diverse strains of *S. pneumoniae*. The immunogenic and protection-eliciting properties of PspA suggest that it may be a good candidate molecule for a protein-based pneumococcal vaccine.

Four distinct domains of PspA have been identified based on DNA sequence. They include a N-terminal highly charged alpha-helical region, a proline-rich 82 amino acid stretch, a C-terminal repeat segment comprised of ten 20-amino acid repeat sequences, and a 17-amino acid tail. A panel of MAbs to Rx1 PspA have been produced and the binding sites of nine of these MAbs were recently localized within the Rx1 *pspA* sequence in the alpha-helical region. Five of the Rx1 MAbs were protective in mice infected with a virulent pneumococcal strain, WU2. Four of these five protective antibodies were mapped to the distal third (amino acids 192-260) of the alpha-helical domain of Rx1 PspA.

Truncated PspAs containing amino acids 192-588 or 192-299, from pneumococcal strain Rx1 were cloned and the recombinant proteins expressed and evaluated for their ability to elicit protection against subsequent challenge with *S. pneumoniae* WU2. As with full-length Rx1 PspA, both truncated PspAs containing the distal alpha-helical region protected mice against fatal WU2 pneumococcal infection. However, the recombinant PspA fragment extending from amino acid 192 to 588 was more immunogenic than the smaller fragment, probably due to its larger size. In addition, the protection elicited by the amino acid fragment 192-588 of Rx1 was comparable to that elicited by full-length Rx1 PspA. Therefore, cross-protective epitopes of other PspAs were also sought in the C-terminal two-thirds of the molecule. As discussed below, PspAs homologous to amino acids 192-588 of strain Rx1 were amplified by PCR, cloned, and expressed in *E. coli*. Then three recombinant PspAs, from capsule type 4 and 5 strains, were evaluated for their ability to confer cross-protection against challenge strains of variant capsular types. The data demonstrate that the truncated PspAs from capsular type 4 and 5 strains collectively protect against or early death caused by challenge with capsular type 4 and 5 parental strains as well as type 3, 6A, and 6B *S. pneumoniae*.

Bacterial strains and culture conditions. All pneumococci were from the culture collection of this laboratory, and have been described (Yother, J. et al., Infect. Immun. 1982;

36: 184-188; Briles, D.E., et al., Infect. Immun. 1992; 60: 111-116; McDaniel, L.S., et al., Microb. Pathog. 1992; 13: 261-269; and McDaniel, L.S., et al., In: Ferretti, J.J. et al., eds. Genetics of streptococci, enterococci, and lactococci. 1995; 283-286), with the exception of clinical isolates TJ0893, 0922134 and BG8740. Pneumococcal strains TJ0893 and 0922134 were recovered from the blood of a 43-year old male and an elderly female, respectively. *S. pneumoniae* BG8743 is a blood isolate from an 8-month old infant. Strains employed in this study included capsular type 3 (A66.3, EF10197, WU2), type 4 (BG9739, EF3296, EF5668, L81905), type 5 (DBL5), type 6A (DBL6A, EF6796), type 6B (BG7322, BG9163, DBL1), type 14 TJ0893), type 19 (BG8090), and type 23 (0922134, BG8743). In addition, strain WG44.1, which expresses no detectable PspA, was employed in PspA-specific antibody analysis. All chemicals were purchased from Fisher Scientific, Fair Lawn, New Jersey unless indicate otherwise.

S. pneumoniae were grown in Todd Hewitt broth (Difco, Detroit, Michigan) supplemented with 5% yeast extract (Difco). Mid-exponential phase cultures were used for seeding inocula in Lactated Ringer's (Abbott laboratories, North Chicago, Illinois) for challenge studies. For pneumococcal strains used in challenge studies, inocula ranged from 2.8 to 3.8 log₁₀ CFU (verified by dilution plating on blood agar). Plates were incubated overnight in a candle jar at 37°C.

E. coli DH1 and BL21(DE3) were cultured in LB medium (1% Bacto-tryptone (Difco), 0.5% Bacto Yeast (Difco), 0.5% NaCl, 0.1% dextrose). For the preparation of cell lysates, recombinant *E. coli* were grown in minimal E medium supplemented with 0.05 M thiamine, 0.2% glucose, 0.1% casamino acids (Difco), and 50 mg/ml kanamycin. Permanent bacterial stocks were stored at -80°C in growth medium containing 10% glycerol.

Construction of plasmid-based strains. pET-9a

(Novagen, Madison, Wisconsin) was used for cloning truncated *pspA* genes from fourteen *S. pneumoniae* strains: DBL5, DBL6A, WU2, BG9739, EF5668, L81905, 0922134, BG8090, BG8743, BG9163, DBL1, EF3296, EF6796, and EF10197 (Table 1). *pspA* gene fragments, from fifteen strains, were amplified by PCR using two primers provided by Connaught Laboratories, Swiftwater, Pennsylvania Primer N192- 5'GGAAGGCCATATGCTCAAAGAGATTGATGAGTCT3' and primer C588 - 5'CCAAGGATCCTTAAACCCATTCACCATTGGC3' were engineered with *NdeI* and *BamHI* restriction endonuclease sites, respectively. PCR-amplified gene products were digested with *BamHI* and *NdeI*, and ligated to linearized pET-9a digested likewise and further treated with bacterial alkaline phosphatase (United States Biochemical Corporation, Cleveland, Ohio) to prevent recircularization of the cut plasmid. Clones were first established in *E. coli* BL21(DE3) which contained a chromosomal copy of the T7 RNA polymerase gene under the control of an inducible *lacUV5* promoter.

E. coli DH1 cells were transformed by the method of Hanahan (Hanahan, D. J. Mol. Biol. 1983; 166: 557-580). Stable transformants were identified by screening on LB-kanamycin plates. Plasmid constructs, isolated from each of these strains, were electroporated (Electro Cell Manipulator 600, BTX Electroporation System, San Diego, California) into *E. coli* BL21(DE3) and their respective strain designations are listed in Table 1. The pET-9a vector alone was introduced into *E. coli* BL21(DE3) by electroporation to yield strain RCT125 (Table 2). All plasmid constructs and PCR-amplified *pspA* gene fragments were evaluated by agarose gel electrophoresis (with 1 kb DNA ladder, Gibco BRL, Gaithersburg, Maryland). Next, Southern analysis was performed using LM*pspA*1, a previously described full-length *pspA* probe (McDaniel. L.S. et al., Microb. Pathog. 1992; 13: 261-269) random primed labeled with digoxigenin-11-dUTP (Genius System, Boehringer Mannheim, Indianapolis, Indiana). Hybridization was detected with chemiluminescent sheets according to the manufacturer's instructions (Schleicher & Schuell, Keene, New Hampshire).

Cell fractionation of recombinant *E. coli* strains.

Multiple cell fractions from transformed *E. coli* were evaluated for the expression of truncated PspA molecules. Single colonies were inoculated into 3 ml LB cultures containing kanamycin and grown overnight at 37°C. Next, an 80 ml LB culture, inoculated with 1:100 dilution of the overnight culture, was grown at 37°C

to mid-exponential phase (A_{600} of ca. 0.5) and a 1 ml sample was harvested and resuspended (uninduced cells) prior to induction with isopropylthiogalactoside (IPTG, 0.3 mM final concentration). Following 1, 2, and 3 hr of induction, 0.5 ml of cells were centrifuges, resuspended, and labeled induced cells. The remaining culture was divided into two aliquots, centrifuged (4000 x g, 10 min, DuPont Sorvall RC 5B Plus), and the supernatant discarded. One pellet was resuspended in 5 ml of 20 mM Tris-HCl pH 7.4 200 mM NaCl, 1 mM (ethylenedinitrilo)-tetraacetic acid disodium salt (EDTA) and frozen at -20°C overnight. Cells were thawed at 65°C for 30 min, placed on ice, and sonicated for five 10-sec pulses (0.4 relative output, Fisher Sonic Dismembrator, Dynatech Laboratories, Inc. Chantilly, Virginia). Next, the material was centrifuged (9000 x g, 20 min) and the supernatant was designated the crude extract-cytoplasmic fraction. The pellet was resuspended in Tris-NaCl-EDTA buffer and labeled the insoluble cell wall and membrane fraction. The other pellet, from the divided induced culture, was resuspended in 10 ml of 30 mM Tris-HCl pH 8.0 containing 20% sucrose and 1 mM EDTA and incubated at room temperature for 10 min with agitation. Cells were then centrifuged, the supernatant removed, and the pellet resuspended in 5 mM $MgSO_4$ (10 ml, 10 min, shaking 4°C bath). This material was centrifuged and the supernatant was designated osmotic shock-periplasmic fraction. Cell fractions were evaluated by SDS-PAGE and immunoblot analysis.

MAbs to PspA. PspA-specific monoclonal antibodies (MAbs) XiR278 and 1A4 were used as previously described (Crain, M.J. et al., 1990, Infect. Immun.; 58: 3293-3299). MAb P50-92D9 was produced by immunization with DBL5 PspA. The PspA-specificity of MAb P50-92D9 was confirmed by Western Analysis by its reactivity with native PspAs from *S. pneumoniae* DBL5, BG9739, EF5668, and L81095 and its failure to recognize the PspA-control strain WG44.1.

SDS-PAGE and immunoblot analysis. *E. coli* cell fractions containing recombinant PspA proteins and biotinylated molecular weight markers (low range, Bio-Rad, Richmond, California) were separated by sodium dodecyl sulfate-polyacrylamide (10%; Bethesda Research Laboratories, Gaithersburg, Maryland) gel electrophoresis (SDS-PAGE) by the method of Laemmli (Laemmli, U.K. Nature 1970; 227: 680-685). Samples were first boiled for 5 min in sample buffer containing 60 mM Tris pH 6.8, 1% 2-B-mercaptoethanol (Sigma, St. Louis, Missouri), 1% SDS, 10% glycerol, and 0.01% bromophenol blue. Gels were subsequently transferred (1 hr, 100 volts) to nitrocellulose (0.45 mM pores, Millipore, Bedford, Massachusetts) as per the method of Towbin et al. Blots were blocked with 3% casein, 0.05% Tween 20 in 10 mM Tris, 0.1 M NaCl, pH 7.4 for 30 min prior to incubating with PspA-specific monoclonal antibodies diluted in PBST for 1 hr at 25°C. Next, the blot was washed 3 times with PBST before incubating with alkaline phosphatase-

labeled goat anti-mouse immunoglobulin (Southern Biotechnology Associates, Inc., Birmingham, Alabama) for 1 hr at 25°C. Washes were performed as before and blots was developed with 0.5 mg/ml 5-bromo-4-chloro-3-indolyl phosphate and 0.01% nitro blue tetrazolium (Sigma) first dissolved in 150 μ l of dimethyl sulfoxide and then diluted in 1.5 M Tris-HCl pH 8.8. Dot blots were analyzed similarly. Lysate samples (2 μ l) were spotted on nitrocellulose filters (Millipore), allowed to dry, blocked, and detected as just described.

Preparation of cell lysates containing recombinant PspA proteins. Transformed *E. coli* strains RCT105, RCT113, RCT117, and RCT125 (Table 2) were grown in mid-exponential phase in minimal E medium before IPTG induction (2 mM final concentration, 2 hours, 37°C). Cultures were harvested by centrifugation (10 min at 9000 x g), resuspended in Tris-acetate pH 6.9, and frozen at -80°C overnight. Samples were thawed at 65°C for 30 min, cooled on ice, and sonicated. Next the samples were treated with 0.2 mM AEBSF (Calbiochem, La Jolla, California) at 37°C for 30 min and finally centrifuged to remove cell wall and membrane components. Dot blot analysis was performed using PspA-specific MAbs to validate the presence of recombinant, truncated PspA molecules in the lysates prior to their use as immunogens in mice. Unused lysate material was stored at -20°C until subsequent immunizations were performed.

Mouse immunization and challenge. CBA/CAHN-XID/J mice (Jackson Laboratories, Bar Harbor, Maine), 6-12 weeks old, were employed for protection studies. These mice carry a X-linked immunodeficiency that prevents them from generating antibody to polysaccharide components, thus making them extremely susceptible to pneumococcal infection. Animals were immunized subcutaneously with cell lysates from *E coli* recombinant strains RCT105, RCT113, RCT117, and RCT125 (Table 2) in complete Freund's adjuvant for primary immunizations. Secondary injections were administered in incomplete adjuvant and subsequent boosts in dH₂O. Immunized and nonimmunized mice (groups of 2 to 5 animals) were challenged with *S. pneumoniae* strains A66.3, BG7322, DBL6A, WU2, DBL5, BG9739, and L81905 intravenously (tail vein) to induce pneumococcal sepsis. Infected animals were monitored for 21 days and mice that survived the 3-week evaluation period were designated protected against death and scored as surviving 22 days for statistical analysis. Protection that resulted in extension of life was calculated as a comparison between mean number of days to death for immunized versus pooled control mice (nonimmunized and RCT125 sham-immunized; total of 6-7 animals).

Determination of PspA serum levels. Mice were bled retro-orbitally following the secondary boost and again prior to challenge. Representative mouse titers were evaluated by enzyme-linked immunosorbent assay (ELISA) using native, parental PspAs isolated from pneumococcal strains DBL5, BG9739, and L81905.

PspAs were immobilized on microtiter plates by incubating in 0.5 NaHCO₃, 0.5 M Na₂CO₃ pH 9.5 at 4°C overnight. Alkaline phosphatase-labeled goat anti-mouse immunoglobulin (Southern Biotechnology Associates, Inc.) was used to detect mouse serum antibodies. Color development was with p-nitrophenyl phosphate (Sigma, 1 mg/ml) in 0.5 M MgCl₂ pH 9.8 with 10% diethanolamine and absorbance was read at 405 nm after a 30 min incubation. Reciprocal titers were calculated as the last dilution of antibody that registered an optical density value of 0.1. Sera from individual mice within a particular immunogen group were evaluated separately and then the respective titers from four mice per group were combined to obtain titer range (Table 3).

Statistics. The one-tailed Fisher exact and two sample rank tests were used to evaluate protection against death and extension of life in the mouse model.

Cloning of truncated *pspA* genes. Using primers N192 and C588, truncated *pspA* genes from fifteen diverse pneumococcal strains representing eight different capsular types (Table 1) were amplified by PCR. Even though variability exists in *pspA* genes from different strain, this result demonstrates that sufficient conservation exists between variant *pspA* genes to allow sequence amplification in all strains examined to date. Successful *pspA* PCR-amplification extended to all capsule types evaluated.

Fourteen of the amplified *pspA* genes were cloned and three clones containing truncated PspA molecules from pneumococcal strains DBL5, BG9739, and L81905 were further studies (Table 2). To verify the constructions, plasmids from recombinant *E. coli* strains (RCT105, RCT113, RCT117, and RCT125 (Table 2) were isolated, digested with *NdeI* and *BAMHI* restriction endonucleases, and electrophoresed in 1% agarose side-by-side with the PCR products used in their respective constructions (Figure 1A). The digestion reaction was complete for pRCT105, while pRCT113 and pRCT117 digestions were incomplete (lanes 5 and 7, respectively). This gel was denatured and DNA transferred to nylon for Southern analysis. Figure 1B depicts the corresponding Southern blot probed with full-length Rx1*pspA* DNA. Lane 1 contains pRCT125, digested vector alone, which does not react with the pneumococcal DNA-specific probe, as expected. The *pspA*-specific probe hybridized with the PCT products and the digested plasmid inserts (see arrow, Figure 1B) as well as the partially undigested pRCT113 and pRCT117 (lane 5 and 7), confirming successful cloning of DBL5, BG9739, and L81905 *pspA* DNA. Constructions were similarly confirmed with the eleven additional recombinant strains containing truncated *pspA* genes from *S. pneumoniae* strains of different capsular and PspA types.

Expression of recombinant PspA in *E. coli* B121(De3).

Transformed *E. coli* strains RCT105, RCT113, RCT117, and RCT125 were cultured to mid-exponential phase prior to the addition of

IPTG to induce expression of the cloned, truncated *pspA* gene in each strain. A cell fractionation experiment was performed to identify the location of recombinant PspA proteins in transformed *E. coli* strains. Samples representing uninduced cells, included cells (1 hr, 2 hr, and 3 hr time intervals), the periplasmic fraction, the cytoplasmic fraction, and insoluble cell wall/membrane material were resolved by SDS-PAGE. Proteins were then transferred to nitrocellulose and Western analysis was performed using monoclonal antibodies specific for PspA epitopes.

Figure 2 reveals that both the cytoplasmic (lane 8) and the insoluble matter fractions (lane 9), from recombinant strain RCT 105, contain a protein of approximately 53.7 kDa that is recognized by MAb XiR278 that is not seen in the uninduced cell sample (lane 3). This protein increases in quantity in direct correlation with the length of IPTG induction (lanes 4-6; 1 hr, 2 hr, and 3 hr respectively). No truncated RCT105 PspA was found in the periplasmic fraction (lane 7), which was expected since the pET-9a vector lacks a signal sequence that would be necessary for directing proteins to the periplasm. The observed molecular weight (ca. 53.5 kDa) is larger than the predicted molecular weight for the 1.2 kb DBL5*pspA* gene product (43.6 kDa; Figure 1A, lane 4). Like full-length Rx1 PspA, the observed and predicted molecular weights for truncated PspAs do not agree precisely. In addition, immunoblot analysis was performed for recombinant *E. coli* strains RCT113, and RCT117 (using MAbs 1A4 and P50-92D,

respectively) and similar results were obtained, while no cell fractions from control strain RCT125 were recognized by MAb XiR278.

Evaluating the protective capacity of recombinant, truncated PspAs. The truncated PspA proteins from strains RCT113, RCT117, and RCT105 were expressed and analyzed for their ability to generate cross-protection against a battery of seven *S. pneumoniae* strains. Control mice (non-immunized and RCT125 sham-immunized) and recombinant PspA-immunized mice were challenged with mouse-virulent strains A66.3, BG7322, DBL6A, WU2, DBL5, BG9739, and L81905. Table 3 presents the day of death for each infected mouse.

Immunization with truncated PspA from RCT113, RCT117, and RCT105 conferred protection against death for all mice challenged with capsular type 3 strains (A66.3 and WU2 (Table 3)). The three truncated PspAs also provided significant protection against death with DBL6A, and BG7322 pneumococci (capsular types 6A and 6B, respectively). In addition, immunization with recombinant RCT113 PspA extended days to death in mice challenged with strains DBL5, BG9739, and L81905, while RCT117 PspA prolonged the lives of mice inoculated with BG9739 pneumococci (Table 3). Truncated BG9739 PspA elicited protection against all challenge strains (100%) evaluated in this study, while recombinant L81905 and DBL5 truncated PspAs conferred protection

against death with 71% and 57% of *S. pneumoniae* challenge strains, respectively.

Anti-PspA antibody titers elicited by the three immunogens vary over approximately a 10-fold range (Table 3). The lowest antibody levels were elicited by RCT105 and this truncated PspA also elicited protection against the fewest number of challenge strains. RCT113 and RCT117 elicited three and nine times as much anti-PspA antibody, respectively. As expected, no antibody to PspA was detected in nonimmunized mice nor was specific-PspA antibody measured in mice immunized with the vector-only control strain (RCT125).

In summary, immunization with RCT113 and RCT117 PspAs protected mice against fatal challenge with capsular type 3 and 6A strains and extended life for mice inoculated with type 4, 5, and 6B pneumococci. RCT105 PspA immunization protected against fatal infection with capsular type 3 and 6B strains and prolonged time to death for type 6A *S. pneumoniae* but offered not protection against type 4 and 5 strains. These data demonstrate that truncated PspAs from capsular type 4 and 5 pneumococci collectively protect mice and ergo other hosts, such as humans, against or delay death caused by each of the seven challenge strains. In general, however, more complete protection was observed against strains of capsular type 3, 6A, and 6B than against type 4 and 5 *S. pneumoniae*.

PspA has been shown to be a protection-eliciting molecule of *S. pneumoniae*. Immunization with PspA has also been shown to be cross-protective, although eliciting more complete protection against certain strains than others. Thus, it is possible that a broadly protective PspA vaccine might need to contain PspAs of more than one pneumococcal strain. The distal third of the alpha-helical region of PspA has been identified as a major protective region of PspA. Moreover, this region is presented in a very antigenic form when expressed with the intact C-terminal half of the molecule. In this Example, the ability to use truncated PspA proteins homologous to the region of Rx1 PspA extending from amino acid residue 192 to the C-terminus at residue 588 is demonstrated.

The C-terminal two-thirds of PspA was cloned from fourteen strains by PCR amplification of a gene fragment of the appropriate size (1.2 kb) which hybridized with full-length Rx1 *pspA*. Successful PCR amplification extended to all capsule types analyzed. Thus, the C-terminal two-third of PspA may be amplified from many, if not all, pneumococcal capsule types with Rx1 *pspA*-specific primers. This technique is thus applicable to the development of antigenic immunological or vaccine compositions containing multiple PspA or fragments thereof.

Of these clones, three truncated PspA proteins were expressed and evaluated in mouse immunization studies to determine their ability to cross-protect against challenge with a

variety of pneumococcal capsular types. All three recombinant PspAs elicited antibody reactive with their respective donor PspA and all three elicited protection against pneumococcal infection. Of the two truncated PspA proteins that elicited the highest antibody responses, 100% and 71% of the challenge strains were protected. RCT105 PspA, which elicited the lowest titers of PspA-specific antibody, yielded protection against 57% of *S. pneumoniae* strains evaluated. With all truncated PspAs, significant levels of protection were observed in four of the seven challenge strains. In fact, in all instances except for on (RCT105-immunized mice challenged with strain BG9739) the trend was for truncated PspA-immunization to elicit protection against pneumococcal challenge. These results demonstrate that truncated Rx1 PspA (amino acids 192-588) cross-protects mice against fatal *S. pneumoniae* WU2 challenge. More importantly, these data show that the homologous regions of diverse PspAs demonstrate comparable cross-protective abilities.

Strains of capsular type 4 and 5 were more difficult to protect against than were type 3, 6A and 6B pneumococcal strains. Serological differences in PspAs might affect cross-protection in some cases. Yet the difficulty in protecting against the type 4 and 5 strains used herein could not be explained on this basis, since the truncated PspA immunogens were cloned from the same three type 4 and 5 strains used for challenge. Both PspAs from the type 4 strains delayed death caused by one or both type 4

challenge strains but neither could prevent death caused by either type 4 pneumococcal strain. Moreover, the truncated PspA from the type 5 strain DBL5 elicited protection against death or delayed death with strains of capsular types 3, 6A and 6B but failed to protect against infection with its donor strain or either type 4 challenge strain.

There may be several reasons why the truncated PspAs from capsular type 4 and type 5 strains failed to protect against death even with their homologous donor *S. pneumoniae* strains. One possibility is that the type 4 and 5 strains chosen for study are especially virulent in the XID mouse model. XID mice fail to make antibodies to polysaccharides and are therefore extremely susceptible to pneumococcal infection with less than 100 CFU of most strains, including those of capsular type 3, 4, 5, 6A, and 6B. The increased mouse virulence of types 4 and 5 is apparent from the fact that in immunologically normal mice these strains have lower LD₅₀s and/or are more consistently fatal than strains of capsular types 3, 6A, or 6B.

Another possibility is that epitopes critical to protection-eliciting capacity with capsular type 4 and 5 strains are not present in the C-terminal two-thirds of PspA (amino acids 192-588), the truncated fragments used for immunization. The critical epitopes for these strains may be located in the N-terminal two thirds of the alpha-helical region of their PspA molecules. Finally, it is also possible that PspA may be less

exposed on some *S. pneumoniae* strains than others. Strain Rx1 PspA amino acid sequence does not contain the cell wall attachment motif LPXTGX described by Schneewind et al. found in many gram-positive bacteria. Rather, PspA has a novel anchoring mechanism that is mediated by choline interactions between pneumococcal membrane-associated lipoteichoic acid and the repeat region in the C-terminus of the molecule. Electron micrographic examination has confirmed the localization of PspA on the pneumococcal surface and PspA-specific MAb data supports the accessibility of surface-exposed PspA. However, it is not known whether *S. pneumoniae* strains differ substantially in the degree to which different PspA regions are exposed to the surrounding environment. Nor is it known if the quantity of PspA expressed on the bacterial cell surface differs widely between strains.

Table 1. *pspA* recombinant strains categorized by pneumococcal capsular type.

| Capsular Type | Parent Strains | Respective Recombinant Strains |
|---------------|-----------------|--------------------------------|
| 3 | WU2, EF10197 | RCT111, RCT137 |
| 4 | BG9739, EF5668 | RCT113, RCT115 |
| | L81905, EF3296 | RCT117, RCT133 |
| 5 | DBL5 | RCT105 |
| 6A | DBL6A, EF6796 | RCT109, RCT135 |
| 6B | BG9163, DBL1 | RCT129, RCT131 |
| 14 | TJO893 | none* |
| 19 | BG8090 | RCT121 |
| 23 | 0922134, BG8743 | RCT119, RCT123 |

* Truncated *pspA* amplified recently, not yet cloned

Table 2. Description of recombinant strains used in evaluating the protection-eliciting capacity of truncated PspAs in mice.

| Recombinant Strain | Description | Capsule Type of Parent PspA |
|-----------------------|--|--------------------------------|
| RCT 105 | BL21(DE3) <u>E. coli</u> with pET-9a:DBL5 | 5 |
| RCT 113 | BL21(DE3) <u>E. coli</u> with pET-9a:BG9739 | 4 |
| RCT 117 | BL21(DE3) <u>E. coli</u> with pET-9a:L81905 | 4 |
| RCT 125 | BL21(DE3) <u>E. coli</u> with pET-9a (vector only) | |

Table 3. Evaluation of the protection elicited by truncated *S. pneumoniae* PspA molecules in mice by days to death post-challenge*.

| Challenge Strain [capsular type] (log ₁₀ dose in CFU) | | | | | | | | |
|--|-----------------------------------|-----------------------------|---------------------------|------------------------------|-------------------------------|----------------------------|------------------------------|------------------------------|
| Immunizing recombinant PspA/ PspA donor strain | Reciprocal anti-PspA titer† | A66.3 [type 3] (2.78) | WU2 [type 3] (3.57) | DBL6A [type 6A] (3.24) | BG7322 [type 6B] (3.11) | DBL5 [type 5] (3.81) | BG9739 [type 4] (3.56) | L81905 [type 4] (3.62) |
| RCT113/BG9739 | 5590 - 50,300 | 4x >21 † | 4x >21 § | 15, 3x >21 † | 12,13,16,>21 † | 3, 3, 4, 5 § | 5, 5, 5, 7 § | 5, 6, 8, 8 † |
| RCT117/L81905 | 5590 - 150,900 | 4x >21 † | 4x >21 § | 7, 16, 2x >21 † | 10,12,13,>21 § | 3, 3, 4, 4 ¶ | 4, 5, 13, >21 § | 3, 4, 6, 8 |
| RCT105/DBL5 | 1860 - 16,770 | 4x >21 † | 4x >21 § | 8, 10, 13, 21 † | 4x >21 † | 2, 2, 2, >21 | 2, 2, 2, 4 | 4, 5, 5, 5 |
| RCT125/vector only | 20 - 620 | 3, 6, 6, >21 | 2, 3, 3, >21 | 3, 6, 6, 6 | 7, 8, 8, 14 | 2, 2, 2, 2 | 2, 2, 3, 4, 5 | 2, 3, 5, 5 |
| none | 0 | 2, 2, 2 | 2, 3 | 3, 3, 4 | 6, 7, 9 | 2, 5 | 3, 5 | 2, 5 |

* Animals surviving the 3-week evaluation period were sacrificed and days to death recorded as >21 days. For statistical analysis, P values were calculated at 22 days for these fully protected mice.

† Range of four sera per group of mice; titers measured against native donor PspAs

‡ P ≤ 0.012

§ P ≤ 0.035

¶ P ≤ 0.057

Note: One-tailed Fisher exact and two sample rank tests were used for statistical analysis.

EXAMPLE 2 - Localization of protection-eliciting epitopes and PspA of *S. pneumoniae*

This Example, the ability of PspA epitopes on two PspA fragments (amino acids 192-588 and 192-299) to elicit cross-protection against a panel of diverse pneumococci is demonstrated. Also, this Example identifies regions homologous to amino acids 192-299 of Rx1 in 15 other diverse pneumococcal strains. The DNA encoding these regions was then amplified and cloned. The recombinant PspA fragments expressed were evaluated for their ability to elicit cross-protection against a panel of virulent pneumococci.

Bacterial strains and media conditions. *S. pneumoniae* strains were grown in Todd Hewitt broth with 0.5% yeast extract (THY) (both from Difco Laboratories, Detroit, Michigan) at 37°C or on blood agar plates containing 3% sheep blood at 37°C under reduced oxygen tension. *E. coli* strains were grown in Luria-Bertani medium or minimal E medium. Bacteria were stored at -80°C in growth medium supplemented with 10% glycerol. *E. coli* were transformed by the methods of Hanahan (Hanahan, D. J. Mol. Biol. 1983; 166: 557). Ampicillin (Ap) was used at a concentration of 100 µg/ml for *E. Coli*.

Construction of pIN-III-ompA3 and pMAL-based *E. Coli* strains. Recombinant plasmids pBC100 and pBAR416 that express and secrete *pspA* fragments from *E. Coli* were constructed with

pIN-III-ompA3 as previously described (McDaniel, L.S. et al., Microb. Pathog. 1994; 17: 323).

The pMAL-p2 vector (New England Biolabs, Protein Fusion & Purification System, catalog #800) was used for cloning *pspA* gene fragments to amino acids 192-299 from strain Rx1 and from 7 other *S. pneumoniae* strains: R36A, D39, A66, BF9739, DBL5, DBL6A, and LM100. Amplification of the *pspA* gene fragments was done by the polymerase chain reaction (PCR) as described previously (McDaniel, L.S. et al., Microb. Pathog. 1994; 17: 323) using primers 5'CCGGATCCGCTCAAAGAGATTGATGAGTCTG3' [LSM4] and 5'CTGAGTCGACTGAGTTTCTGGAGCTGGAGC3' [LMS6] made with *Bam*HI and *Sal*I restriction endonuclease sites, respectively. Primers were based on the sequence of Rx1 PspA. PCR products and the pMAL vector were digested with *BAM*HI and *Sal*I, and ligated together. Clones were transformed into *E. Coli* DH5 α by the methods of Hanahan. Stable transformants were selected on LB plates containing 100 μ g/ml Ap. These clones were screened on LB plates containing 0.1 mM IPTG, 80 μ g/ml X-gal and 100 μ g/ml Ap and replica LB plates with 100 μ g/ml Ap according to the manufacturer's instructions. The strain designations for these constructs are listed in Table 6. Positive clones were evaluated for the correct *pspA* gene fragment by agarose gel electrophoresis following plasmid isolation by the methods of Birnboim and Doly (Birnboim, H.C. et al., Nucl. Acids Res. 1979, 7: 1513). Southern analysis was done as previously described using a full-

length *pspA* probe (McDaniel, L.S. et al., Microb. Pathog. 1994; 17: 323), randomly primed labeled with digoxigenin-11-dUTP (Genius System, Boehringer Mannheim, Indianapolis, Indiana) and detected by chemiluminescence.

Expression of recombinant PspA protein fragments. For induction of expression of strains BC100 and BAR416, bacteria were grown to an optical density of approximately 0.6 at 660 nm at 37°C in minimal media, and IPTG was added to a final concentration of 2 mM. The cells were incubated for an additional 2 hours at 37°C, harvested, and the periplasmic contents released by osmotic shock. For strains BAR36A, BAR39, BAR66, BAR5668, BAR9739, BARL5, BAR6A and BAR100, bacteria were grown and induced as above except LB media + 10 mM. glucose was the culture medium. Proteins from these strains were purified over an amylose resin column according to the manufacturer's instructions (New England Biolabs, Protein Fusion & Purification System, Catalog #800). Briefly, amylose resin was poured into a 10 mL column and washed with column buffer. The diluted osmotic shock extract was loaded at a flow rate of approximately 1 mL/minute. The column was then washed again with column buffer and the fusion protein eluted off the column with column buffer containing 10 mM maltose. Lysates were stored at -20°C until further use.

Characterization of truncated PspA proteins used for immunization. The truncated PspA molecules, controls and

molecular weight markers (Bio-Rad, Richmond, CA) were electrophoresed in a 10% sodium dodecyl (SDS) - polyacrylamide gel and electroblotted onto nitrocellulose. Rabbit polyclonal anti-PspA serum and rabbit antimaltose binding protein were used as the primary antibodies to probe the blots.

A direct binding ELISA procedure was used to quantitatively confirm reactivities observed by immunoblotting. For all protein extracts, osmotic shock preparations were diluted to a concentration of 3 μ g/ml in phosphate buffered saline (PBS), and 100 μ l was added to the wells of Immulon 4 microtitration plates (Dynatech Laboratories, Inc., Chantilly, VA). After blocking with 1.5% bovine serum albumin in PBS, unfractionated tissue culture supernates of individual MABs were titered in duplicated by three-fold serial dilution through seven wells and developed using an alkaline phosphatase-labeled goat anti-mouse immunoglobulin secondary antibody (Southern Biotech Associates, Birmingham, AL) and alkalinephosphatase substrate (Sigma, St. Louis, MO). The plates were read at 405 nm in a Dynatech plate reader after 25 minutes, and the 30% end point was calculated for each antibody with each preparation.

Immunization and Protection Assays. Six to nine week old CBA/CAHN-XID/J (CBA/N) mice were obtained from the Jackson Laboratory, Bar Harbor, Maine. CBA/N mice carry an X-linked immunodeficiency trait, which renders them relatively unable to respond to polysaccharide antigens, but they do respond with

normal levels of antibodies against protein antigens. Because of the absence of antibodies reactive with the phosphocholine determinant of C-polysaccharide in their serum, the mice are highly susceptible to pneumococcal infection. Mice immunized with the BC100 fragment were injected inguinally with antigen emulsified in CFA, giving an approximate dose of 3 μ g of protein per mouse. Fourteen days later the mice were boosted intraperitoneally with 3 μ g of antigen diluted in Ringer's lactate without adjuvant. Control mice were immunized following the same protocol with diluent and adjuvant, but no antigen. Mice immunized with the BAR416 fragment were injected with 0.2 ml at two sites in the sublingual area with antigen emulsified in CFA. The mice were boosted inguinally fourteen days later with antigen emulsified in IFA and were boosted a second time fourteen days later intraperitoneally with 0.2 ml of antigen diluted in Ringer's lactate without adjuvant.

Mice that were immunized with the homologues of Rx1 BAR416 were immunized as described above. The control animals followed the same immunization protocol but received maltose binding protein (MBP) diluted 1:1 in CFA for their immunization and were also boosted with MBP.

Serum analysis. Mice were retro-orbitally bled with a 75 μ l heparinized microhematocrit capillary tube (Fisher Scientific) before the first immunization and then once approximately 2 hours before challenge with virulent pneumococci.

The serum was analyzed for the presence of antibodies to PspA by an enzyme-linked immunosorbent assay (ELISA) using native full-length R36A PspA as coating antigen as previously described (McDaniel, L.S. Microb. Pathog. 1994; 17: 323).

Intravenous infection of mice. Pneumococcal cultures were grown to late log phase in THY. Pneumococci were diluted to 10^4 CFU based on the optical density at 420 nm into lactated Ringer's solution. Seven days following the last boost injection for each group, diluted pneumococci were injected intravenously (tail vein) in a volume of 0.2 ml and plated on blood agar plates to confirm the numbers of CFU per milliliter. The final challenge dose was approximately 50-100 times the LD₅₀ of each pneumococcal strain listed in Tables 4-6. The survival of the mice was followed for 21 days. Animals remaining alive after 21 days were considered to have survived the challenge.

Statistical analysis. Statistical significance of differences in days to death was calculated with the Wilcoxon two-sample rank test. Statistical significance of survival versus death was made using the Fisher exact test. In each case, groups of mice immunized with PspA containing preparations were compared to unimmunized controls, or controls immunized with preparations lacking PspA. One-tailed, rather than two-tailed, calculations were used since immunization with PspA or fragments of PspA has never been observed to cause a statistically significant decrease in resistance to infection.

Cloning into pMAL vector. Using primers based on the sequence of Rx1 PspA, LSM4 and LSM6, *pspA* gene fragments were amplified by PCR from fifteen out of fifteen pneumococcal strains examined. Seven of the eleven gene fragments were cloned into pMAL-p2 and transformed into *E. coli* (Table 6). The correct insert for each new clone was verified by agarose gel electrophoresis and Southern hybridization analysis. Plasmids from recombinant *E. coli* strains BAR36A, BAR39, BAR66, BAR9739, BARL5, BAR6A and BAR100 were isolated, digested with *Bam*HI and *Sal*I restriction endonucleases and electrophoresed on a 0.7% TBE agarose gel. The gel was then denatured and the DNA transferred to a nylon membrane for southern hybridization. The blot was probed with full-length Rx1 *pspA* DNA at high stringency conditions. The cloning of R36A, D39, A66, BG9739, DBL5, DBL6A and LM100 *pspA* DNA into pMal-p2 was confirmed by the recognition of all *Bam*HI and *Sal*I digested DNA inserts by the Rx1 probe.

Expression and conformation of truncated recombinant proteins. The transformed *E. coli* strains BAR36A, BAR39, BAR66, BAR9739, BARL5, BAR6A and BAR100 were grown in LB media supplemented with 10 mM glucose and induced with 2 mM IPTG for expression of the truncated PspA protein fused with maltose binding protein. Transformed *E. coli* strains BC100 and BAR416, which express PspA fragments fused to the OmpA leader sequence in the pIN-III-ompA3 vector, were grown in minimal medium and induced with 2 mM IPTG for expression. Both vectors, pIN-III-

ompA3 and pMal-p2, are vectors in which fusion proteins are exported to the periplasmic space. Therefore, an osmotic shock extract from the pMal-p2 containing bacteria was then run over an amylose column for purification and resolved by SDS-PAGE western blotting. The western blot of the protein extracts from BAR36A, BAR39, BAR66, BAR9739, BARL5, BAR6A and BAR100 were recognized by a rabbit polyclonal antibody made to strain BC100 PspA. The apparent M_r of full-length PspA from WU2 is 91.5 kD. The M_r of maltose binding protein is 42 kD and the expected M^r for the PspA portion of the fusion is 12 kD. All extracts exhibited molecular weights that ranged from 54 to 80 kD. This range of molecular weights can be attributed to the variability of *pspA* among different pneumococcal strains. An ELISA, with plates coated with the various cloned fragments quantitatively confirmed the reactivities that were observed in the western blots with all protein extracts.

Protection and cross-protection against fatal pneumococcal infection elicited by cloned PspA fragments. CBA/N mice were immunized with the truncated PspA fragment encoded by pBC100, which is composed of amino acids 192 to 588 of Rx1 PspA, and challenged with 13 different *S. pneumoniae* strains representing 7 different capsular types (Table 4). With all 13 strains, the immunization resulted in protection from death or an extended time to death. With 10 of the strains the difference was statistically significant. With strains of capsular types 3,

6A, and 6B, all immunized mice were protected against death. Although there were fewer survivors in the case of capsular types 2, 4, and 5, the immunization with BC100 resulted in significant increases in times to death.

The BC100 immunization studies made it clear that epitopes C-terminal to residue 192 could elicit cross-protection. The BAR416 fragment, which includes amino acids 192-299, could elicit protection from fatal infection with a single challenge strain WU2. This Example shows the ability of BAR416 immunization to protect against the 6 strains that had been best protected against by immunization with BC100. Immunization with the BAR416 construct resulted in increased time to death for all 6 challenge strains examined (Table 5). BAR416 provided significant protection against death with WU2, A66, BG7322 and EF6796 pneumococci (capsular types 3, 3, 6B and 6A respectively). It also prolonged the lives of mice challenged with ATCC6303 and DBL6A pneumococci (capsular types 3 and 6A respectively). Serum from mice immunized with the BAR416 fragment yielded a geometric mean reciprocal anti-PspA ELISA titer to full-length Rx1 PspA of 750. Mice immunized with BC100 had geometric mean reciprocal titers of close to 2000, while non-immunized mice had anti-PspA titers of <10.

The above data indicates that the BAR416 fragment from Rx1 elicits adequate cross-reactive immunity to protect against diverse pneumococci and suggests that this region must be

serologically conserved among PspAs. This hypothesis was confirmed by immunized with recombinant BAR416 homologous regions from the 7 different clones and then challenging with strain WU2 (Table 6). All 7 immunogens elicited significant protection. PspA fragments from capsular types 2 and 22 and the rough R36A strain elicited complete protection against death with all challenged mice. All of the other immunogens were able to extend the time to death of all the mice with the median days to death being 21 days or >21 days. Serum from mice immunized with the BAR416 homologous fragments had anti-PspA reciprocal titers that ranged from 260 to 75,800 with an average of 5700 while control animals immunized with only maltose binding protein had anti-PspA reciprocal titers of <10.

Antibody reactivities. All of the above immunization studies attest to the cross-reactivity of epitopes encoded by amino acids from position 192-299. This region includes the C-terminal third of the α -helical region and the first amino acids of the proline rich region. Other evidence that epitopes within this region are cross-reactive among different PspAs comes from the cross-reactivity of a panel of nine MAbs all of which were made by immunization with Rx1 PspA. The epitopes of four of the antibodies in the panel reacted with epitopes mapping between amino acids 192-260. The epitopes of the other five MAbs in the panel map between amino acids 1 and 115 (McDaniel, L.S., et al., Microb. Pathog. 1994; 17: 323). Each of these 9 MAbs were tested

for its ability to react with 8 different PspAs in addition to Rx1. The 5 MAbs whose epitopes were located within the first 115 amino acids, reacted on average with only 1 other PspA. Three of the 5 in fact, did not react with any of the other 8 PspAs. In contrast the MAbs whose epitopes map between 192 and 260 amino acids each cross-reacted with an average of 4 of the 8 non-Rx1 PspAs, and all of them reacted with at least two non-Rx1 PspAs. Thus, based on this limited section of individual epitopes, it would appear that epitopes in the region from 192-260 amino acids are generally much more cross-reactive than epitopes in the region from 1-115 amino acids.

The BC100 fragment of Rx1 PspA can elicit protection against the encapsulated type 3 strain WU2. Although the PspAs of the two strains can be distinguished serologically they are also cross-reactive (Crain, M.J., et al., Infect. Immun. 1990; 58: 3293). The earlier finding made it clear that epitopes cross-protective between Rx1 and WU2 PspAs exist. The importance of cross-reactions in the region C-terminal to residue 192 is demonstrated in this Example where 13 mouse virulent challenge strains have been used to elicit detectable protection against all of them. The first indication that epitopes C-terminal to position 192 might be able to elicit cross-protection came from our earlier study where we showed the MAbs Xi64, XiR278, XiR1323, and XiR1325, whose epitopes mapped between amino acids 192 and 260 of strain Rx1 PspA, could protect against infection with WU2.

Moreover, immunization with PspA fragments from 192-588 and 192-299 were able to elicit protection against infection against WU2. This Example shows that the BC100 Rx1 fragment (192-588) elicits significant protection against each of 13 different mouse virulent pneumococci, thereby firmly establishing the ability of epitopes C-terminal to position 192 to elicit a protective response. The observation that a fusion protein containing amino acids 192-299 fused C-terminal to maltose binding protein could also elicit cross-protection, permits the conclusion that epitopes in this 107 amino acid region of PspA are sufficient to elicit significant cross-protection against a number of different strains.

Evidence that a comparable region of other PspAs is also able to elicit cross-protection came from the studies where sequences homologous to the 192-299 region of Rx1 PspA were made from 5 other PspAs. All 5 of these fragments elicited significant protection against challenge with strain WU2. These data provide some suggestion for serologic differences in cross-protection elicited by the 192-299 region.

Based on present evidence, without wishing to be bound by any one particular theory, it is submitted that the PspAs in strains D39, Rx1 and R36A are identical. All of the 9 mice immunized with the 192-299 fragments from R36A and D39 survived challenge with WU2. Only LM100, one of the non-R36A/D39 PspAs, protected as high a percentage of mice from WU2. The difference

in survival elicited by the R36A/D39 PspAs and the non-Rx1 related PspAs was statistically significant.

The data does indicate however, that all of the differences in protection against different strains are not due to differences in serologic cross-reactivity. BC100, which is made from Rx1, protected against death in 100% of the mice challenged with 7 different strains of *S. pneumonia*, but only delayed death with strain D39, which is thought to have the same PspA as strain Rx1. Thus, some of the differences in cross-protection are undoubtedly related to factors other than PspA cross-reactivity. Whether such factors are related to differences in virulence of the different strains in the hypersusceptible Xid mouse, or differences in requirements for epitopes N-terminal to amino acid 192, or difference in the role of PspA in different strains is not yet known.

These results suggest that a vaccine containing only the recombinant PspA fragments homologous with Rx1 amino acids 192-299 is effective against pneumococcal infection. Moreover, the results demonstrate that utility of PspA a.a. 192-299, a.a. 192-260 and DNA coding therefor, e.g. primers N192 or 588 (variants of LSM4 and LSM2) as useful for detecting the presence of pneumococci by detecting presence of that which binds to the amino acid or to the DNA, or which is amplified by the DNA, e.g., by using that DNA as a hybridization probe, or as a PCR primer, or by using the amino acids in antibody-binding kits, assays or

tests; and, the results demonstrate that a.a. 192-299 and a.a. 192-260 can be used to elicit antibodies for use in antibody-binding kits assays or tests.

Table 4 Protection of mice by immunization with BC100 from Rx1 PspA

| Challenge Strain* | Capsule type | PspA type | BC100 Immunogen | | | Controls | | | P Value [§] |
|-------------------|--------------|-----------|------------------|------------|-------------------|------------------|------------|-------------------|----------------------|
| | | | # alive / # dead | % Survival | Median days alive | # alive / # dead | % Survival | Median days alive | |
| D39 | 2 | 25 | 0 / 5 | 0% | 5 | 0 / 3 | 0% | 2 | 0.02 |
| WU2 | 3 | 1 | 4 / 0 | 100% | >21 | 0 / 3 | 0% | 3 | 0.002 |
| ATCC6303 | 3 | 7 | 5 / 0 | 100% | >21 | 0 / 5 | 0% | 7 | 0.004 |
| A66 | 3 | 13 | 4 / 0 | 100% | >21 | 0 / 3 | 0% | 1 | 0.03 |
| EF10197 | 3 | 18 | 5 / 0 | 100% | >21 | 0 / 3 | 0% | 2 | 0.02 |
| EF5668 | 4 | 12 | 1 / 3 | 25% | 9 | 0 / 3 | 0% | 4 | N.S. |
| EF3296 | 4 | 20 | 1 / 3 | 25% | 5 | 0 / 3 | 0% | 3 | N.S. |
| L81905 | 4 | 23 | 1 / 4 | 20% | 4 | 0 / 6 | 0% | 2 | 0.02 |
| BG9739 | 4 | 26 | 0 / 4 | 0% | 6.5 | 0 / 3 | 0% | 2 | N.S. |
| DBL5 | 5 | 33 | 0 / 5 | 0% | 5 | 0 / 3 | 0% | 2 | 0.02 |
| BG7322 | 6 | 24 | 4 / 0 | 100% | >21 | 1 / 2 | 33.3% | 6 | 0.03 |
| EF6796 | 6A | 1 | 4 / 0 | 100% | >21 | 0 / 3 | 0% | 1 | 0.03 |
| DBL6A | 6A | 19 | 5 / 0 | 100% | >21 | 0 / 3 | 0% | 7 | 0.03 |

* Mice were challenged with approximately 10^5 CFU/mL of each strain
[§] P values were based on comparison of days alive by a one-tailed Wilcoxon 2 sample-rank test

Table 5 Protection of mice by immunization with BAR416 from Rx1 PspA

| Challenge Strain | Capsule type | PspA type | BAR416 Immunogen | | | Controls | | | P Value [§] |
|------------------|--------------|-----------|------------------|------------|-------------------|------------------|------------|-------------------|----------------------|
| | | | # alive / # dead | % Survival | Median days alive | # alive / # dead | % Survival | Median days alive | |
| WU2 | 3 | 1 | 4 / 1 | 80% | >21 | 0 / 3 | 0% | 1 | 0.002 |
| ATCC6303 | 3 | 7 | 2 / 3 | 40% | 13 | 1 / 4 | 20% | 4 | 0.048 |
| A66 | 3 | 13 | 5 / 0 | 100% | >21 | 0 / 5 | 0% | 2 | 0.004 |
| BG7322 | 6 | 24 | 3 / 2 | 60% | >21 | 0 / 4 | 0% | 7 | 0.02 |
| EF6796 | 6A | 1 | 3 / 2 | 60% | >21 | 0 / 5 | 0% | 5 | 0.004 |
| DBL6A | 6A | 19 | 0 / 5 | 0% | 7 | 0 / 5 | 0% | 2 | 0.008 |

Note, mice were challenged with about 10^5 CFU of each strain
[§] P values were based on comparison of days alive by a one-tailed Wilcoxon 2 sample-rank test

Table 6 Protection of mice against *S. pneumoniae* WU2 by immunization with BAR416 Analogs of 7 PspAs

| Immunogen | Parent Strain | Capsule type | PspA type | # alive / total # | % Survival | Median days alive | P value* vs. MBP |
|-----------|---------------|--------------|-----------|-------------------|------------|-------------------|------------------|
| BAR36A | R36A | - | 25 | 4 / 4 | 100 % | >21 | 0.002 |
| BAR39 | D39 | 2 | 25 | 5 / 5 | 100 % | >21 | 0.0008 |
| BAR66 | A66 | 3 | 13 | 7 / 8 | 88 % | >21 | <0.0001 |
| BAR9739 | BG9739 | 4 | 26 | 5 / 8 | 63 % | >21 | 0.0002 |
| BARL5 | DBL5 | 5 | 33 | 4 / 8 | 50 % | 21 | 0.03 |
| BAR6A | DBL6A | 6A | 19 | 3 / 5 | 60 % | >21 | 0.05 |
| BAR100 | LM100 | 22 | ND | 5 / 5 | 100 % | >21 | 0.0008 |
| MBP | - | - | - | 0 / 8 | 0 % | 2 | - |

* P values were based on comparison of days alive by a one-tailed Wilcoxon 2 sample-rank test

Note, the PspA fragments used for immunization were cloned from products amplified with primers LSM4 and LSM6. In addition to the strains listed above, PCR reactions with LSM4 and LSM6 amplified products of the appropriate size from strains BG9163, WU2, L81905, EF6796, EF5668, BG7376, BG7322, and BG5-8A.

Table 7 Reactivity of MAbs with PspAs of Different Pneumococci

| Donor of test PspA | | | MAb mapping to 1-115 amino acids | | | | | MAb mapping to 192-260 amino acids | | | |
|--------------------|--------------|-----------|----------------------------------|-------------|---------------|-------------|-------------|------------------------------------|----------|-------------|---------------|
| Strain | Capsule Type | PspA Type | Xi126 IgG2b | XiR1224 IgM | XiR1526 IgG2b | XiR35 IgG2a | XiR16 IgG2a | XiR1323 IgM | Xi64 IgM | XiR278 IgG1 | XiR1325 IgG2a |
| Rx1 | rough | 25 | ++ | ++ | ++ | ++ | ++ | ++ | ++ | ++ | ++ |
| ATCC101813 | 3 | 3 | ++ | - | - | - | - | ++ | ++ | ++ | ++ |
| EF10197 | 3 | 18 | - | - | - | - | - | - | - | ++ | +/- |
| BG9739 | 4 | 26 | - | - | - | - | - | ++ | - | + | ++ |
| L81905 | 4 | 23 | - | - | - | - | - | - | - | - | - |
| BG-5-8A | 6A | 0 | +/- | + | - | - | - | + | - | + | - |
| BG9163 | 6B | 21 | - | - | - | - | - | - | - | + | - |
| LM100 | 22 | N.D. | +/- | - | - | - | - | - | - | - | - |
| WU2 | 3 | 1 | ++ | - | - | - | - | ++ | ++ | ++ | ++ |

Note, immunoblot analysis was carried out with the nine MAbs from this study against a panel of nine different pneumococcal strains. Rx1 served as a positive control. The results are presented as ++ (strong reaction), + (weak, but clearly positive reaction), +/- (difficult to detect), and - (no reaction). The PspA of all strains gave a positive reaction with rabbit

antiserum against PspA. N.D. means not determined. Mapping of epitopes was to fragments of strain Rx1 PspA

**EXAMPLE 3 - Isolation of PspA and Truncated Forms
Thereof, and Immunization Thereby**

PspA is attached to the pneumococcal surface through a choline binding site on PspA. This allows for successful procedures for the isolation of FL-PspA. PspA can be released from the surface of pneumococci by elution with 2 percent choline chloride (CC), or by growth in a chemically defined medium (CDM) containing 1.2 percent CC (CDM-CC) or medium in which the choline had been replaced by ethanolamine (CDM-ET). Since CDM-ET supernatants lack high concentrations of choline, the PspA released into them can be adsorbed to a choline (or choline analog) column and isolated by elution from the column with 2 percent choline chloride (CC).

This Example describes the ability to obtain PspA by these procedures, and the ability of PspA obtained by these procedures to elicit protection in mice against otherwise fatal pneumococcal sepsis. Native PspA from strains R36A, Rx1, and WU2 was used because these strains have been used previously in studies of the ability of PspA to elicit protective immunity (see, e.g., Examples *infra* and *supra*). The first MAbs to PspA were made against PspA from strain R36A and the first cloned fragments of PspA and PspA mutants came from strain Rx1. Strain Rx1 was derived from strain R36A, which was in turn derived from the encapsulated type 2 strain, D39. PspAs from these three strains appears to be identical based on serologic and molecular weight analysis. Molecular studies have shown no differences in the *pspA* genes of strains D39, Rx1, and R36A. The third strain that provided PspA in this Example is the mouse virulent capsular type 3 strain WU2. Its PspA is highly cross-reactive with that from R36A and Rx1, and immunization with Rx1 and D39 PspA can protect against otherwise fatal infections with strain WU2.

S. pneumoniae

Strains of *S. pneumoniae* used in this study have been described previously (Table 6). Bacteria were grown in either Todd-Hewitt broth with 0.5 percent yeast extract (THY), or a chemically defined medium (CDM) described previously^{32, 43}. Serial passage of stock cultures was avoided. Strains were maintained frozen in THY + 20 percent glycerol and cultured from a scraping of the frozen culture.

Recovery of PspA from pneumococci

PspA is not found in the medium of growing pneumococci unless they have reached stationary phase and autolysis has commenced ³⁶. To release PspA from pneumococci three procedures were used. In one approach were grow pneumococci in 100 ml of THY and collect the cells by centrifugation at mid-log phase. The pellet was washed three times in lactated Ringer's solution (Abbot Lab. North Chicago, IL), suspended in a small volume of 2 percent choline chloride in phosphate buffered saline (PBS) (pH 7.0), incubated for 10 minutes at room temperature, and centrifuged to remove the whole pneumococci. From immunoblots with anti-PspA MAb Xi126 ⁴⁸ at serial dilutions of the original culture, the suspended pellet, and the supernatant, it was evident that this procedure released about half of the PspA originally present on the pneumococci. Analysis of silver stained polyacrylamide gels showed this supernatant to contain proteins in addition to PspA ³⁶.

The CDM used in the remaining two procedures was modified from that of Van der Rijn ⁴³. For normal growth it contained 0.03% CC. To cause PspA to be released during bacterial growth, the pneumococci were grown in CDM containing 1.2 percent choline chloride (CDM-CC), or in CDM containing 0.03 percent ethanolamine and only 0.000,001 percent choline (CDM-ET). In media lacking a normal concentration of choline the F-antigen and C-polysaccharide contain phosphoethanolamine rather than phosphocholine ⁴⁹. In CDM-CC and CDM-ET, PspA is released from the pneumococcal surface because of its inability to bind to the cholines in the lipoteichoic acids ³⁶. In addition to releasing PspA from the pneumococcal surface, growth in CDM-CC or CDM-ET facilitates PspA isolation by its other effects on the cell wall. In these media pneumococci do not autolyse ⁴⁹, thus permitting them to be grown into stationary phase to maximize the yield of PspA. In these media septation does not occur and the pneumococci grow in long chains ^{36, 49}. As the pneumococci reach stationary phase they die, cease making PspA, and rapidly settle out. Preliminary studies, using serial dilution dot blots to quantitate PspA, indicated that the production of PspA ceases at about the time the pneumococci begin to settle out, with the formation of visible strands of the condensed pneumococcal chains. When the pneumococci began to settle out, the medium was recovered by centrifugation at 2900 x g for 20 minutes, and filtered with a low protein-binding filter (.45µ Nalgene Tissue Culture Filter #158-0045).

For growth in CDM-CC or CDM-ET, the pneumococci were first adapted to the defined medium and then, in the case of CDM-ET, to very low choline concentrations. To do this, strains were first inoculated into 1 part of THY and 9 parts of CDM medium containing 0.03 percent choline and 0.03 percent ethanolamine. After two subsequent subcultures in CDM containing 0.03 percent choline and 0.03 percent ethanolamine (0.1 ml of culture + 0.9 ml of pre-warmed fresh medium), the culture was used to inoculate CDM with only 0.003 percent choline (and 0.03 percent ethanolamine). These steps was repeated until the strain would grow in CDM-ET containing 0.000,001 percent choline and 0.03 percent ethanolamine. It was critical that cultures be passed while in exponential growth phase (at about 10⁷ CFU/ml). Even trace contamination of the medium by

exogenous choline resulted in the failure of the PspA to be released from the pneumococcal surface³⁶. Thus, disposable plastic ware was used for the preparation of CDM-ET media and for growth of cultures. Once a strain was adapted to CDM-ET it was frozen in 80 percent CDM-ET and 20 percent glycerol at -80° C. When grown subsequently the strain was inoculated directly into the CDM-ET.

Isolation of native (full-length) PspA

PspA was isolated from the medium of cells grown in CDM-ET using choline-Sepharose prepared by conjugating choline to epoxy-activated Sepharose⁵⁰. A separate column was used for media from different strains to avoid cross-contamination of their different PspAs. For isolation of PspA from clarified CDM-ET, we used a 0.6 ml bed volume of choline-Sepharose. The column bed was about 0.5 cm high and 1.4 cm in diameter. The flow rate during loading and washing was approximately 3 ml/min. After loading 300 ml CDM-ET supernatant, the column was washed 10 times with 3 ml volumes of 50 mM Tris acetate buffer, pH 6.9 containing 0.25 M NaCl (TAB). The washed column was eluted with sequential 3 ml volumes of 2 percent CC in TAB. Protein eluted from the column was measured (Bio-Rad protein assay, Bio-Rad, Hercules, CA). The column was monitored by quantitative dot blot. The loading material, washes, and the eluted material were dot blotted (1 µl) as undiluted, 1/4, 1/16, 1/64, 1/256, and 1/1024 on nitrocellulose. The membranes were then blocked with 1 percent BSA in PBS, incubated for 1 hr with PspA-specific MAbs Xi126 or XiR278, and developed with biotinylated goat-anti-mouse Ig, alkaline phosphatase conjugated streptavidin (Southern Biotechnology Associates Inc. Birmingham, AL), and nitrobluetetrazolium substrate with 5-bromo 4-chloro-3-indoyl phosphate *p*-toluidine salt (Fisher Scientific, Norcross GA)¹⁷. The purity of eluted PspA was assessed by silver-stained (silver stain kit, Bio Rad, Hercules, CA) SDS-PAGE gels run as described previously³². Immunoblots of SDS-PAGE gels were developed with MAbs Xi126 and XiR278¹⁷.

Isolation of 29 kDa PspA

The 29 kDa fragment comprising the N-terminal 260 amino acids of PspA was produced in DH1 *E. coli* from pJY4306^{31, 37}. An overnight culture of JY4306 was grown in 100 ml of Luria Broth (LB) containing 50 µg/ml ampicillin. The culture was grown at 37° C in a shaker at 225 rpm. This culture was used to inoculate 6 one liter cultures that were grown under the same conditions. When the culture O.D. at 600 nm reached 0.7, 12 grams of cells, as a wet paste, were harvested at 4° C at 12,000 xg. The pellet was washed in 10 volumes of

25 mM Tris pH 7.7 at 0° C and suspended in 600 ml of 20% sucrose, 25 mM Tris pH 7.7 with 10 mM ethylenediamine tetraacetic acid (EDTA) for 10 minutes. The cells were pelleted by centrifugation (8000 xg) and rapidly suspended in 900 ml of 1 percent sucrose with 1 mM Pefabloc SC hydrochloride (Boehringer Mannheim Corp., Indianapolis, IN.) at 0° C. The suspension was pelleted at 8000 xg at 4° C for 15 minutes and the PspA-containing supernatant (periplasmic extract) ⁵¹ recovered. The recombinant PspA was precipitated from the periplasmic extract by 70 percent saturated ammonium sulfate overnight at 4° C. The precipitated material was collected by centrifugation at 12,000 xg at 4° C for 30 minutes. The precipitated protein was resuspended in 35 ml of 20 mM histidine 1 percent sucrose at pH 6.6 (HSB). Insoluble materials were removed at 1,000 xg at 4° C for 10 minutes. The clarified material was dialyzed versus HSB, passed through a 0.2µm filter and further purified on a 1 ml MonoQ HR 5/5 column (Pharmacia Biotech, Inc., Piscataway, N.J) equilibrated with HSB. The clarified material was loaded on the column at 1 ml/min, and the column was washed with 10 column volumes of HSB. The column was then eluted with a gradient change to 5 mM NaCl per minute at a flow rate of 1 ml/min. As detected by immuno blot with Xi126, SDS-PAGE and absorbance, PspA eluted as a single peak at approximately 0.27 to 0.30 M NaCl. By SDS-PAGE the material was approximately 90 percent pure. The yield from 6 liters of culture was 2 mg (Bio-Rad protein assay) of recombinant PspA.

Growth of pneumococci for challenge

Mice were challenged with log-phase pneumococci grown in THY. For challenge, the pneumococci were diluted directly into lactated Ringer's without prior washing or centrifugation. To inject the desired numbers of pneumococci, their concentration in lactated Ringer's solution was adjusted to an O.D. of about 0.2 at 420 nm (LKB Ultrospec III spectrophotometer). The number of pneumococci present was calculated at 5×10^8 CFU per ml / O. D. and confirmed by colony counts (on blood agar) of serial dilutions of the inoculum.

Immunization, challenge, and bleeding of mice

CBA/CAHN/XID/J (CBA/N) and BALB/cByJ (BALB/c) mice were purchased from Jackson Laboratory Bar Harbor, ME. Mice were given two injections two weeks apart and challenged i.v. two weeks later. Injections without CFA were given intraperitoneally in a 0.1 ml of Ringers. Where indicated, the first injection was given in complete Freund's adjuvant (CFA) consisting of approximately a 1:1 emulsion of antigen solution and CFA oil (Difco, Detroit MI). Antigen in CFA was injected inguinally in 0.2 ml divided between the two hind legs. All mice were boosted i.p. without adjuvant. When mice were injected with media supernatants or 2 percent choline chloride eluates of whole bacteria, the amounts of material injected were expressed as the

volume of media from which the injected material was derived. For example, if the clarified medium from pneumococci grown in CDM-CC or CDM-ET was used for immunization without dilution or concentration, the dose was described as 100 μ l. If the material was first diluted 1/10, or concentrated 10 fold, the dose was referred to as 10 or 1000 μ l respectively.

ELISA for antibodies to PspA

Specific modifications of previously reported ELISA conditions¹⁷, are described. Microtitration plates (Nunc Maxisorp, P.G.C. Scientific, Gaithersburg MD.) were coated with undiluted supernatants of Rx1 and WG44.1 pneumococci grown in CDM-ET or 1 percent BSA in PBS. Mice were bled retro-orbitally (75 μ l) in a heparanized capillary tube (Fisher Scientific, Fair Lawn, N.J.) The blood was immediately diluted in 0.5 ml of one percent bovine serum albumin in PBS. The dilution of the resultant sera was 1/15 based on an average hematocrit of 47 percent. The sera were diluted in 7 three fold dilution in microtitration wells starting at 1/45. Mab Xi126 was used as a positive control. The maximum reproducible O. D. observed with Xi126 was defined as "maximum O. D." The O. D. observed in the absence of immune sera or MAb was defined as "minimum O.D." Antibody titers were defined as the dilution that gives 33 percent of maximum O. D. The binding to the Rx1 CDM-ET coated plates was shown to be PspA-specific, since in no case did we observe ≥ 33 percent of maximum binding of immune sera or Xi126 on plates coated with WG44.1 CDM-ET or BSA.

Statistical analysis. Unless otherwise indicated *P* values refer to comparisons using the Wilcoxin two-sample rank test to compare the numbers of days to death in different groups. Mice alive at 21 days were assigned a value of 22 for the sake of calculation. *P* values of >0.05 have been regarded as not significant. Since we have never observed immunization with PspA or other antigens to make pneumococci more susceptible to infection the *P* values have been calculated as single tailed tests. To determine what the *P* value would have been if a two tailed test had been used the values given should be multiplied by two. In some cases *P* values were given for comparisons of alive versus dead. These were always calculated using the Fisher exact test. All statistical calculations were carried out on a Macintosh computer using InStat (San Diego, CA).

PspA is the major protection-eliciting component released from pneumococci grown in CDM-ET or CDM-CC, or released from conventionally grown pneumococci by elution with 2% CC.

PspA-containing preparations from pneumococci were able to protect mice from fatal sepsis following i.v. challenge with 3×10^3 (100 x LD₅₀) capsular type 3 *S. pneumoniae* (Table 9). Comparable preparations from the strains unable to make PspA (WG44.1 and JY1119), or unable to make full length PspA (LM34 and JY2141) were unable to elicit protection. Regardless of the method of isolation the minimum protective dose was derived from pneumococci grown in from 10 - 30 μ l of medium. We also observed that supernatants of log phase pneumococci grown in normal THY or CDM media could not elicit protection (data not shown). This finding is consistent with earlier studies^{36, 37} indicating the PspA is not normally released in quantity into the medium of growing pneumococci.

Isolated PspA can elicit protection against fatal infection

Although PspA was necessary for these preparations to elicit protection it was possible that it did not act alone. Mice were thus, immunized with purified FL-PspA to address this question.

Isolation of FL-PspA from CDM-ET growth medium. We isolated the FL-PspA from CDM-ET rather than from CDM-CC medium or a 2 percent choline chloride elution of live cells, because the high levels of choline present in the latter solutions prevents adsorption of the PspA to the choline residues on the choline-Sepharose column. PspA for immunization was isolated from strain R36A, as the strain is non-encapsulated and the isolated PspA could not be contaminated with capsular polysaccharide. As a control we have conducted mock isolations from WG44.1 since this strain has an inactivated *pspA* gene and produces no PspA. The results shown in Table 10 are typical of isolations from 300 ml of CDM-ET medium from R36A grown pneumococci. We isolated 84 μ g of PspA from 300 ml of medium, or about 280 μ g/liter. Based on the dot blot results this appears to be about 75% of the PspA in the original medium, and that CDM-ET from R36A cultures contains about 400 μ g/liter of PspA, or about 0.4 μ g/ml.

No serologically detectable PspA was seen in the CDM-ET from WG44.1 cultures. More significantly there was undetectable protein recovered from the choline-Sepharose column after adsorption of CDM-ET from a WG44.1 culture, indicating that PspA is the only protein that could be isolated by this procedure. Moreover by silver stained SDS PAGE gel the PspA isolated from R36A appeared to be homogenous (Figure 3). Although autolysin can also be isolated on choline-Sepharose^{20, 50}, we did not expect it to be isolated by this procedure since autolysin is not released from pneumococci grown in choline deficient medium³⁶. The

immunologic purity of the isolated PspA was emphasized by the fact that immunization with it did not elicit any antibodies detectable on plates coated with CDM-ET supernatants of WG44.1.

Loading more than 300 ml on the 0.6 ml bed volume column did not result in an increased yield, which suggested that the column capacity had been reached. However, increasing the depth of the choline-Sepharose bed to greater than 0.5 cm, decreased the amount of PspA eluted from the column, presumably because of non-specific trapping of aggregates in the column matrix. The elution buffer contains 50 mM Tris acetate 0.25 M NaCl and 2% choline chloride. Elution without added NaCl or with 1M NaCl resulted in lower yields. Elution with less than 1% CC also reduced yields.

Immunization of mice with purified R36A PspA. For immunization we used only the first 3 ml fraction of the R36A column. Mice were immunized with two injections of 1, 0.1, or 0.01 μ g of R36A PspA, spaced two weeks apart. As controls, some mice were inoculated with a comparable dilutions of the first 3 ml fraction from the WG44.1 column. Purified FL-PspA elicited antibody to PspA at all doses regardless of whether CFA was used as an adjuvant (*Table 1*). In the absence of CFA the highest levels of antibody were seen with the 1 μ g dose of PspA. In the presence of CFA, however, the 0.1 μ g dose was as immunogenic as the 1 μ g dose.

To test the ability of the different doses of PspA to elicit protection against challenge we infected the immunized mice with two capsular type 3 strains, WU2 and A66. Although both of these strains are able to kill highly susceptible CBA/N XID mice at challenge doses of less than 10^2 , the A66 strain is several logs more virulent when BALB/c mice are used ^{47, 52}. The difference in virulence of A66 and WU2, was partially compensated for by challenging the immunized CBA/N mice with lower doses of strain A66 than WU2.

After immunization of CBA/N mice with 1 and 0.1 μ g doses of PspA we observed protection against WU2 challenge regardless of whether or not CFA was used as an adjuvant (*Table 4*). At the lowest dose, 0.01 μ g PspA, most of the mice immunized with PspA + CFA lived whereas most immunized with PspA alone did not; however, the difference was not statistically significant. When immunized mice were challenged with the more virulent strain A66 ^{47, 53}, survivors were only observed among mice immunized with the 1 and 0.1 μ g doses. There was slightly, more protection against fatal A66 infection among mice immunized with CFA than without, but the difference was not statistically significant. When the two sample rank test was used to analyze the time to death of mice infected with A66 we observed a statistically significant delay in the time to death in each immunized group as compared to the pooled controls.

↓

The 29 kDa N-terminal fragment of PspA can elicit protection against infection when injected with CFA

We have compared the immunogenicity, with and without CFA, of an isolated 29 kDa fragment composed of the first 260 amino acids of PspA. Unlike the case with FL-PspA, adjuvant was required for the 29 kDa fragment to elicit a protective response. This was observed even though the immunizing doses of the 29 kDa antigen used were 10 and 30 µg/mouse, or about 100 and 300 times the minimum dose of FL-PspA that can elicits protection in the absence of adjuvant.

Injection with CFA revealed the presence of additional protection eliciting antigen(s) in CDM-CC, and CDM-ET growth medium but not in the 2 percent choline chloride eluates of live cells

The observation that Freund's adjuvant could have such a major effect on the immunogenicity of the 29 kDa fragment (*Table 12*), prompted us to reexamine the immunogens described in *Table 2* to determine if immunization with adjuvant might enhance protection elicited by PspA-containing preparations or provide evidence for protection eliciting antigens in addition to PspA. By using CFA with the primary injection, the dose of PspA-containing growth medium (CDM-CC and CDM-ET) required to elicit protection was reduced from 10 - 30 µl (*Table 9*) down to 1 to 3 µl (*Table 13*). When CFA was used as an adjuvant with CDM-CC and CDM-ET from PspA⁻ strains WG44.1 and JY1119 we were able to elicit protective immune responses if material from ≥100 µl or more of media were injected. Thus, although there were apparently some protection eliciting components other than PspA in CDM-CC and CDM-ET growth media, PspA remained the major protection eliciting component even in the presence of adjuvant.

One of the media used for injection was CDM-ET in which JY2141 had been grown. This medium elicited protection against WU2 challenge even when injected at doses as low as 1 µl. It should be noted that although this strain does not make full-length PspA, it secretes a truncated molecule comprising the first 115 amino acids of PspA into the growth medium. Thus, unlike CDM-ET from WG44.1 and JY1119, CDM-ET from JY2141 has the potential to elicit PspA-specific immunity. In contrast to these results, the material eluted from JY2141 with 2 percent CC was relatively non-immunogenic even when emulsified with CFA. This result is consistent with the fact that the 115 amino acid N-terminal PspA fragment of JY2141 is not surface attached³⁷, and would be expected to be washed away prior to the elution with 2 percent CC.

Extension of studies to BALB/c mice and i.p. challenge route

The studies above all involve i.v. challenge of CBA/N mice expressing with the XID genetic defect. The i.v. route, used in the present studies provides a relevant model for bacteremia and sepsis, but pneumococci have higher LD₅₀s when injected i.v. than i.p. CBA/N mice are hypersusceptible to pneumococcal infection because of the XID defect. This genetic defect prevents them from having circulating naturally occurring antibody to phosphocholine. The absence of these antibodies has been shown to make XID mice several logs

immunologic purity of the isolated PspA was emphasized by the fact that immunization with it did not elicit any antibodies detectable on plates coated with CDM-ET supernatants of WG44.1.

Loading more than 300 ml on the 0.6 ml bed volume column did not result in an increased yield, which suggested that the column capacity had been reached. However, increasing the depth of the choline-Sepharose bed to greater than 0.5 cm, decreased the amount of PspA eluted from the column, presumably because of non-specific trapping of aggregates in the column matrix. The elution buffer contains 50 mM Tris acetate 0.25 M NaCl and 2% choline chloride. Elution without added NaCl or with 1M NaCl resulted in lower yields. Elution with less than 1% CC also reduced yields.

Immunization of mice with purified R36A PspA. For immunization we used only the first 3 ml fraction of the R36A column. Mice were immunized with two injections of 1, 0.1, or 0.01 μ g of R36A PspA, spaced two weeks apart. As controls, some mice were inoculated with a comparable dilutions of the first 3 ml fraction from the WG44.1 column. Purified FL-PspA elicited antibody to PspA at all doses regardless of whether CFA was used as an adjuvant (*Table 1*). In the absence of CFA the highest levels of antibody were seen with the 1 μ g dose of PspA. In the presence of CFA, however, the 0.1 μ g dose was as immunogenic as the 1 μ g dose.

To test the ability of the different doses of PspA to elicit protection against challenge we infected the immunized mice with two capsular type 3 strains, WU2 and A66. Although both of these strains are able to kill highly susceptible CBA/N XID mice at challenge doses of less than 10^2 , the A66 strain is several logs more virulent when BALB/c mice are used ^{47, 52}. The difference in virulence of A66 and WU2, was partially compensated for by challenging the immunized CBA/N mice with lower doses of strain A66 than WU2.

After immunization of CBA/N mice with 1 and 0.1 μ g doses of PspA we observed protection against WU2 challenge regardless of whether or not CFA was used as an adjuvant (*Table 4*). At the lowest dose, 0.01 μ g PspA, most of the mice immunized with PspA + CFA lived whereas most immunized with PspA alone did not; however, the difference was not statistically significant. When immunized mice were challenged with the more virulent strain A66 ^{47, 53}, survivors were only observed among mice immunized with the 1 and 0.1 μ g doses. There was slightly, more protection against fatal A66 infection among mice immunized with CFA than without, but the difference was not statistically significant. When the two sample rank test was used to analyze the time to death of mice infected with A66 we observed a statistically significant delay in the time to death in each immunized group as compared to the pooled controls.

more susceptible to pneumococci than isogenic mice lacking the immune defect. From the data in Table 14 it is clear, however, that immunization with PspA can protect against infection in mice lacking the XID defect even when the challenge is by the i.p. route. Thus, there is no reason to suspect that the results presented are necessarily dependent on the use of the CBA/N XID mouse or the i.v. route.

PspA is highly immunogenic

These studies provide the first quantitative data on the amount of purified FL-PspA that is required to elicit protective immunity in mice. The isolated PspA for these studies was obtained by taking advantage of the fact that the C-terminal half of PspA binds to cell surface choline³⁶. The isolated FL-PspA was found to be highly immunogenic in the mouse. Only two injections of 100 ng of PspA in the absence of adjuvant were required to elicit protection against otherwise fatal sepsis with greater than 100 LD₅₀ of capsular type 3 *S. pneumoniae*. When the first injection was given with adjuvant, doses as small as 10 ng could elicit protective response. The potent immunogenicity of PspA, and the ability to isolate it on choline-Sepharose columns provides a demonstration for the possible use of PspA as a vaccine in humans.

A large body of published^{17, 29, 37} as well as unpublished evidence indicates that the major protection eliciting epitopes of PspA are located in the α -helical (N-terminal) half of the molecule. From the present studies, it is clear that immunization with N-terminal fragments containing the first 115 or 260 of the 288 amino acid α -helical region are able to elicit protection when given with CFA. However, these fragments were not able to elicit protective responses without CFA. In the case of both the 115 and 260 amino acid fragments, even immunization at 100 times the minimum dose that is immunogenic for FL-PspA failed to elicit a protective response. This result is consistent with previous results showing that a fragment composed of the N-terminal 245 amino acids^{31, 37} could elicit protection against otherwise fatal pneumococcal infection of mice when the immunization was given with CFA³². In that study no immunization without CFA was attempted. Even though the C-terminal half of PspA may not contain major protection-eliciting epitopes it appears to contain sequence important in the immunogenicity of the molecule as a whole, since the full length molecule elicited much greater protection than the N-terminal fragments. The effect of the C terminal half on antigenicity may be in part that it doubles the size of the immunogen. Molecules containing the C-terminal half of PspA may also be especially immunogenic because they exhibit more extensive aggregation than is seen with fragments expressing only the α -helical region³⁸. Protein aggregates are known to generally be more antigenic and less tolerogenic than individual free molecules⁵⁴.

PspA is the major protection eliciting component of our pneumococcal extracts

Evidence that PspA is the major protection eliciting component of the CDM-ET, CDM-CC growth media and the two percent CC eluates was dependent on the use of mutant pneumococci that lacked the ability to produce FL-PspA. More than one *pspA* mutant strain was used to insure that the failure to elicit protection in the absence of FL-PspA was not a spurious result of non-PspA mutation blocking the production of some other antigen. Strains WG44.1 and JY1119 contain identical deletions that include the 5' end of the *pspA* genes and extend about 3 kb upstream of *pspA*³⁷. WG44.1 is a mutant of the non-encapsulated strain Rx1 and JY1119 was made by transforming capsular type 3 strain WU2 with the WG44.1 *pspA* mutation. In no case were preparations from WG44.1 and JY1119 as efficient at eliciting protection as those from the PspA⁺ strains. To rule out the possibility that protection elicited by preparations from the PspA⁺ strains was elicited by some non-PspA molecule also encoded by a 3 kb deletion linked to the mutant *pspA* genes of WG44.1 and JY1119, we also used strains JY2141 and LM34^{26, 37}. In these strains the Rx1 *pspA* gene has been insertionally inactivated causing the production of N-terminal fragments of 115 and 245 amino acids respectively. These strains have no other known mutations. Although Rx1 and R36A are closely related non-encapsulated strains, some of the studies included Rx1 as the PspA⁺ control since it is the isogenic partner to WG44.1, LM34, and JY2141. The N-terminal fragments produced by JY2141 and LM34 lack the surface anchor and are secreted into the medium³⁶. Two percent CC eluates of JY2141 were non-protection eliciting even in the presence of adjuvant. In the absence of adjuvant, CDM-ET from JY2141 was not protection-eliciting. LM34 was tested without CFA in only 3 mice, but gave results consistent with those obtained with JY2141.

Anticapsular antibodies are known to be protective against pneumococcal infection^{5, 19}. However, in these studies it is unlikely that they account for any of the protection we attributed to PspA. Our challenge strain bore the type 3 capsular polysaccharide and our primary source of PspA was strain R36A, which is a spontaneous non-encapsulated mutant of a capsular type 2 strain^{39, 41}. The R36A strain has been recently demonstrated to lack detectable type 3 capsule on the surface or in its cytoplasm⁵⁵. Furthermore, the CBA/N mice used in most of the studies are unable to make antibody responses to capsular type 3 polysaccharide⁵⁶.

Non-PspA protection eliciting components

The observation that CDM-CC and CDM-ET supernatants of WG44.1 could elicit protection when injected in large amounts with adjuvant, suggested that these supernatants contained at least trace amounts of non-PspA protection eliciting molecules. In the case of preparations containing PspA eluted from the surface of live washed pneumococci with 2 percent CC, there was no evidence for any protection eliciting components other than PspA, presumably because the protection-eliciting non-PspA proteins released into the media were removed by the previous washing step. The identity of the protection eliciting molecules in the WG44.1

supernatant are unknown. In this regard, it is of interest that unlike R36A, strain Rx1 has been shown to contain a very small amount of cytoplasmic type 3 polysaccharide (but totally lacks surface type 3 polysaccharide⁵⁵). This difference from Rx1 apparently came about through genetic manipulations in the construction of Rx1 from R36A^{39, 41}. Thus, preparations made from Rx1 or from its daughter strains WG44.1, LM34, or JY2141 could potentially contain small amounts of capsular polysaccharide. For a number of reasons however, it seems very unlikely that the non-PspA protection-eliciting material identified in these studies was type 3 capsular polysaccharide (expressed by the WU2 challenge strain: 1) growth of these strains was either in CDM-CC or CDM-ET, each of which prevent autolysin activity and lysis⁵⁷ that would be required to release the small amount of type 3 polysaccharide from the cytoplasm of the Rx1 family of strains; 2) CBA/N mice made protective responses to the non-PspA antigens, but express the XID immune response deficiency which permits responses to proteins, but blocks antibody to most polysaccharides⁴⁶, including type 3 capsular polysaccharide⁵⁶; and 3) immunogenicity of the non-PspA component required CFA, an adjuvant known to stimulate T-dependent (protein) rather than T-independent (polysaccharide) antibody responses.

A number of non-PspA protection eliciting pneumococcal proteins have been identified: pneumolysin, autolysin, neuraminidase, and PsaA which are 52, 36.5, 107 and 37 kDa respectively^{21, 58, 59, 60}. The non-PspA protection eliciting components reported here could be composed of a mixture of these and/or other non-identified proteins. Attempts to identify lambda clones producing non-PspA protection eliciting proteins as efficacious as PspA have not been successful²⁵.

Isolation of PspA

The protective capacity of the CDM-CC, CDM-ET and material eluted from live cells with 2% CC were similar in terms of the volume of the original culture from which the injected dose was derived. The major advantage of eluting the PspA from the surface of pneumococci with 2 percent CC is that the pneumococci may be grown in any standard growth medium, and do not have to be first adapted to a defined medium. Moreover, concentration of PspA can be accomplished by centrifugation of the pneumococci prior to the elution of the PspA. An advantage of using either CDM-CC and CDM-ET media was that these media prevented lysis and pneumococci could be grown into stationary phase without contaminating the preparations with cytoplasmic contents and membrane and wall components. A particular advantage of CDM-ET growth medium is that since it lacks high concentrations of choline the PspA contained in it can be adsorbed directly to a choline-Sepharose column for affinity purification.

One liter of CDM-ET growth medium contains about 400 µg of PspA, and we were able to isolate about 3/4 of it to very high purity. At 0.1 µg/dose, a liter of CDM-ET contains enough PspA to immunize about 4,000

mice; or possibly 40 - 400 humans. Our present batch size for a single column run is only 300 ml of CDM-ET. This could presumably be increased by increasing the amount of the adsorbent surface by increasing the diameter of the column. Using our present running buffer we have found that a choline-Sepharose resin depth of 0.5 cm was optimal; increases beyond 0.5 cm caused the overall yield to decrease rather than increase, even in the presence of larger loading volumes of R36A CDM-ET

Table 8 Pneumococcal Strains

| Strain | Capsule type | PspA expressed | Parent strain | Construction technique | References |
|--------|------------------|------------------------------|---------------|--|------------|
| D39 | 2 | full length | -- | clinical isolate | 26, 44 |
| R36A | non-encapsulated | full length | D39 | non-encapsulated mutant | 23, 44, 45 |
| Rx1 | non-encapsulated | full length | R36A | derived from R36A | 26, 39, 41 |
| WG44.1 | non-encapsulated | none | Rx1 | aberrant insertion inactivation with pKSD300 | 26, 37 |
| LM34 | non-encapsulated | aa 1-245 of Rx1 ^a | Rx1 | insertional inactivation with pKSD300 | 26, 37, 42 |
| JY2141 | non-encapsulated | aa 1-115 of Rx1 ^a | Rx1 | insertional inactivation with pJY4208 | 37 |
| WU2 | 3 | full length | -- | clinical isolate | 25, 46 |
| JY1119 | 3 | none | WU2 | transformation with WG44.1 DNA | 37 |
| A66 | 3 | full length | -- | clinical isolate | 44, 47 |

^a LM34 and LY2141 express fragments containing the first 245 and first 115 amino acids of Rx1 PspA respectively.

Table 9 PspA is the major protection-eliciting component in antigen preparations made by three different methods

| Preparation | Strain (PspA status) | Dose as volume of media in μ l ^a | Median Days Alive | Alive: Dead | P versus controls ^b |
|------------------------------|-----------------------------|---|-------------------|-------------|--------------------------------|
| 2% CC eluate from live cells | R36A (PspA ⁺) | 1000 | >21 | 2:0 | 0.03 |
| | | 200 | >21 | 2:0 | |
| | | 20 | >21 | 2:0 | |
| | | 2 | 1.5 | 0:2 | |
| | | all R36A | >21 | 6:2 | |
| | JY2141 (as 1-115) | 1000 | 3, >21 | 1:1 | |
| | | 200 | 1 | 0:2 | |
| | | 20 | 1 | 0:2 | |
| | | | | | |
| | | | | | |
| CDM-CC clarified medium | Rx1 (PspA ⁺) | 100 | >21 | 9:0 | <0.0001 |
| | | 30 | >21 | 2:1 | |
| | | 10 | 2 | 1:2 | |
| | | 3 | 2 | 0:3 | |
| | | ALL | 2, >21 | 12:6 | |
| | LM34 | 100 | 2, 2, >21 | 1:2 | 0.0004 |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | WG44.1 (PspA ⁻) | 100 | 2 | 0:9 | |
| | | 30 | 2 | 0:3 | |
| | | 10 | 2 | 0:3 | |
| | | 4 | 2 | 0:3 | |
| | | | | | |
| | WU2 (PspA ⁺) | 1000 | >21 | 3:0 | 0.05 |
| | | 100 | >21 | 1:0 | |
| | | ALL | >21 | 4:0 | |
| | JY1119 (PspA ⁻) | 1000 | 4 | 0:3 | 0.03 |
| | | | | | |
| | | | | | |
| | CDM-CC | 100 | 2 | 0:2 | |
| CDM-ET clarified medium | R36A (PspA ⁺) | 100 | >21 | 8:0 | <0.0001 |
| | | 10 | 3, >21 | 5:5 | |
| | | 1 | 1.5 | 3:5 | |
| | | 0.1 | 2 | 0:2 | |
| | | ALL | >21 | 16:12 | |
| | JY2141 (as 1-115) | 100 | 1.5 | 0:2 | 0.006 |
| | | 10 | 1.5 | 0:2 | |
| | WG44.1 (PspA ⁻) | 100 | 3 | 0:2 | |
| | | 10 | 1.5 | 0:2 | |
| | None | - | 2 | 0:14 | - |

^a Antigen dose is given as the volume of growth media from which the 0.1 ml of injected material was derived. Each mouse was injected twice i.p. with the indicated dose diluted as necessary in lactated Ringer's injection solution.

^b Controls used for statistical comparisons: 2% CC, all JY2141; CDM-CC Rx1, all WG44.1; CDM-CC WU2, JY1119; CDM-ET, all WG44.1 + all JY2141.

Table 10 Isolation of PspA from 300ml of CDM-ET media after the growth of R36A or WG44.1 pneumococci^a

| fraction | R36A | | | | WG44.1 | | |
|--------------|---------------|-------------------------------|---------------------------------------|--------------------------------------|-------------------|-------------------------------|---------------------------------------|
| | µg protein/ml | total µg protein ^b | max. reciprocal dot blot ^c | total dot blot units ^{b, d} | µg protein per/ml | total µg protein ^b | max. reciprocal dot blot ^c |
| growth media | 13.3 | 3,990 | 4 | 1200 | 13.7 | 4,110 | <1 |
| fall-through | 13.6 | 4,080 | 1 | 300 | 13.5 | 4,050 | <1 |
| 1st wash | | | <1 | | | | <1 |
| 10th wash | | | <1 | | | | <1 |
| elution #1 | 26 | 78 | 256 | 770 | <1 | — | <1 |
| elution #2 | 2 | 6 | 16 | 48 | <1 | — | <1 |
| elution #3 | <1 | — | 4 | 12 | <1 | — | <1 |
| total eluted | | 84 | | 830 | | — | <1 |

^a The columns were loaded with 300 ml of clarified CDM-ET medium after the growth of R36A or WG44.1.

The column was washed with 10 sequential 3 ml fractions of TBA. Elution was with TBA plus 2 percent CC.

^b Total µg protein or total dot blot units reflect the total protein in the 300 ml of the loading material or the 3 ml size of the eluted fractions.

^c MAb XiR278 was used in the immunoblots to detect PspA in dot blots.

^d Dot blot units were calculated as the reciprocal dot blot titer times the volume in ml.

Table 11 Purified full-length PspA is able to elicit protection against fatal sepsis in mice.

| Antigen | Dose ^a | Adjuvant or Diluent | Anti- PspA titer ^b (Log mean ± S.E.) | Challenge with 10 ^{5.1} WU2 | | | Challenge with 10 ^{4.2} A66 | | |
|--------------------------------|-------------------|---------------------------|--|---|-------------------------|---|---|-------------------------|--|
| | | | | Alive : Dead | Median Days Alive | P vs. pooled control ^c | Alive : Dead | Median Days Alive | P vs. pooled controls ^c |
| R36A (PspA ⁺) | 1 µg | Ringer's | 3.3 ± 0.2 | 5 : 0 | >21 | 0.015 | 2 : 3 | 4 | 0.002 |
| | 0.1 | Ringer's | 2.6 ± 0.2 | 4 : 0 | >21 | 0.041 | 1 : 4 | 4 | 0.0032 |
| | 0.01 | Ringer's | 2.7 ± 0.2 | 1 : 4 | 4 | n.s. | 0 : 5 | 3 | 0.0058 |
| | 1 µg | CFA | 3.5 ± 0.2 | 5 : 0 | >21 | 0.027 | 3 : 2 | >21 | 0.0012 |
| | 0.1 | CFA | 3.6 ± 0.1 | 5 : 0 | >21 | 0.013 | 2 : 3 | 4 | 0.0012 |
| | 0.01 | CFA | 3.1 ± 0.2 | 4 : 1 | >21 | 0.015 | 0 : 5 | 3 | 0.0058 |
| WG44.1 (PspA ⁻) | 3600 µl | Ringer's | <1.6 | n.d. | n.d. | | 1 : 4 | 3 | n.s. |
| | 360 | Ringer's | <1.6 | n.d. | n.d. | | 0 : 5 | 2 | n.s. |
| | 36 | Ringer's | <1.6 | n.d. | n.d. | | 0 : 5 | 2 | n.s. |
| | 3600 µl | CFA | <1.6 | n.d. | n.d. | | 0 : 5 | 2 | n.s. |
| | 360 | CFA | <1.6 | n.d. | n.d. | | 1 : 4 | 2 | n.s. |
| | 36 | CFA | <1.6 | n.d. | n.d. | | 0 : 5 | 2 | n.s. |
| saline | — | CFA | <1.6 | 1 : 5 | 4 | — | n.d. | n.d. | — |
| pooled controls | | | <1.6 | 1 : 5 | 4 | | 2 : 28 | 2 | — |

^a For comparison with the data in Table 2, it should be noted that the 1, 0.1, and 0.01 µg doses were derived from 3600, 360, and 36 µl of R36A growth media. Equivalent dilutions of the PspA⁻ eluate from strain WG44.1 were injected as controls. The amount of the WG44.1 preparations injected is listed as 3600, 360, and 36 µl and corresponds to the volume original growth medium from which the doses of WG44.1 was prepared.

^b Antibody values were expressed as reciprocal ELISA titer.

^c P values calculated by the Wilcoxon two sample rank test. By Kruskal-Wallis nonparametric ANOVA for the WU2 challenge was significant at P=0.01, for A66 significance was at P<0.0001.

Table 12 The 29 kDa N-terminal fragment of Rxl PspA must be injected with adjuvant to elicit protection against WU2^a

| μ g 29 kDa PspA | Adjuvant or diluent | Median Days Alive | Alive : Dead | P versus none ^b |
|---------------------|---------------------|-------------------|--------------|----------------------------|
| 30 | CFA | >21 | 3 : 0 | 0.0006 |
| 3 | CFA | >21 | 3 : 0 | |
| 30 | Ringer's | 2 | 0 : 3 | |
| 3 | Ringer's | 2 | 1 : 2 | |
| none | CFA | 2 | 0 : 7 | |
| none | Ringer's | 2 | 0 : 7 | |

^aThe 29 kDa fragment comprises the first 260 amino acids of PspA.

^bFor the calculation of P values the 30 μ g and 3 μ g data were pooled; mice immunized with PspA + CFA were compared to CFA controls; mice immunized with PspA + Ringer's were compared to controls immunized with Ringer's. Only the statistically significant P values are shown. The calculated P value of PspA + CFA versus CFA alone, was 0.0006 by both the Wilcoxon two sample rank test and the Fisher exact test.

Table 13 PspA is not the only protection eliciting molecule released from pneumococci by interference with binding to choline on the surface of pneumococci

| Preparation | Strain (PspA status) | Dose (as volume in µl) | Median Day Alive | Alive: Dead | P values ^a |
|---|--|------------------------------|------------------------|----------------|-----------------------|
| | | | | | P vs. all JY2141 |
| 2% CC eluate from live cells | R36A (PspA ⁺) | 1000 | >21 | 2 : 0 | |
| | | 200 | >21 | 5 : 0 | 0.02 |
| | | 20 | >21 | 5 : 0 | 0.02 |
| | | 2 | >21 | 5 : 0 | 0.02 |
| | | all R36A | >21 | 17 : 0 | 0.001 |
| | JY2141 (as 1 - 115) | 1000 | >21 | 2 : 0 | |
| | | 200 | 1 | 0 : 2 | |
| | | 20 | 1 | 0 : 2 | |
| | | 2 | 1 | 0 : 2 | |
| | | all JY2141 | 1 | 2 : 6 | |
| | | | | | P versus pooled cont. |
| CDM-CC clarified medium + CFA | Rx1 (PspA ⁺) | 1000 | >21 | 3 : 0 | 0.002 |
| | | 100 | >21 | 3 : 0 | 0.002 |
| | WU2 (PspA ⁺) | 1000 | >21 | 3 : 0 | 0.002 |
| | | 100 | >21 | 3 : 0 | 0.002 |
| | | 3 | >21 | 3 : 0 | 0.002 |
| | WG44.1 (PspA ⁻) | 1000 | >21 | 5 : 1 | <0.0001 |
| | | 100 | 2.5 | 2 : 4 | 0.002 |
| | JY1119 (PspA ⁻) | 1000 | >21 | 3 : 0 | 0.002 |
| | | 100 | >21 | 3 : 0 | 0.002 |
| | CDM-ET clarified medium + CFA | R36A (PspA ⁺) | 1000 | >21 | 3 : 1 |
| 10 | | | >21 | 4 : 0 | 0.004 |
| 1 | | | >21 | 3 : 1 | 0.004 |
| 0.2 | | | 2 | 0 : 4 | |
| JY2141 (as 1 - 115) | | 10 | >21 | 2 : 0 | |
| | | 1 | >21 | 2 : 0 | |
| | | all JY2141 | - | >21 | 4 : 0 |
| WG44.1 (PspA ⁻) | | 100 | >21 | 2 : 0 | |
| | 10 | 2 | 0 : 2 | | |
| CDM-ET only None | + CFA none | | 2 | 0 : 9 | |
| | | | 1.5 | 0 : 4 | |
| Pooled Controls ^b | | | 2 | 0 : 13 | |

^a In cases where there were not statistically significant results no P value was shown.

^b "Pooled Controls" refers to "CDM-ET only" Data and "None" data.

Table 14 Immunization of BALB/c mice with isolated PspA elicits protection against WU2 *S. pneumoniae*

| Antigen | | Adjuvant or diluent | Challenge | | Days to Death | <i>P</i> vs. controls TSR/FE ^b |
|---------------------------------|-------------------|------------------------|-----------|--------------|------------------|--|
| Source | Dose ^a | | Log CFU | Route | | |
| R36A (PspA ⁺) | 1μg | CFA | 4 | i.p. | 2, >21, >21, >21 | 0.06/0.03 |
| WG44.1 (PspA ⁻) | 100μl | CFA | 4 | i.p. | 2, 3 | |
| None | - | CFA | 4 | i.p. | 2, 2, 2, 4 | |
| R36A (PspA ⁺) | 1μg | none | 6 | i.v. | 2, >21, >21, >21 | 0.06/0.03 |
| WG44.1 (PspA ⁻) | 100μl | none | 6 | i.v. | 5, 7 | |
| None | - | none | 6 | i.v. | 2, 2, 2, 3 | |
| Pooled i.v. and i.p. results | | | | i.v. or i.p. | | 0.008/0.0007 |

^a The 1μg dose of R36A PspA was isolated from 100μl of CDM-ET medium. As a control mice were injected with an corresponding volume of choline-column effluent from a mock isolation of PspA from the PspA⁻ strain WG44.1. The dose of WG44.1 material is expressed as 100 μl since this is the volume CDM-ET from which the injected column effluent was derived.

^b *P* values calculated by Wilcoxon two-sample rank test, TSR, or Fisher exact, FE versus pooled controls for each group. "Pooled controls" include data obtained with by injection of "WG44.1" and "none". The i.p. and i.v. studies gave comparable results. When the data from the two studies were pooled the *P* values by both tests were ≤0.008. In cases where there were not statistically significant results no *P* value was shown.

REFERENCES

1. Anonymous. Pneumococcal polysaccharide vaccine. *MMWR* 1981, 30, 410-419
2. Farley, J.J., King, J.C., Nair, P., al., e. Invasive pneumococcal disease among infected and uninfected children of mothers with immunodeficiency virus infection. *J. Pediatr.* 1994, 124, 853-858
3. Schwartz, B., Gove, S., Lob-Lovit, J., Kirkwood, B.R. Potential interactions for the prevention of childhood pneumonia in developing countries: etiology of acute lower respiratory infections among young children in developing countries. *Ped. Infect. Dis.* in Press,
4. Avery, O.T., Goebel, W.F. Chemoimmunological studies of the soluble specific substance of pneumococcus. I. The isolation and properties of the acetyl polysaccharide of pneumococcus type 1. *J. Exp. Med.* 1933, 58, 731 - 755
5. Austrian, R. Pneumococcal Vaccine: Development and Prospects. *Am. J. Med* 1979, 67, 547-549
6. Shapiro, E.D., Berg, A.T., Austrian, R., Schroeder, D., Parcells, V., Margolis, A., Adair, R.K., Clemmens, J.D. Protective efficacy of polyvalent pneumococcal polysaccharide vaccine. *N. Engl. J. Med* 1991, 325, 1453-1460
7. Fedson, D.S. Pneumococcal vaccination in the prevention of community-acquired pneumonia: an optimistic view of cost-effectiveness. *Sem. Resp. Infect.* 1993, 8, 285-293
8. Robbins, J.B., Austrian, R., Lee, C.-J., Rastogi, S.C., Schiffman, G., Henrichsen, J., Makela, P.H., Broome, C.V., Facklam, R.R., Tiesjema, R.H., Parke, J.C., Jr. Considerations for formulating the second-generation pneumococcal capsular polysaccharide vaccine with emphasis on the cross-reactive types within groups. *J Infect Dis* 1983, 148, 1136-1159
9. Gotschlich, E.C., Goldschneider, I., Lepow, M.L., Gold, R. *The immune response to bacterial polysaccharides in man*. Antibodies in human diagnosis and therapy, . New York, Raven, 1977, 391-402.
10. Cowan, M.J., Ammann, A.J., Wara, D.W., Howie, V.M., Schultz, L., Doyle, N., Kaplan, M. Pneumococcal polysaccharide immunization in infants and children. *Pediatrics* 1978, 62, 721-727

11. Mond, J.J., Lees, A., Snapper, C.M. T cell-independent antigens type 2. *Ann. Rev. Immunol.* 1995, 13, 655-692
12. Chiu, S.S., Greenberg, P.D., Marcy, S.M., Wong, V.K., Chang, S.J., Chiu, C.Y., Ward, J.I. Mucosal antibody responses in infants following immunization with *Haemophilus influenzae*. *Pediatric Res. Abstracts* 1994, 35, 10A
13. Kauppi, M., Eskola, J., Kathry, H.H. *influenzae* type b (Hib) conjugate vaccines induce mucosal IgA1 and IgA2 antibody responses in infants and children. *ICAAC Abstracts* 1993, 33, 174
14. Dagen, R., Melamed, R., Abramson, O., Figlansky, L., Greenberg, D., Mendelman, P.M., Bohidar, N., Ter-Minassian, D., Cvanovich, N., Lov, D., Rusk, C., Donnelly, J., Yagupsky, P. Effect of heptavalent pneumococcal-OMPC conjugate vaccine on nasopharyngeal carriage when administered during the 2nd year of life. *Pediat. Res.* 1995, 37, 172A
15. Fattom, A., Vann, W.F., Szu, S.C., Sutton, A., Bryla, D., Shiffman, G., Robbins, J.B., Schneerson, R. Synthesis and physiochemical and immunological characterization of pneumomcoccus type 12F polysaccharide-diphtheria toxoid conjugates. *Infect. Immun.* 1988, 56, 2292-2298
16. Kennedy, D., Derosse, C., E., A. Immunologic response of 12 -18 month children to licensed pneumococcal polysaccharide vaccine primed with *Streptococcus pneumoniae* 19F conjugate vaccine. *ICAAC* 1994, Abstract, G89
17. McDaniel, L.S., Ralph, B.A., McDaniel, D.O., Briles, D.E. Localization of protection-eliciting epitopes on PspA of *Streptococcus pneumoniae* between amino acid residues 192 and 260. *Microbial Pathogenesis* 1994, 17, 323-337
18. Langermann, S., Palaszynski, S.R., Burlein, J.E., Koenig, S., Hanson, M.S., Briles, D.E., Stover, C.K. Protective humoral response against pneumococcal infection in mice elicited by recombinant Bacille Calmette-Guérin vaccines expressing PspA. *J. Exp. Med.* 1994, 180, 2277-2286

19. Siber, G.R. Pneumococcal Disease: Prospects for a New Generation of Vaccines. *Science* 1994, 265, 1385-1387
20. Lock, R.A., Hansman, D., Paton, J.C. Comparative efficacy of autolysin and pneumolysin as immunogens protecting mice against infection by *Streptococcus pneumoniae*. *Microbial Pathogenesis* 1992, 12, 137-143
21. Sampson, J.S., O'Connor, S.P., Stinson, A.R., Tharpe, J.A., Russell, H. Cloning and nucleotide sequence analysis of *psaA*, the *Streptococcus pneumoniae* gene encoding a 37-kilodalton protein homologous to previously reported *Streptococcus* sp. adhesins. *Infect. Immun.* 1994, 62, 319
22. Paton, J.C., Lock, R.A., Lee, C.-J., Li, J.P., Berry, A.M., Mitchell. Purification and immunogenicity of genetically obtained pneumolysin toxoids and their conjugation to *Streptococcus pneumoniae* type 19F polysaccharide. *Infect. Immun.* 1991, 59, 2297-2304
23. McDaniel, L.S., Scott, G., Kearney, J.F., Briles, D.E. Monoclonal antibodies against protease sensitive pneumococcal antigens can protect mice from fatal infection with *Streptococcus pneumoniae*. *J. Exp. Med.* 1984, 160, 386-397
24. Briles, D.E., Forman, C., Horowitz, J.C., Volanakis, J.E., Benjamin, W.H., Jr., McDaniel, L.S., Eldridge, J., Brooks, J. Antipneumococcal effects of C-reactive protein and monoclonal antibodies to pneumococcal cell wall and capsular antigens. *Infect. Immun.* 1989, 57, 1457 - 1464
25. McDaniel, L.S., Sheffield, J.S., Delucchi, P., Briles, D.E. PspA, a surface protein of *Streptococcus pneumoniae*, is capable of eliciting protection against pneumococci of more than one capsular type. *Infect. Immun.* 1991, 59, 222-228
26. McDaniel, L.S., Yother, J., Vijayakumar, M., McGarry, L., Guild, W.R., Briles, D.E. Use of insertional inactivation to facilitate studies of biological properties of pneumococcal surface protein A (PspA). *J. Exp. Med.* 1987, 165, 381-394
27. Yother, J., McDaniel, L.S., Crain, M.J., Talkington, D.F., Briles, D.E. *Pneumococcal surface protein A: Structural analysis and biological significance* In: Dunny, G.M., Cleary, P.P., McKay, L.L. ed. Genetics

- and Molecular Biology of Streptococci, Lactococci, and Enterococci. Washington, DC: American Society for Microbiology, 1991, 88-91
28. Waltman, W.D., II, McDaniel, L.S., Gray, B.M., Briles, D.E. Variation in the molecular weight of PspA (Pneumococcal Surface Protein A) among *Streptococcus pneumoniae*. *Microb. Pathog.* 1990, 8, 61-69
 29. Crain, M.J., Waltman, W.D., II, Turner, J.S., Yother, J., Talkington, D.E., McDaniel, L.M., Gray, B.M., Briles, D.E. Pneumococcal surface protein A (PspA) is serologically highly variable and is expressed by all clinically important capsular serotypes of *Streptococcus pneumoniae*. *Infect. Immun.* 1990, 58, 3293-3299
 30. McDaniel, L.S., Scott, G., Widenhofer, K., Carroll, Briles, D.E. Analysis of a surface protein of *Streptococcus pneumoniae* recognized by protective monoclonal antibodies. *Microb. Pathog.* 1986, 1, 519-531
 31. Yother, J., Briles, D.E. Structural properties and evolutionary relationships of PspA, a surface protein of *Streptococcus pneumoniae*, as revealed by sequence analysis. *J. Bact.* 1992, 174, 601-609
 32. Talkington, D.F., Crimmins, D.L., Voellinger, D.C., Jother, J., Briles, D.E. A 43-kilodalton pneumococcal surface protein, PspA: isolation, protective abilities, and structural analysis of the amino-terminal sequence. *Infect. Immun.* 1991, 59:, 1285-1289
 33. McDaniel, L.S., McEdaniel, D.O. Genetic analysis of the gene encoding type 12 PspA of *Streptococcus pneumoniae* strain EF5668 In: Feretti, J.J., Gilmore, M.S., Khenhammer, T.R., Brown, F. ed. Genetics of the streptococci, enterococci, and lactococci. Basel: Dev. Biol. Stand. Basel Krager, 1995, 283-286
 34. Fischetti, V.A., Pancholi, V., Schneewind, O. Conservation of a hexapeptide sequence in the anchor region of surface proteins from gram-positive cocci. *Molec. Microbiol.* 1990, 4, 1603-1605
 35. Schneewind, O., Fowler, A., Faull, K.F. Structure of cell wall anchor of cell surface proteins in *Staphylococcus aureus*. *Science* 1995, 268, 103-106
 36. Yother, J., White, J.M. Novel surface attachment mechanism for the streptococcus pneumoniae protein PspA. *J. Bact.* 1994, 176, 2976-2985

37. Yother, J., Handsome, G.L., Briles, D.E. Truncated forms of PspA that are secreted from *Streptococcus pneumoniae* and their use in functional studies and cloning of the *pspA* gene. *J. Bact.* 1992, 174, 610-618
38. Talkington, D.F., Voellinger, D.C., McDaniel, L.S., Briles, D.E. Analysis of pneumococcal PspA microheterogeneity in SDS polyacrylamide gels and the association of PspA with the cell membrane. *Microbial Pathogenesis* 1992, 13, 343-355
39. Smith, M.D., Guild, W.R. A plasmid in *Streptococcus pneumoniae*. *J. Bacteriol.* 1979, 137, 735-739
40. Shoemaker, N.B., Guild, W.R. Destruction of low efficiency markers is a slow process occurring at a heteroduplex stage of transformation. *Mol. Gen. Genet.* 1974, 128, 283-290
41. Raven, A.W. Reciprocal capsular transformations of pneumococci. *J. Bact.* 1959, 77, 296-309
42. McDaniel, L.S., Sheffield, J.S., Swiatlo, E., Yother, J., Crain, M.J., Briles, D.E. Molecular localization of variable and conserved regions of *pspA*, and identification of additional *pspA* homologous sequences in *Streptococcus pneumoniae*. *Microbial Pathogenesis* 1992, 13, 261-269
43. Rijn, V.D., Kessler, R.E. Growth characteristics of Group A Streptococci in a new chemically defined medium. *Infect. Immun.* 1980, 27, 444-448
44. Avery, O.T., MacLeod, C.M., McCarty, M. Studies on the chemical nature of the substance inducing transformation of pneumococcal types. Induction of transformation by a desoxyribonucleic acid fraction isolated from pneumococcus type III. *J. Exp. Med.* 1944, 79, 137-158
45. McCarty, M. *The transforming principle*. New York, Norton, 1985, 252.
46. Briles, D.E., Nahm, M., Schroer, K., Davie, J., Baker, P., Kearney, J., Barletta, R. Antiphosphocholine antibodies found in normal mouse serum are protective against intravenous infection with type 3 *Streptococcus pneumoniae*. *J. Exp. Med.* 1981, 153, 694-705
47. Briles, D.E., Crain, M.J., Gray, B.M., Forman, C., Yother, J. A strong association between capsular type and mouse virulence among human isolates of *Streptococcus pneumoniae*. *Infect. Immun.* 1992, 60, 111-116

48. Waltman, W.D., II, McDaniel, L.S., Andersson, B., Bland, L., Gray, B.M., Svanborg-Eden, C., Briles, D.E. Protein serotyping of *Streptococcus pneumoniae* based on reactivity to six monoclonal antibodies. *Microb. Pathog.* 1988, 5, 159-167
49. Tomasz, A. Surface components of *Streptococcus pneumoniae*. *Rev. Infect. Dis* 1981, 3, 190-211
50. Garcia, J.L., Garcia, E., Lopez, R. Overproduction and rapid purification of the amidase of *Streptococcus pneumoniae*. *Arch. Microbiol.* 1987, 149, 52-56
51. Osborn, M.J., Munson, J. Separation of the inner (cytoplasmic) and outer membranes of gram negative bacteria. *Methods Enzymol.* 1974, 31A, 642-653
52. Briles, D.E., Horowitz, J., McDaniel, L.S., Benjamin, W.H., Jr., Claflin, J.L., Booker, C.L., Scott, G., Forman, C. Genetic control of susceptibility to pneumococcal infection. *Curr. Top. Microbiol. Immunol.* 1986, 124, 103-120
53. Briles, D.E., Forman, C., Crain, M. Mouse antibody to phosphocholine can protect mice from infection with mouse-virulent human isolates of *Streptococcus pneumoniae*. *Infect. Immun.* 1992, 60, 1957-1962
54. Weigle, W.O. *Immunological unresponsiveness*. Academic Press, New York, New York, 1973.
55. Dillard, J.P., Yother, J. Genetic and molecular characterization of capsular polysaccharide biosynthesis in *Streptococcus pneumoniae* type 3. *Molec. Microbiol.* 1994, 12, 959-972
56. Amsbaugh, D.F., Hansen, C.T., Prescott, B., Stashak, P.W., Barthold, D.R., Baker, P.J. Genetic control of the antibody response to type III pneumococcal polysaccharide in mice. I. Evidence that an X-linked gene plays a decisive role in determining responsiveness. *J. Exp. Med* 1972, 136, 931-949
57. Tomasz, A. Biological consequences of the replacement of choline by ethanolamine in the cell wall of pneumococcus: chain formation, loss of transformability, and loss of autolysis. *Proc. Natl. Acad. Sci. USA* 1968, 59, 86-93

58. Paton, J.C., Lock, R.A., Hansman, D.C. Effect of immunization with pneumolysin on survival time of mice challenged with *Streptococcus pneumoniae*. *Infect. Immun.* 1983, 40, 548-552
59. Berry, A.M., Lock, R.A., Hansman, D., Paton, J.C. Contribution of autolysin to virulence of *streptococcus pneumoniae*. *Infect. Immun.* 1989, 57, 2324-2330
60. Lock, R.A., Paton, J.C., Hansman, D. Purification and immunologic characterization of neuraminidase produced by *Streptococcus pneumoniae*. *Microbial Pathogenesis* 1988, 4, 33-43
61. Tuomanen, E., Liu, H., Hengstler, B., Zak, O., Tomasz, A. The Induction of meningeal inflammation by components of the pneumococcal cell wall. 1985, 151, 859-868
62. Tuomanen, E., Tomasz, A., Hengstler, B., Zak, O. The relative role of bacterial cell wall and capsule in the induction of inflammation in pneumococcal meningitis. *J. Infect. Dis.* 1985, 151, 535-540
63. Paton, J.C. *Pathogenesis of pneumococcal disease*. 1993, 363-368
64. Briese, T., Hakenbeck, R. Interaction of the pneumococcal amidase with lipoteichoic acid and choline. 1985, 146, 417-427

EXAMPLE 4 - Evidence For Simultaneous Expression of Two PspAs

From Southern blot analysis there has been an issue as to whether most isolates of *S. pneumoniae* has two DNA sequences that hybridize with both 5' and 3' halves of Rx1 pspA, or whether this is an artifact of Southern blot. When bacterial lysates have been examined by Western blot, the results have always been consistent with the production of a single PspA by each isolate. This Example provides evidence for the first time that two PspAs of different apparent molecular weights and different serotypes can be simultaneously expressed by the same isolate.

Different PspAs frequently share cross-reactive epitopes, and an immune serum to one PspA was able to recognize PspAs on all pneumococci. In spite of these similarities, PspAs of different strains can generally be distinguished by their molecular weights and by their reactivity with a panel of PspA-specific monoclonal antibodies (MAbs).

A serotyping system for PspA has been developed which uses a panel of seven MAbs. PspA serotypes are designated based on the pattern of positive or negative reactivity in immunoblots with this panel of MAbs. Among a panel of 57 independent isolates of 9 capsular groups/types, 31 PspA serotypes were observed. The large diversity of PspA was substantiated in a subsequent study of 51 capsular serotype 6B isolates from Alaska, provided by Alan Parkinson at the Arctic Investigations Laboratory of the Centers for Disease Control and Prevention.

Among these 51 capsular type 6B isolates were observed 22 different PspAs based on PspA serotype and molecular weight variations of PspA.

While most pneumococcal strains appear to have two DNA sequences homologous with both the 5' and 3' halves of *pspA*, site-specific truncation mutations of *Rx1* have revealed that one these, *pspA*, encodes PspA. The other sequence has been provisionally designated as the *pspA*-like sequence. At present whether the *pspA*-like sequence makes a gene product is unknown. Evidence that the *pspA* and *pspA*-like genes are homologous but distinct groups of alleles comes from Southern blot analysis at high stringencies. Additional evidence that *pspA* and the *pspA*-like loci are distinct comes from studies using PCR primers that permit amplification of a single product approximately 2Kb in size from 70% of pneumococci. For the remaining 30% of pneumococci no amplification was observed with the primers used.

Evidence for two PspAs:

When the strains of MC25-28 were examined with the panel of seven MAbs specific for different PspA epitopes, all four demonstrated the same patterns of reactivity (Fig. 4). The MAbs XiR278 and 2A4 detected a PspA molecule with an apparent molecular weight of 190 KDa in each isolate. In accordance with the previous PspA serotyping system, the 190 KDa molecule was designated as PspA type 6 because of its reactivity with XiR278 and 2A4, but none of the five other MAbs in the typing system.

Each isolate also produced a second PspA molecule with an apparent molecular weight 82 KDa. The 82 KDa PspA in each isolate was detected only with the MAb 7D2 and was designated as type 34. No reactivity was detected with MAbs Xi126, Xi64, 1A4, or SR4W4. The fact that all four capsular 6B strains exhibit two PspAs, based on both molecular weights and PspA serotypes, suggested that they might be members of the same clone.

Simultaneous production of both PspAs:

Results from the colony immunoblotting showed that both PspAs were present simultaneously in each colony of these isolates when grown in vitro. All colonies on each plate of the original culture, as well as all of the progeny colonies from a single colony, reacted with MAbs XiR278, 2A4, and 7D2.

Number of *pspA* genes:

One explanation for the second PspA molecule was that these strains contained an extra *pspA* gene. Since most strains contain a *pspA* gene and a *pspA*-like gene it was expected that if an extra gene were present one might observe at least three *pspA* homologous loci in isolates MC25-28. In *Hind* III digests of MC25-28 each strain revealed a 7.7 and 3.6 Kb band when probed with pLSM*pspA*13/2 (Figure 5A). In comparison, when Rx1 DNA was digested with *Hind* III and hybridized with pLSM*pspA*13.2, homologous sequences were detected on 9.1 and 4.2 Kb fragments as expected from previous studies (9) (Figure 5A). Results consistent with only two *pspA*-homologous genes in MC25-28 were

also obtained with digestion using four additional enzymes (Table 15).

In previous studies it has been reported that probes for the 5' half of *pspA* (encoding the alpha-helical half of the protein) bind the *pspA*-like sequence of most strains only at a stringency of around 90%. With chromosomal digests of MC25-28 we observed that the 5' Rx1 probe of pLSM*pspA*12/6 bound both *pspA* homologous bands at a stringency of greater than 95 percent. The same probe bound only the *pspA* containing fragment Rx1 at a stringency above 95 percent (Figure 5B).

Further characterization of the *pspA* gene was done by RFLP analysis of PCR amplified *pspA* from each strain. Since previous studies indicated that individual strains yielded only one product, and since the amplification is carried out with primers based on a known *pspA* sequence, it seems likely that in each case the amplified products represent the *pspA* rather than the *pspA*-like gene. When MC25-28 were subjected to this procedure, an amplified *pspA* product of 2.1 Kb was produced in each case. When digested with *Hha* 1 digest the sum of the fragments obtained with each enzyme was approximately equal to the size of the 2.1 Kb amplified product (Figure 6). These results suggest that the 2.1 Kb amplified DNA represents the amplified product of only a single DNA sequence. Rx1, by comparison, produced an amplified product of 2.0 Kb and five

fragments of 0.76, 0.468, 0.390, 0.349 and 0.120, when digested with *Hha* I as expected from its known *pspA* sequence.

The four isolates examined in this Example are the first in which two PspAs have unambiguously been observed. The interpretation that two PspAs are simultaneously expressed by a single pneumococcal isolate is based on the observation that bands of different molecular weights were detected by different MAbs to PspA. Isolates used in this study were from a group originally selected for study by Brian Spratt because of their resistance to penicillin. It is very likely that all four of the isolates making two PspAs are related since they share PspA serotypes, amplified *pspA* RFLPs, chromosomal *pspA* RFLPs, capsule type, and resistance to penicillin.

The interpretation of studies presented here, showing the existence of two PspAs in the four strains MC25-28, must be set in the context of what is known about the serology PspA as detected by Western blots. PspAs of different strains have been shown previously to exhibit apparent molecular weight sizes ranging from 60 to 200 KDa as detected by Western blots. At least part of this difference in size is attributable to secondary structure. Even for the PspA of a single isolate, bands of several sizes are generally observed. Mutation and immunochemistry studies have demonstrated, however, that all of the different sized PspA bands from Rx1 are made by a single gene capable of encoding a 69 KDa protein. The heterogeneity of band

size on Western blots of PspA made by a single strain appears to be due to both degradation and polymerization.

PspA was originally defined by reciprocal absorption studies demonstrating that a panel of MAbs to Rx1 surface proteins each reacted with some protein and later by studies using Rx1 and WU2 derivatives expressing various truncated forms of PspA. In both cases it was clear that each MAbs to the PspA of a given strain reacted with the same protein. Such detailed studies have not been done with each of the several hundred human isolates. It is possible that with some isolates, reactivity of the MAbs with two PspAs may have gone unnoticed. This could have happened if all reactive antibodies detected both PspAs of the same isolate, or if the most prominent migration bands from each of the two PspAs co-migrated. With isolates MC25-28 the observation of two PspAs was possible because clearly distinguishable bands of different molecular weights reacted preferentially with different MAbs.

Applicants favor the interpretation that isolates MC25-28 each make two PspAs, because an alternative possibility, namely, that the 190 KDa PspA detected by MAbs XiR278 and 2A4 might be a dimer of the 84 KDa monomer detected by MAb 7D2, if the epitopes recognized by the different MAbs were dependent on either the dimeric or monomeric status of the protein, seems unlikely since whenever MAbs react with the PspA of a strain, they usually detect both the monomeric and the dimeric forms. No

other isolates have been observed where some MABs detected only the apparent dimeric form of PspA while others detected only the monomeric form.

There could be several possible explanations for the failure to observe two PspAs produced by most strains. 1) All pneumococci might make two *pspAs* in culture, but MABs generally recognize only one of them (perhaps in this isolate there has been a recombination between *pspa* DNA and the *pspA*-like locus, thus allowing that locus to make a product detected by MAB to PspA). 2) All pneumococci can have two *pspAs* but the expression of one of them generally does not occur under *in vitro* growth conditions. 3) The *pspA*-like locus is normally a nonfunctional pseudogene sequence that for an unexplained reason has become functional in these isolates.

It seems unlikely that the expression of only a single PspA by most strains is the result of a phase shift that permits the expression of only the *pspA* or *pspA*-like gene at any one time, since many of the strains examined repeatedly and consistently produce the same PspA. In the case of strains MC25-28, the appearance of two PspAs is apparently not the result of a phase switch, since individual colonies produced both the type 6 and the type 34 PspAs.

Presumably in these four strains, the second PspA protein is produced by the *pspA*-like DNA sequence. At high stringency, the probe comprising the coding region of the alpha-

helical half of PspA recognized both *pspA* homologous sequences of MC25-28 but not the *pspA*-like sequence of Rx1. This finding indicates that the *pspA*-like sequence of MC25-28 is more similar to the Rx1 *pspA* sequence than is the Rx1 *pspA*-like sequence. If the *pspA*-like sequence of these strains is more similar to *pspA* than most *pspA*-like sequences, it could explain why we were able to see the products of *pspA*-like genes of these strains with our MAbs. The finding of two families of PspAs made *in vivo* by pneumococci, allows for use of the second PspA in compositions, as well as the use of DNA primers or probes for the second gene for more conclusive detecting, determining or isolating of pneumococci.

Isolates and Bacterial Cell Culture:

Pneumococcal isolates described in these studies were cultured from patients in Barcelona, Spain (one adult at Bellvitge Hospital, and three children at San Juan de Dios) between 1986 and 1988 (Table 2). These penicillin resistant pneumococci originally in the collection of Dr. Brian Spratt were shared with applicants by Dr. Alexander Tomasz at the Rockefeller Institute. Rx1 is a rough pneumococcus used in previous studies, and it is the first isolate in which *pspA* was sequenced. Bacteria were grown in Todd-Hewitt broth with 0.5% yeast extract or on blood agar plates overnight in a candle jar. Capsular serotype was confirmed by cell agglutination using Danish antisera (Statens Seruminstitut, Copenhagen, Denmark) as

previously described. The isolates were subsequently typed as 6B by Quellung reaction, utilizing rabbit antisera against 6A or 6B capsule antigen prepared by Dr. Barry Gray.

Bacterial lysates:

Cell lysates were prepared by incubating the bacterial cell pellet with 0.1% sodium deoxycholate, 0.01% sodium dedecylsulfate (SDS), and 0.15 M sodium citrate, and then diluting the lysate in 0.5M Tris hydrochloride (pH 6.8) as previously described. Total pneumococcal protein in the lysates was quantitated by the bicinchonic acid method (BCA Protein Assay Reagent; Pierce Chemical Company, Rockford, IL).

PspA serotyping:

Serotyping of PspA was performed according to previously published methods. Briefly, pneumococcal cell lysates were subjected to SDS-PAGE, transferred to nitrocellulose membranes, and developed as Western blots using a panel of seven MAbs to PspA. PspA serotypes were assigned based on the particular combination of MAbs with which each PspA was reactive.

Colony Immunoblotting:

A ten ml tube of Todd-Hewitt broth with 0.5% yeast extract was inoculated with overnight growth of MC23 from a blood agar plate. The isolate was allowed to grow to a concentration of 10^7 cells/ml as determined by an O.D. of 0.07 at 590nm. MC23 was serially diluted and spread-plated on blood agar plates to give approximately 100 cells per plate. The plates were allowed

to grow overnight in a candle jar, and a single block agar plate with well-defined colonies was selected. Four nitrocellulose membranes were consecutively placed on the plate. Each membrane was lightly weighted and left in place for 5 minutes. In order to investigate the possibility of phase-variation between the two proteins detected on Western blots a single colony was picked from the plate, resuspended in ringers, and spread-plated onto a blood agar plate. The membranes were developed as Western blots according to PspA serotyping methods.

Chromosomal DNA Preparation:

Pneumococcal chromosomal DNA was prepared as in Example 9. The cells were harvested, washed, lysed, and digested with 0.5% (wt/vol) SDS and 100µg/ml proteinase K at 37°C for 1 hour. The cell wall debris, proteins, and polysaccharides were complexed with 1% hexadecyl trimethyl ammonium bromide (CTAB) and 0.7M sodium chloride at 65°C for 20 minutes, then extracted with chloroform/isoamyl alcohol. DNA was precipitated with 0.6 volumes isopropanol, washed, and resuspended in 10mM Tris-HCL, 1mM EDTA, pH 8.0. DNA concentration was determined by spectrophotometric analysis at 260nm.

Probe preparation:

5' and 3' oligonucleotide primers homologous with nucleotides 1 to 26 and 1967 to 1990 of Rx1 *pspA* (LSM 13 and LSM2, respectively) were used to amplify the full length *pspA* and construct probe LSM_{pspA}13/2 from Rx1 genomic DNA. 5' and 3'

oligonucleotide primers homologous to nucleotides 161 to 187 and nucleotides 1093 to 1117 (LSM 12 and LSM 6, respectively) were used to amplify the variable alpha-helical region to construct probe LSMpspA12/6. PCR generated DNA was purified by Gene Clean (Bio101 Inc., Vista, CA) and random prime-labeled with digoxigenin-11-dUTP using the Genius 1 Nonradioactive DNA Labeling and Detection Kit as described by the manufacturer (Boehringer Mannheim, Indianapolis, IN).

DNA electrophoresis:

For Southern blot analysis, approximately 10 μ g of chromosomal DNA was digested to completion with a single restriction endonuclease, (*Hind* III, *Kpn* 1, *EcoR* 1, *Dra* 1, or *Pst* 1) then electrophoresed on a 0.7% agarose gel for 16-18 hours at 35 volts. For PCR analysis, 5 μ l of product were incubated with a single restriction endonuclease, (*Bcl* 1, *BamH* 1, *Pst* 1, *Sac* 1, *EcoR* 1 *Sma* 1, and *Kpn* 1) then electrophoresed on a 1.3% agarose gel for 2-3 hours at 90 volts. In both case, 1 Kb DNA ladder was used for molecular weight makers (BRL, Gaithersburg, MD) and gels were stained with ethidium bromide for 10 minutes and photographed with a ruler.

Southern blot hybridization

The DNA in the gel was depurinated in 0.25N HCL for 10 minutes, denatured in 0.5M NaOH and 1.5M NaCl for 30 minutes, and neutralized in 0.5M Tris-HCl (pH 7.2), 1.5M NaCl and 1mM disodium EDTA for 30 minutes. DNA was transferred to a nylon membrane

(Micron Separations INC, MA) using a POSIBLOT pressure blotter (Stratagene, La Jolla, CA) for 45 minutes and fixed by UV irradiation. The membranes were prehybridized for 3 hours at 42°C in 50% formamide, 5X SSC, 5X Denhardt solution, 25mM sodium phosphate (pH 6.5), 0.5% SDS 3% (wt/vol) dextran sulfate and 500µg/ml of denatured salmon containing 45% formamide, 5X SSC, 1X Denhardt solution, 20mM sodium phosphate (pH 6.5), 0.5% SDS, 3% dextran sulfate, 250µg/ml denatured sheared salmon sperm DNA and about 20ng of heat-denatured diogoxigenin-labeled probe DNA. After hybridization, the membranes were washed twice in 0.1% SDS and 2X SSC for 3 minutes at room temperature. The membranes were washed twice to a final stringency of 0.1% SDS in 0.3X SSC at 65°C for 15 minutes. This procedure yields a stringency greater than 95 percent. The membranes were developed using the Genius 1 Nonradioactive DNA Labeling and Detection Kit as described by the manufacturer (Boehringer Mannheim, Indianapolis, IN). To perform additional hybridization with other probes, the membranes were stripped in 0.2N NaOH/0.1%SDS at 40°C for 30 minutes and then washed twice in 2X SSC.

Polymerase Chain Reaction (PCR):

5' and 3' primers homologous with the DNA encoding the N- and C-terminal ends of PspA (LSM13 and LSM2, respectively) were used in these experiments. Amplifications were made using Taq DNA polymerase, MgCl² and 10X reaction buffer obtained from Promega (Madison, WI). DNA used for PCR was prepared using the

method previously described in this paper. Reactions were conducted in 50ml volumes containing 0.2mM of each dNTP, and 1ml of each primer at a working concentration of 50mM. $MgCl_2$ was used at an optimal concentration of 1.75mM with 0.25 units of Taq DNA polymerase. Ten to thirty ng of genomic DNA was added to each reaction tube. The amplification reactions were performed in a thermal cycler (M.J. Research, Inc.) using the following three step program. Step 1 consisted of a denaturing temperature of 94°C for 2 minutes. Step 2 consisted of 9 complete cycles of a denaturing temperature of 94°C for 1 minute, an annealing temperature of 50°C for 2 minutes, and an extension temperature of 72°C for 3 minutes. Step 3 cycled for 19 times with a denaturing temperature of 94°C for 1 minute, an annealing temperature of 60°C for 2 minutes, and an extension temperature of 72°C for 3 minutes. At the end of the last cycle, the samples were held at 72°C for 5 minutes to ensure complete extension.

Band size estimation:

Fragment sizes in the molecular weight standard and in the Southern blot hybridization patterns were calculated from migration distances. The standard molecular sizes were fitted to a logarithmic regression model using Cricket Graph (Cricket Software, Malvern, PA). The molecular weights of the detected bands were estimated by entering the logarithmic line equation obtained by Cricket Graph into Microsoft Excel (Microsoft

Corporation, Redmond, WA) in order to calculate molecular weights based in migration distances observed in the Southern blot.

Table 15.

| Restriction Enzyme | Strains Examined | | | | | Restriction Fragments (sizes in kilobases) | |
|--------------------|------------------|------|------|------|-----|---|-----------|
| | MC25 | MC26 | MC27 | MC28 | RX1 | MC25-MC28 | RX1 |
| <i>Hind</i> III | + | + | + | + | + | 7.7, 3.6 | 9.1, 4.2 |
| <i>Kpn</i> I | + | + | + | + | + | 11.6, 10.6 | 10.6, 9.8 |
| <i>Eco</i> RI | + | | | | + | 8.4, 7.6 | 7.8, 6.6 |
| <i>Dra</i> I | + | | | | + | 2.1, 1.1 | 1.9, 0.9 |
| <i>Pst</i> I | + | | | | + | >14, 6.1 | 10.0, 4.0 |

Table 16. Penicillin Resistant Capsular Serogroup 6 Strains from Spain

| Isolate | Penicillin | MIC (µg/ml) | Year | Site | Hospital |
|---------|------------|-------------|------|--------|------------------|
| MC25 | | 1 | 1986 | sputum | Bellvitge |
| MC26 | | 4 | 1988 | ear | San Juan de Dios |
| MC27 | | 1 | 1988 | ear | San Juan de Dios |
| MC28 | | 2 | 1988 | ? | San Juan de Dios |

References

1. Crain M. J., W. D. Waltman II, J. S. Turner, J. Yother, D. E. Talkington, L. S. McDaniel, B. M. Gray and D. E. Briles. Pneumococcal surface protein A (PspA) is serologically highly variable and is expressed by all clinically important capsular serotypes of *Streptococcus pneumoniae*. Infect Immun 1990; 58:3293-3299.
2. Briles D. E., J. Yother and L. S. McDaniel. Role of pneumococcal surface protein A in the virulence of *Streptococcus pneumoniae*. Rev Infect Dis 1988; 10:S372-374.
3. McDaniel L. S., J. Yother, M. Vijayakumar, L. McGarry, W. R. Guild and D. E. Briles. Use of insertional inactivation to facilitate studies of biological properties of pneumococcal surface protein A (PspA). J Exp Med 1987; 165:381-394.
4. Crain M. J. Unpublished data.

5. Yother J. and D. E. Briles. Structural properties and evolutionary relationships of PspA, a surface protein of *Streptococcus pneumoniae*, as revealed by sequences analysis. J Bact 1992; 174:601-609.
6. Talkington D. F., D. L. Crimmins, D. C. Voellinger, J. Yother and D. E. Briles. A 43-kilodalton pneumococcal surface protein, PspA: isolation, protective abilities, and structural analysis of the amino-terminal sequence. Infect Immun 1991; 59:1285-1289.
7. McDaniel L. S., B. A. Ralph, D. O. McDaniel and D. E. Briles. Localization of protection-eliciting epitopes on PspA of *Streptococcus pneumoniae* between amino acid residues 192 and 260. Microb Pathogen 1994; 17:323-337.
8. Yother J., G. L. Handsome and D. E. Briles. Truncated forms of PspA that are secreted from *Streptococcus pneumoniae* and their use in functional studies and cloning of the PspA gene. J Bact 1992; 174:610-618.
9. McDaniel L. S., J. S. Sheffield, E. Swiatlo, J. Yother, M. J. Crain and D. E. Briles. Molecular localization of variable and conserved regions of *pspA*, and identification of additional *pspA* homologous sequences in *Streptococcus pneumoniae*. Microb Pathogen 1992; 13:261-269.
10. Waltman W. D. II, L. S. McDaniel, B. M. Gray and D. E. Briles. Variation in the molecular weight of PspA (Pneumococcal Surface Protein A) among *Streptococcus pneumoniae*. Microb Pathogen 1990; 8:61-69.

11. Munoz R., J. M. Musser, M. Crain, D. E. Briles, A. Marton, A. J. Parkinson, U. Sorensen and A. Tomasz. Geographic distribution of penicillin-resistant clones of *Streptococcus pneumoniae*: characterization by penicillin-binding protein profile, surface protein A typing, and multilocus enzyme analysis. *Clinic Infect Dis* 1992; 15:112-118.
12. Brooks-Walter A. and L. S. McDaniel. 1994. Unpublished data.
13. Talkington D. F., D. C. Voellinger, L. S. McDaniel and D. E. Briles. Analysis of pneumococcal PspA microheterogeneity in SDS polyacrylamide gels and the association of PspA with the cell membrane. *Microb Pathogen* 1992; 13:343-355.
14. McDaniel L. S., G. Scott, K. Widenhofer, Carroll and D. E. Briles. Analysis of a surface protein of *Streptococcus pneumoniae* recognized by protective monoclonal antibodies. *Microb Pathogen* 1986; 1:519-531.
15. Sheffield J. S., W. H. Benjamin and L. S. McDaniel. Detection of DNA in Southern Blots by Chemiluminescence is a sensitive and rapid technique. *Biotechniques* 1992; 12:836-839.
16. Briles D. E., M. J. Crain, B. M. Gray, C. Forman and J. Yother. A strong association between capsular type and mouse virulence among human isolates of *Streptococcus pneumoniae*. *Infect Immun* 1992; 60:111-116.

17. Smith P. K., R. I. Krohn , G. T. Hermanson, A. K. Mallia, F.-H. Gartner, M. D. Provenzano, E. K. Fujimoto, N. M. Goeke, B. J. Olson and D. C. Klenk. Measurement of Protein using Bicinchoninic Acid. Anal Biochem 1985; 150:76-85.
18. Meade H. M., S. R. Long, C. B. Ruvkin, S. E. Brown and F. M. Ausubel. Physical and genetic characterization of symbiotic and auxotrophic mutants of *Rhizobium melioli* induced by transposon Tn5 mutagenesis. J Bacteriol 1982; 149:114-122.
19. Silhavy T. J., M. L. Berman and L. W. Enquist. Experiments with gene fusions. Cold Springs Harbor: Cold Springs Harbor Laboratory. 1984.
20. Murray M. G. and W. F. Thompson. Rapid isolation of high molecular weight plant DNA. Nucl Acids Res 1980; 8:4321-4325.
21. Use of PCR to amplify pspAs and fragments of pspAs:
Example 5, infra.
22. Southern E. M. Detection of specific sequences among DNA fragments separated by gel electrophoresis. J Mol Biol 1975; 98:503-517.
23. Badenes M. J., and Dan E. Parfitt. Reducing background and interference on Southern Blots probed with nonradioactive chemiluminescent probes. BioTechniques 1994; 17:622-624.

EXAMPLE 5 - Southern blot analysis of *pspA*s and Fragments of *pspA*

In this example, Applicants used oligonucleotides derived from the DNA sequence of *pspA* of *S. pneumoniae* Rx1 both as hybridization probes and as primers in the polymerase chain reaction to investigate the genetic variation and conservation of the different regions of *pspA* and *pspA*-like sequences. The probes used ranged in size from 17 to 33 bases and included sequences representing the minus 35, the leader, the α -helical region, the proline-rich regions, the repeat regions, and the C-terminus. Applicants examined 18 different isolates representing 12 capsular and 9 PspA serotypes. The proline-rich, repeat, and leader, regions were highly conserved among *pspA* and *pspA*-like sequence.

In the previous Example, it was shown that strain Rx1 and most other strains of *S. pneumoniae* had two homologous sequences that could hybridize with probes encoding the N terminal and C terminal halves of PspA. This conclusion that these were separate sequences was supported by the fact that no matter which restriction enzymes was used there were always at least two (generally two sometimes three or four) restriction fragments of Rx1 and most other strains hybridized with the *pspA* probes. When the genome of Rx1 was digested with *HindIII* and hybridized with these, two *pspA*-homologous sequences were found to be in 4.0 and 9.1 kb fragments. Using derivative of Rx1 which had insertion mutations in *pspA*, it was possible to determine

that the 4.0 kb fragment contained the functional *pspA* sequence. The *pspA*-homologous sequence included within the 9.1 kb band was referred to as the *pspA*-like sequence. Whether or not the *pspA*-like sequences makes a product is not know, and none has been identified *in vitro*. Since *pspA*-specific mutants can be difficult to produce in most strains, and exist for only a limited number of pneumococcal isolates, this Example identifies oligonucleotide probes that could distinguish between the *pspA* and *pspA*-like sequences.

The purpose of this Example was to further define both the conserved and variable regions of *pspA*, and to determine whether the central proline-rich region is variable or conserved, and identify those domains of *pspA* that are most highly conserved in the *pspA*-like sequence (and ergo, provide oligonucleotides that can distinguish between the two). Oligonucleotides were used and are therefore useful as both hybridization probes and as primers for polymerase chain reaction (PCR) analysis.

Hybridization with oligonucleotide probes.

The oligonucleotides used in this study were based on the previously determined sequence of Rx1 PspA. Their position and orientation relative to the structural domains of Rx1 PspA are shown in Figure 7. The reactivity of these oligonucleotide probes with the *pspA* and *pspA*-like sequences was examined by hybridization with a *Hind*III digest of Rx1 genomic DNA (Table 17). As expected, each of the eight probes recognized the *pspA*-

containing 4.0 kb fragment of the *Hind*III digested Rx1 DNA. Five of the 8 probes (LSM1, 2, 3, 7, and 12) could also recognize the *pspA*-like sequence of the 9.1 kb band at least at low stringency. At high stringency four of the probes (LSM2, 3, 4 and 5) were specific for the 4.0 kb.

These 8 probes were used to screen *Hind*III digest of the DAN from 18 strains of *S. pneumoniae* at low and high stringency. For comparison to earlier studies each of the strains was also screened using a full-length *pspA* probe. Table 23 illustrates the results obtained with each strain at high stringency. Table 18 summarizes the reactivities of the probes with the strains at high and low stringency. Strain Rx1 is a laboratory derivative of the clinical isolate, D39. The results obtained with both strains were identical. They are listed under a single heading in Table 23 and are counted as a single strain in Table 28. Although AC17 and AC94 are related clinical isolates, they have distinguishable *pspAs* and are listed separately. All of the other strains represent independent isolates.

The only strain not giving at least two *pspA*-homologous *Hind*III fragments was WU2. This observation was expected since WU2 was previously shown to have only one *pspA*-homologous sequence and to give only a single *Hind*III fragment that hybridizes with Rx1 *pspA*. Even at high stringency 6 of the 8 probes detected more than one fragment in at least one of the 18

strains Tables 18 and 23. Probes LSM7, 10 and 12 reacted with DNA from a majority of the strains and detected two fragments in over 59% of the strains they reacted with. In almost every case the fragments detected by the oligonucleotide probes were identical in size to those detected by the full-length *pspA* probe. Moreover, the same pairs of fragments were frequently detected by probes from the 3' as well as the 5' half Rx1 *pspA*. These results are consistent with earlier findings that the pairs of *HindIII* fragments from individual isolated generally include two separate but homologous sequences, rather than fragments of a single *pspA* gene.

The differences in the frequency with which the oligonucleotides reacted with (at least one fragment) of the strains in the panel was significant at $P < 0.0001$ by 2 x 8 chi square). When the oligonucleotides were compared in terms of their ability to react with both fragments of each strain the P value was also < 0.0001 . Table 18 gives the percentage of strains reactive with each probe, the percentage in which only one fragment was reactive, and the percentage in which two (or more) fragments were reactive.

The last column in Table 18 give the ratio of strains that showed one reactive *HindIII* fragment at high stringency divided by the total number of reactive strains. In this column values of 1 were obtained with probes that only reacted with one band in each reactive strain. Such probes are assumed to be

those that are most specific for *pspA*. The lowest values were obtained with probes that generally see two bands in each strain. Such probes are assumed to be those that represent regions relatively conserved between the *pspA* and *pspA*-like sequences. At high stringency, probes LSM3 and LSM4 detected only a single *HindIII* fragment in the DNA of strains they reacted with. These findings suggested probes LSM3 and LSM4 were generally detecting alleles of *pspA* rather than the *pspA*-like sequence. The observation that the fragments detected by LSM3 or LSM4 were also detected by all of the other reactive probes, strengthened the conclusion that these probes generally detected the *pspA* rather than the *pspA*-like sequence. WU2 has only one *pspA*-homologous DNA sequence and secretes a serologically detectable PspA. The fact that LSM3 reacts with the single *HindIII* fragment of WU2 is consistent with the interpretation that LSM3 detects the *pspA* sequences. Sequences representing the second proline region (LSM1) and the C-terminus (LSM2) appeared to also be relatively specific for the *pspA* sequences since they were generally detected in only one of the *HindIII* fragments of each strain.

Oligonucleotides, LSM12, and LSM10 detected the most conserved epitopes of *pspA* and generally reacted with both *pspA*-homologous fragments of each strain (Table 18). LSM7 was not quite as broadly cross-reactive but detected two PspAs in 41% of strains including almost 60% of the strains it reacted with. Thus, sequences representing the leader, first proline region,

and the repeat region appear to be relatively conserved not only within *pspA* but between the *pspA* and *pspA*-like sequences. LSM3, 4, and 5 reacted with the DNA from the smallest fraction of strains of any oligonucleotide (29 - 35 percent), suggesting that the portion of *pspA* encoding the α -helical region is the least conserved region of *pspA*.

With two strains BG85C and L81905, the oligonucleotides detected more than two *HindIII* fragments containing *pspA*-homologous sequences. Because of the small size of the oligonucleotide probes and the absence of *HindIII* restriction sites within any of them, it is very unlikely that these multiple fragments were the results of fragmentation of the target DNA within the probed regions. In almost every case the extra oligonucleotides were detected at high stringency by more than one oligonucleotide. These data strongly suggest that at least in these two strains there are 3 or 4 sequences homologous to at least portions of the *pspA*. The probes most reactive with these additional sequences are those for the leader, the α -helical region and the proline rich region. The evidence for the existence of these additional *pspA*-related sequences was strengthened by results with BG58C and L81905 at low stringency where the LSM3 (α -helical) primer picked up the extra 1.2 kb band of L81905 (in addition to the 3.6 kb band) and the LSM7 (proline-rich) primer picked up the extra 3.2 and 1.4 kb bands (in addition to the 3.6 kb band) of BG58C.

Amplification of *pspA*

The utility of these oligonucleotides as PCR primers was examined by determining if they could amplify fragments of *pspA* from the genomic DNA of different pneumococcal isolates. Applicants attempted to amplify *pspAs* from 14 diverse strains of *S. pneumoniae* comprising 12 different capsular types using primers based on the Rx1 *pspA* sequence. Applicants observed that the 3' primer LSM2, which is located at the 3' end of *pspA*, would amplify an apparent *pspA* sequence from each of the 14 pneumococcal strains when used in combination with LSM1 located in the region of *pspA* encoding the proline-rich region (Table 19). LSM2 was also used in combination with four other 5' primers LSM1, 3, 7, 8 and 12. LSM8 is located 5' of the *pspA* start site (near the -35 region).

If a predominant sequence of the expected length was amplified that could be detected on a Southern blot with a full-length *pspA* probe, we assumed that *pspA* gene of the amplified DNA had homologous sequences similar to those of the *pspA* primers used. Based on these criteria the primer representing the α -helical sequence was found to be less conserved than the primers representing the leader, proline, and C-terminal sequences. These results were consistent with those observed for hybridization. The lowest frequency of amplification was observed with LSM8 which is from the Rx1 sequence 5' of the *pspA*

start site. This oligonucleotide was not used in the hybridization studies.

Further evidence for variability comes from differences in the sizes of the amplified *pspA* gene. The Example showed that when PCR primers LSM12 and LSM2 were used to amplify the entire coding region of *PspA*, PCR products from different pneumococcal isolates ranged in size from 1.9 and 2.3 kb (Table 20). The regions within *pspA* encoding the α -helical, proline-rich, and repeats were also amplified from the same isolates. As seen in Table 20, the variation in size of *pspA* appeared to come largely from variation in the size of *pspA* encoding encodes the α -helical region.

Using probes that consisted of approximately the 5' and 3' halves of *pspA* it has been determined that the portion of *pspA* that encodes the α -helical regions is less conserved than the portion of *pspA* that encodes the C-terminal half of the molecule. This Example show using 4 oligonucleotide probes from within each half of the DNA encoding *PspA*. Since a larger number of smaller probes were used, Applicants have been able to obtain a higher resolution picture of conserved and variable sequences within *pspA* and have also been able to identify regions of likely differences and similarities between *pspA* and the *pspA*-like sequences.

The only strains in which the *pspA* gene has been identified by molecular mutations are Rx1, D39 and WU2. Rx1 and

D39 apparently have identical *pspA* molecules that are the result of the common laboratory origin of these two strains. WU2 lacks the *pspA*-like gene. Thus, when most pneumococci are examined by Southern blotting using full length-*pspA* as a probe, it is not possible to distinguish between the *pspA* and *pspA*-like loci, since both are readily detected. A major aim of these studies was to attempt to identify conserved and variable regions within the *pspA* and *pspA*-like loci. A related aim was to determine whether probes based on the Rx1 *pspA* could be identified that would permit one to differentiate *pspA* from the *pspA*-like sequence. Ideally such probes would be based on relatively conserved portion of the *pspA* sequence that was quite different in the *pspA*-like sequence. A useful *pspA* specific probe would be expected to identify the known Rx1 and WU2 *pspA* genes and identify only a single *HindIII* fragment in most other strains. Two probes (LSM3 and LSM4) never reacted with more than one *pspA*-homologous sequence in any particular strain. Both of reacted with Rx1 *pspA* and LSM3 reacted with WU2 *pspA*. Each of these probes reacted with 4 of the other 15 strains. When these probes identified a band, however, the band was generally also detected by all other Rx1 probes reactive with that strain's DNA. Additional evidence that the LSM3 and LSM4 were restricted to reactivity with *pspA* was that they reacted with the same bands in all three non-Rx1 strains. Each probe identifies *pspA* in certain strains and even when used in combination they recognized *pspA* in

over 40 percent of strains. Probes for the second proline-rich region (LSM1) and the C-terminus of *pspA* (LSM2) generally, but not always, identified only one *pspA*-homologous sequence at high stringency. Collectively LSM1, 2, 3, and 4 reacted with 16 of the 17 isolates and in each case revealed a consensus band recognized by most to all of the reactive probes.

By making the assumption that in different strains the Rxl *pspA* probes are more likely to recognize *pspA* than the *pspA*-like sequences, it is possible to make some predictions about areas of conservation and variability within the *pspA* and *pspA*-like sequences. When a probe detected only a single *pspA*-homologous sequence in an isolate, it was assumed that it was *pspA*. If the probe detected two *pspA*-homologous sequences, it was assumed that it was reacting with both the *pspA* and *pspA*-like sequence. Thus, the approximate frequency with which a probe detects *pspA* can be read from Table 18 as the percent of strains where it detects at least one *pspA*-homologous band. The approximate frequency with which the probes detect the *pspA*-like sequence is the percent of strains in which two or more *pspA*-homologous band are detected.

Using these assumptions the most variable portion of portion of the *pspA* gene was observed to be the -35 region and the portion encoding α -helical region. The most conserved portion of *pspA* was found to be the repeat region, the leader and the proline rich region. Although only one probe from the region

was used, the high degree of conservation among the 10 repeats in the Rx1 sequence makes it likely that other probes for the repeat regions give similar results.

The portion of the *pspA*-like sequence most similar to Rx1 *pspA* was that encoding the leader sequence, the 5' portion of the proline rich region, and the repeat region, and those portions encoding the N-terminal end of the proline-rich and repeat regions. The repeat region of PspA has been shown to be involved in the attachment to PspA to the pneumococcal surface. The conservation of the repeat region among both *pspA* and *pspA*-like genes suggests that if is PspA-like protein is produced, that it may have a surface attachment mechanism similar to that of PspA. The need for a functional attachment site may explain the conservation of the repeat region. Moreover, the conservation in DNA encoding the repeat regions of the *pspA* and *pspA*-like genes suggests that the repeat regions may serve as a potential anti-pneumococcal drug target. The conservation in the leader sequence between *pspA* and the *pspA*-like sequence was also not surprising since similar conservation has been reported for the leader sequence of other gram positive proteins, such as M protein of group A streptococci. It is noteworthy, however, that there is little evidence at the DNA level that the PspA lead is shared by many genes other than PspA and the possible gene product of the *pspA*-like locus.

Although the region encoding the C-terminus of *pspA* (LSM12) or the 3' portion of the proline-rich sequence (LSM1) appear to be highly conserved within *pspA* genes, corresponding regions in the *pspA*-like sequences are either lacking, or very distinct from those in *pspA*. The reason for conservation at these sites is not apparent. In the case of the PspA, its C-terminus does not appear to be necessary for attachment, since mutants lacking the C-terminal 49 amino acids are apparently as tightly attached to the cell surface as those with the complete sequence. Whether these difference from *pspA* portends a subtle difference in the mechanism of attachment of proteins produced by these two sequences is unknown. If the C-terminal end of the *pspA*-like sequence, or the 3' portion of the proline-rich sequence in the *pspA*-like sequence are as conserved within the *pspA*-like family of genes as it is within *pspA*, then this region of *pspA* and the *pspA*-like sequence serve as targets for the development of probes to distinguish between all *pspA* and *pspA*-like genes.

With two strains, some of the oligonucleotide probes identified more than two *pspA*-homologous sequences. In the case of each of these strains, there was a predominant sequence recognized by almost all of the probes, and two or three additional sequences that were each recognized by at least two of the probes. One interpretation of the data is that there may be more than two *pspA*-homologous genes in some strains. The

significance of such sequences is far from established. It is of interest however, that although the additional sequences is far from established. It is of interest however, that although the additional sequences share areas of homology with the leader, α -helical, and proline region, they exhibited no homology with the repeat region of the C-terminus of *pspA*. These sequences, thus, might serve as elements that can recombine with *pspA* and/or the *pspA*-like sequences to generate sequence diversity.

Alternatively the sequences might produce molecules with very different C-terminal regions, and might not be surface attached. If these *pspA*-like sequences make products, however, they, like *PspA*, may be valuable as a component of a pneumococcal antigenic, immunological vaccine compositions.

Bacterial strains, growth conditions and isolation of chromosomal DNA.

S. pneumoniae strains used in this study are listed in Table 5. Strains were grown in 100 ml of Todd-Hewitt broth with 0.5% yeast extract at 37°C to an approximate density of 5×10^8 cells/ml. Following harvesting of the cells by centrifugation (2900xg, 10 minutes), the DNA was isolated as previously described and stored at 4°C in TE (10mM Tris, 1mM EDTA, pH 8.0).

Amplification of *pspA* sequences.

Polymerase chain reaction (PCR) primers, which were also used as oligonucleotide probes in Southern hybridizations, were designed based on the sequence of *pspA* from pneumococcal

strain Rx1. These oligonucleotides were obtained from Oligos Etc. (Wilsonville, OR) and are listed in Table 22.

PCRs were done with a MJ Research, Inc., Programmable Thermal Cycler (Watertown, MA) as previously described using approximately 10 ng of genomic pneumococcal DNA with appropriate 5' and 3' primer pair. The sample was brought to a total volume of 50 μ l containing a final concentration of 50mM KCl, 10mM Tris-HCl (PH 8.3), 1.5 mM MgCl₂, 0.001% gelatin, 0.5 mM each primer, 200mM of each deoxynucleotide triphosphate, and 2.5 U of Taq DNA polymerase. Following overlaying of the samples with 50 μ l of mineral oil, the samples were denatured at 94°C for 2 minutes. Then the samples were subjected to 10 cycles consisting of 1 minute at 94°C, 2 minutes at 50°C, and 3 minutes at 72°C followed by another 20 cycles of 1 minute at 94°C, 2 minutes at 50°C, and 3 minutes at 72°C followed by another 20 cycles of 1 minute at 94°C, 2 minutes at 60°C, and 3 minutes at 72°C. After all 30 cycles, the samples were held at 72°C for an additional 5 minutes prior to cooling to 4°C. The PCR products were analyzed by agarose gel electrophoresis.

DNA hybridization analysis.

Approximately 5 μ g of chromosomal DNA was digested with HindIII according to the manufacturer's instructions (Promega, Inc., Madison, WI). The digested DNA was electrophoresed at 35 mV overnight in a 0.8% agarose gels and then vacuum-blotted onto Nytran membranes (Schleicher & Schuell, Keene, NH).

Labeling of oligonucleotide with and detection of probe-target hybrids were both performed with the Genius System according to the manufacturer's instructions (Mannheim, Indianapolis, IN). All hybridizations were done for 18 hours at 42°C without formamide. By assuming that 1% base-pair mismatching results in a 1°C decrease in T_m designations of "high" and "low" stringency were defined by salt concentration and temperature of post-hybridization washes. Homology between probe and target sequences was derived using calculated T_m the established method. High stringency is defined as 90% or greater homology, and low stringency is 80-85% sequence homology.

Table 17: Hybridization of oligonucleotides with *Hind*III restriction fragments of Rx1 DNA.

| Oligonucleotide | Region | Stringency | |
|-----------------|-----------------|------------|----------|
| | | Low | High |
| LSM12 | Leader | N.D. | 4.0, 9.1 |
| LSM5 | α -helix | N.D. | 4.0 |
| LSM3 | α -helix | 4.0, 9.1 | 4.0 |
| LSM4 | α -helix | 4.0 | 4.0 |
| LSM7 | Proline | 4.0, 9.1 | 4.0, 9.1 |
| LSM1 | Proline | 4.0, 9.1 | 4.0, 9.1 |
| LSM10 | Repeats | N.D. | 4.0, 9.1 |
| LSM2 | C-terminus | 4.0, 9.1 | 4.0 |

Note. Values indicated are the sizes of restriction fragments expressed as kb.

**Table 18. Summary of Hybridization at High and Low Stringency of
8 Oligonucleotides with *Hind*III Restriction Fragments of
the 17 Pneumococcal Isolates Listed in Figure ?**

| Oligonucleotide | Percent with ≥ 1 band | | Percent with ≥ 2 bands | | Percent with 1 band | | 1 band/ ≥ 1 band | |
|-----------------|-------------------------------|------|--------------------------------|------|------------------------|------|--------------------------|------|
| | Low | High | Low | High | Low | High | Low | High |
| LSM12 | | 82 | | 59 | | 24 | | 0.29 |
| LSM5 | | 29 | | 18 | | 12 | | 0.40 |
| LSM3 | 65 | 35 | 41 | 0 | 24 | 35 | 0.36 | 1.00 |
| LSM4 | 35 | 29 | 0 | 0 | 35 | 29 | 1.00 | 1.00 |
| LSM7 | 94 | 71 | 71 | 41 | 24 | 29 | 0.25 | 0.42 |
| ✓ LSM1 | 100 | 65 | 53 | 12 | 47 | 53 | 0.47 | 0.82 |
| LSM10 | | 94 | | 59 | | 35 | | 0.37 |
| LSM2 | 88 | 53 | 41 | 12 | 47 | 41 | 0.53 | 0.78 |

Note, for all values listed all 17 strains were examined. If no value is listed, then no strains were examined.

Table 1. Amplification of Pneumococcal Isolates using the Indicated 5' Prime Combination with the 3' Primer LSM2 at the 3' end of *pspA*

| 5' Primer | Region | Nucleotide Position | Amplified/ Tested | Percent Amplified |
|-----------|-------------------|---------------------|-------------------|-------------------|
| LSM8 | - 35 | 47 to 70 | 2/14 | 14 |
| LSM12 | leader | 162 to 188 | 8/14 | 57 |
| LSM3 | α -helical | 576 to 598 | 3/14 | 21 |
| LSM7 | proline | 1093 to 1117 | 12/14 | 86 |
| LSM1 | proline | 1312 to 1331 | 14/14 | 100 |

Note, by 2x5 chi square analysis the different primers amplified different frequencies of *pspAs* ($P < 0.0001$). The tendency for there to be more amplification with the 3' most primers was significant at $P < 0.0001$.

Table 20 Size of amplified *pspA* fragments in kilobases

| <i>pspA</i> Region | Primer Pairs | number <i>pspAs</i> examined | Size | Range | S.D. |
|--------------------|--------------|---------------------------------|----------|-------|------|
| Full length | LSM12 + LSM2 | 9 | 1.9-2.3 | 0.4 | 0.17 |
| α -helical | LSM12 + LSM6 | 6 | 1.1-1.5 | 0.4 | 0.17 |
| Proline | LSM7 + LSM9 | 3 | 0.23 | 0 | 0 |
| Repeats | LSM1 + LSM2 | 19 | 0.6-0.65 | 0.05 | 0.01 |

Note: amplification was attempted with each set of primers on a panel of 19 different *pspAs*. Data is shown only for *pspAs* that could be amplified with the indicated primer pairs.

Table 3 Pneumococcal strains

| Strain | Relevant characteristics |
|----------|--|
| WU2 | Capsular type 3, PspA type 1 |
| D39 | Capsular type 2, PspA type 25 |
| R36A | Nonencapsulated mutant of D39, PspA type 25 |
| Rx1 | Nonencapsulated variant of R36A, PspA type 25 |
| DBL5 | Capsular type 5, PspA type 33 |
| DBL6A | Capsular type 6A, PspA type 19 |
| A66 12/2 | Capsular type 3, PspA type 13 |
| AC94 | Capsular type 9L, PspA type 0 |
| AC17 | Capsular type 9L, PspA type 0 |
| AC40 | Capsular type 9L, PspA type 0 |
| AC107 | Capsular type 9V, PspA type 0 |
| AC100 | Capsular type 9V, PspA type 0 |
| AC140 | Capsular type 9N, PspA type 18 |
| D109-1B | Capsular type 23, PspA type 12 |
| BG9709 | Capsular type 9, PspA type 0 |
| BG58C | Capsular type 6A, PspA type ND |
| L81905 | Capsular type 4, PspA type 25 |
| L82233 | Capsular type 14, PspA type 0 |
| L82006 | Capsular type 1, PspA type 0 |

98E

| Table 82 PCR primers. | |
|-----------------------|-----------------------------------|
| Primer | Sequence (5' to 3') |
| LSM1 | CCGGATCCAGCTCCTGCACCAAAAC |
| LSM2 | GCGCGTCGACGGCTTAAACCCATTACCATTTGG |
| LSM3 | CCGGATCCTGAGCCAGAGCAGTTGGCTG |
| LSM4 | CCGGATCCGCTCAAAGAGATTGATGAGTCTG |
| LSM5 | GCGGATCCCGTAGCCAGTCAGTCTAAAGCTG |
| LSM6 | CTGAGTCGACTGGAGTTTCTGGAGCTGGAGC |
| LSM7 | CCGGATCCAGCTCCAGCTCCAGAACTCCAG |
| LSM8 | GCGGATCCTTGACCAATATTTACGGAGGAGGC |
| LSM9 | GTTTTTGGTGCAGGAGCTGG |
| LSM10 | GCTATGGCTACAGGTTG |
| LSM11 | CCACCTGTAGCCATAGC |
| LSM12 | CCGGATCCAGCGTGCCTATCTTAGGGGCTGGTT |
| LSM13 | GCAAGCTTATGATATAGAAATTTGTAAC |

TABLE 23

| Hybridization at high stringency of eight different PspA probes with <i>Hind</i> III digests of 18 strains of <i>Streptococcus pneumoniae</i> | | | | | | | | | | | | | | | | | |
|---|----------|-----|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|
| Probe | Strain | | | | | | | | | | | | | | | | |
| | Rx1/D39 | WU2 | DBL5 | DBL6A | A66 | AC94 | AC17 | AC40 | AC107 | AC100 | AC140 | DC109 | BG9709 | BG58C | L81905 | L82233 | L82006 |
| FL-Rx1 | 4.0, 9.1 | 3.8 | 3.7, 5.8 | 3.0, 3.4 | 3.6, 4.3 | 3.6, 6.3 | 3.6, 6.3 | 3.2, 3.6 | 3.6, 6.3 | 4.0, 8.0 | 3.0, 4.0 | 3.3, 4.7 | 2.2, 9.6 | 1.4, 3.2 | 3.6, 5.2 | 3.7 | 4.3, 6.4 |
| LSM 12 | 4.0, 9.1 | 3.8 | 3.7, 5.8 | 3.0, 3.4 | 4.3 | | 3.6, 6.3 | 3.2, 3.6 | | 4.0, 8.0 | 4.0 | 3.3, 4.7 | 2.2, 9.6 | 1.4, 3.2 | 3.6 | 1.3, 3.7 | |
| LSM5 | 4.0 | | | | | 3.6, 6.3 | | | | | | | 2.2, 9.6 | 3.6 | 1.2, 2.3 | | |
| LSM3 | 4.0 | 3.8 | | | | 6.3 | | | | | | | 2.2 | 3.6 | 3.6 | | |
| LSM4 | 4.0 | | | | | | | | | | | | 2.2 | 3.6 | 3.6 | 3.7 | |
| LSM7 | 4.0, 9.1 | 3.8 | 3.7 | 3.0, 3.4 | 3.6 | | | 3.2, 3.6 | | | 3.0, 4.0 | 3.3, 4.7 | 2.2, 9.6 | 3.6 | 2.3 | 3.7 | |
| LSM1 | 4.0, 9.1 | 3.8 | 3.7, 5.8 | 3.4 | | 6.3 | | 3.2 | 3.6 | 4.0 | 4.0 | | 2.2 | | 5.2 | | |
| LSM 10 | 4.0, 9.1 | 3.8 | 3.7 | 3.4 | 3.6, 4.3 | | 3.6, 6.3 | 3.2 | 3.6, 6.3 | 4.0 | 4.0 | 3.3, 4.7 | 2.2, 9.6 | 3.6 | 3.6, 5.2 | 1.3, 3.7 | 4.3, 6.4 |
| LSM2 | 4.0 | | 3.7 | | | 3.6 | 3.6 | | 3.6, 6.3 | 4.0 | 3.0, 4.0 | 4.7 | | | | | 4.3 |

Note: All probes were tested versus *Hind*III digests of all strains. If no bands are listed none were detected. Strains Rx1 and D39 gave identical results and are shown in a single column. The full name of strain AC109 is AC109-1B

EXAMPLE 6 - Restriction Fragment Length Polymorphisms of *pspA*
Reveals Grouping

Pneumococcal surface A (PspA) is a protection eliciting protein of *Streptococcus pneumoniae*. The deduced amino acid sequence of PspA predicts three distinct domains; an α helical coiled-coil region, followed by two adjacent proline-rich regions, and ten 20 amino acid repeats. Almost all PspA molecules are cross-reactive with each other in variable degrees. However, using a panel of monoclonal antibodies specific for individual epitopes, this protein has been shown to exhibit considerable variability even within strains of the same capsular type. Oligonucleotide primers based on the sequence of *pspA* from *S. pneumoniae* Rx1 were used to amplify the full-length *pspA* gene and the 5' portion of the gene including the α -helical and the proline-rich region. PCR-amplified product were digested with *Hha* I or *Sau*3A I to visualize restriction fragment length polymorphism of *pspA*. Although strains were collected from around the world and represented 21 different capsular types, isolates could be grouped into 17 families or subfamilies based on their RFLP pattern. The validity of this approach was confirmed by demonstrating that *pspA* of individual strains which are known to be clonally related were always found within a single *pspA* family.

Numerous techniques have been employed in epidemiological surveillance of pneumococci which include

serotyping, ribotyping, pulsed field electrophoresis, multilocus enzyme electrophoresis, penicillin-binding protein patterns, and DNA fingerprinting. Previous studies have also utilized the variability of pneumococcal surface protein A (PspA) to differentiate pneumococci. This protein, which can elicit protective antipneumococcal antibodies, is a virulence factor found on all pneumococcal isolates. Although PspA molecules are commonly cross-reactive, they are seldom antigenically identical. This surface protein is the most serologically diverse protein known on pneumococci; therefore, it is an excellent marker to be used to follow individual strains. Variations in PspA and the DNA surrounding its structural gene have proven useful for differentiation of *S. pneumoniae*.

When polyclonal sera are used to identify PspA, cross-reaction is observed between virtually all isolates. Conversely, when panels of monoclonal antibodies are used to compare PspA of independent isolation they are almost always observed to express different combinations of PspA epitopes. A typing system based on this approach has limitations because it does not easily account for differences in monoclonal binding strength to different PspA molecules. Moreover, some strains are weakly reactive with individual monoclonal antibodies and may not always give consistent results.

A less ambiguous typing system that takes advantage of the diversity of PspA was therefore necessary to develop and was

used to examine the clonality of strains. This method involves examination of the DNA within and adjacent to the *pspA* locus. Southern hybridizations of pneumococcal chromosomal DNA digested with various endonucleases, such as *Hind* III, *Dra* I, or *Kpn* I, and probed with labeled *pspA* provided a means to study the variability of the chromosome surrounding *pspA*. When genomic DNA is probed, the *pspA* and the *pspA*-like loci are revealed. In most digests the *pspA* probe hybridizes to 2-3 fragments and, digests of independent isolates were generally dissimilar.

Like the monoclonal typing system, the Southern hybridization procedure permitted the detection of clones of pneumococci. However, it did not provide a molecular approach for following *pspA* diversity. Many of the restriction sites defining the restriction fragment length polymorphism (RFLP) were outside of the *pspA* gene, and it was difficult to differentiate the *pspA* gene from the *pspA*-like locus. In an effort to develop a system to follow *pspA* diversity Applicants examined the RFLP of PCR-amplified *pspA*. Amplified *pspA* was digested with *Sau*3A I and *Hha* I, restriction enzymes with four base recognition sites. To evaluate the utility of this approach *pspA* from clinical and laboratory strains known to be clonally related as well as random isolates were examined.

Bacterial strains

Derivatives of the *S pneumoniae* D39-Rx1 family were kindly provided by Rob Massure and Sanford Lacks (Figure 8).

Eight clinical isolates from Spain and four isolates from Hungary, a gift from Alexander Tomasz. Seventy-five random clinical isolates from Alabams, Sweden, Alaska, and Canada were also studied.

PCR amplifications

The oligonucleotide primers used in this study are listed in Table 24. Chromosomal DNA, which was isolated according to procedures described by Dillard et al., was used as template for the PCR reactions. Amplification was accomplished in a 50 μ l reaction containing approximately 50 ng template DNA, .25U Taq, 50 μ M of each primer, 175 μ M $MgCl_2$, and 200 μ M dNTP in a reaction buffer containing 10 μ M Tris-HCl, pH 9.0, 50 μ M KCl, 0.1% Triton X-100, 0.01% wt/vol. gelatin. The mixture was overlaid with mineral oil, and placed in a DNA thermal cycler. The amplification program consisted on an initial denaturation step at 94°C, followed by 29 cycles opf 94°C for 1 min, 55°C for 2 min, and 72°C for 3 min. The final cycle included an incubation at 72°C for 5 min.

Restriction fragment analysis of PCR-amplified product

Aliquots of the PCR mixtures were digested with *Hha* I or *Sau*3A I in a final volume of 20 μ l according to manufacturer's protocols. After digestion the DNA fragments were electrophoresed on a 1.3% TBE agarose gel and stained with ethidium bromide. Fragment sizes were estimated by comparison to a 1kb DNA ladder (Gibco BRL).

Because of the variability of *pspA*, and the fact that the entire *pspA* sequence is known for only one gene, it has not been possible to design primers which amplify *pspA* from 100% of pneumococcal strains. However, oligonucleotide primers, LSM2 and LSM1, can amplify an 800 bp region of the C-terminal end in 72 of the 72 strains tested. Based on hybridizations at different stringencies, this region was found to be relatively conserved in pneumococcal strains, and thus would not be expected to be optimal for following restriction polymorphisms within the *pspA* molecule. LSM13 and LSM2, primers which amplify the full length *pspA* gene, can amplify *pspA* from approximately 79% 55/75 of the strains tested (Table 25).

Stability of amplified RFLP pattern within clonally related pneumococci

To determine the stability of *pspA* during long passages in vitro, we examined the RFLP pattern of the *pspA* gene of the derivatives of the *S. pneumoniae* D39-Rx1 family. Rx1 is an acapsular derivative of *S. pneumoniae* D39, the prototypical pneumococcal laboratory strain isolated by Avery in 1914. Throughout the 1900's spontaneous and chemical mutations have been introduced into D39 by different laboratories (Figure 8). During this period unencapsulated strains were maintained in vitro, and D39 was passed both in vivo and in vitro passage. All the derivatives of D39, including Rx1, R6, RNC, and R36A, produced a 1.9 kb fragment upon PCR amplification of full length

pspA. All members of the family exhibited the RFLP pattern. Digestion with *Sau3A* I of PCR amplified full length *pspA* revealed a .83, .58, .36 and a .27 kb fragment in all of the D39-rX1 derivatives of the family. Digesting the full length *pspA* with *Hha* I resulted in bands which were .76, .47, .39, .35, and .12 kb (Figure 9 or Table 26).

The stability of *pspA* polymorphism was also investigated using pneumococcal isolates which had previously been shown to be clonally related by other criteria, including capsule type, antibiotic resistance, enzyme electrophoresis, and *PspA* serotype. Three sets of isolates, all of which were highly penicillin resistant, were collected from patients during an outbreak in Hungary and two separate outbreaks in Spain. PCR amplified full length *pspA* from the capsular type 19A pneumococcal strains from the outbreak in Hungary, DB18, DB19, DB20, and DB21, resulted in a band approximately 2.0 kb. After digesting full length *pspA* with *Hha* I, four fragments were visualized, .89, .48, and .28 kb. Digestion with *Sau3A* I yielded five fragments .880, .75, .35, .34, and .10kb. Capsule type 6B pneumococcal strains, DB1, DB2, DB3, and DB4, were obtained from an outbreak in Spain. Full length *pspA* from these strains were approximately 1.9 kb. Digestion of the PCR-amplified fragment with *Hha* I resulted in four fragments which were .83, .43, .33, and .28 kb. *Sau3A* I digestion yielded a .88, .75, .34, and .10 kb fragments. DB6, DB8, and DB9, which are capsular serotype 23F

strains, were isolated from a second outbreak in Spain. DB6, DB8, and DB9 had an amplified *pspA* product which was 2.0 kb. *Hha* I digested fragments were .90, .52, .34, and .30 kb and *Sau*3A I fragments were .75, .52, .39, .22, .20, and .10 kb in size (Figure 10). DB7 had a 19A capsular serotype and was not identical to DB6, DB8, and DB9. In the D39/Rx1 family and in each of the three outbreak families the size of the fragments obtained from the *Hha* I and the *Sau*3A I digests totaled approximately 2.0 kb which is expected if the amplified product represents a single *pspA* sequence.

Diversity of RFLP pattern of amplified *pspA* from random pneumococcal isolates

PCR amplification of the *pspA* gene from 70 random clinical pneumococcal isolates yielded full-length *pspA* ranging in size from 1.8 kb to 2.3 kb. RFLP analysis of PCR-derived *pspA* revealed two to six DNA fragments ranging in size from 100 bp to 1.9 kb depending on the strain. The calculated sum of the fragments never exceeded the size of the original amplified fragment. Not all pneumococcal strains had a unique *pspA*, and some seemingly unrelated isolates from different geographical regions and different capsular types exhibited similar RFLP patterns. Isolates were grouped into families based on the number of fragments produced by *Hha* I and *Sau*3A I digests and the relative size of these fragments.

Based on the RFLP patterns it was possible to identify 17 families with four of the families containing pairs of subfamilies. Within families all of the restriction fragments were essentially the same regardless which restriction enzyme was used. The subfamilies represent situations where two families share most but not all the restriction fragments. With certain strains an FRLP pattern was observed where detectable fragment size differed from the pattern of the established family by less than 100 bp. Since the differences were considered small compared to the differences in the fragment size and the number of fragments between families, they were not considered in family designation. The RFLP pattern of two isolates from six of the families is pictured in Figure 11, Table 27. These families were completely independent of the capsular type or the protein type as identified by monoclonal antibodies (Table 28 and 29).

Previous DNA hybridization studies have demonstrated that the *pspA* gene of different isolates are the most conserved in their 3' region of the gene and more variable in the 5' region of the gene. Thus, it seemed likely that the differences in the *pspA* families reflected primarily differences in the 5' end of the gene. To confirm this theory, the α helical and proline region of *pspA* was examined without the amino acid repeats. Nucleotide primers LSM13 and KSH2 were used to amplify this fragment which is approximately 1.6 kb. Examination of this region of *pspA* afforded two things.

This primer pair permitted amplification of 90% of the strains which is greater than the 75% of the strains which can be amplified with oligonucleotides which amplify the full length gene. Second, it allowed Applicants to examine if the original groupings which were based on the full length gene coincide with the fingerprint patterns obtained by looking at the 5' half of the gene.

Figure 12 contains the same strains which were examined in Figure 11 but the PCR products were amplified with SKH2 and LSM13. The RFLP patterns obtained from digestion of the amplified α helical and proline rich region confirms the original designated families. However, these primers amplify a smaller portion of the *psaA* and therefore the difference in the families is not as dramatic as the RFLP patterns obtained from the RFLP pattern of the full length gene.

The polymerase chain reaction has simplified the process of analyzing *pspA* gene and have provided a means of using *pspA* diversity to examine the epidemiology of *S. pneumoniae*. Because not all strains contained a unique fingerprint of *pspA*, RFLP patterns of *pspA* cannot be used alone to identify the clonality of a strain. These results indicate the RFLP of PCR-amplified *pspA* from pneumococcal strains in conjunction with other techniques may be useful for identifying the clonal relatedness among pneumococcal isolates, and that this pattern is stable over long passages in vitro.

These findings suggests that the population of *pspA* is not as diverse as originally believed. PCR-RFLP of *pspA* may perhaps represent a relatively simplistic technique to quickly access the variability of the gene within a population. Further, these findings enable techniques to diagnose *S. pneumoniae* via PCR or hybridization by primers on probes to regions of *pspA* common within groupings.

The sequence studies divide the known strains into several families based on sequence homologies. Sequence data demonstrates that there have been extensive recombinations occurring in nature within *pspA* genes. The net effect of the recombination is that the "families" identified by specific sequences differ depending upon which part of the *pspA* molecule is used for analysis. "Families" or "grouping" identified by the 5' half of the alpha-helical region, the 3' half of the α -helical region and the proline rich region are each distinct and differ slightly from each other. In addition there is considerable evidence of other diversity (including base substitutions and deletions and insertions in the sequences) among otherwise closely related molecules.

This result indicates that it is expected that there will be a continuum of overlapping sequences of PspAs, rather than a discrete set of sequences.

The findings indicate that there is the greatest conservation of sequence in the 3' half of the α -helical region and in the immediate 5' tip. Because the diversity in the mid half of the α -helical region is greater, this region is of little use in predicting cross-reactivity among vaccine components and challenge strains. Thus, the sequence of 3' half of the alpha-helical region and the 5' tip of the coding sequence are likely to be the critical sequences for predicting PspA cross-reactions and vaccine composition.

The sequence of the proline-rich region may not be particularly important to composition of a vaccine because this region has not been shown to be able to elicit

cross-protection even though it is highly conserved. The reason for this is presumably because antibodies to epitopes in this region are not surface exposed.

Based on our present sequences of 27 diverse *pspAs* we have found that there are 4 families of the 3' half of the α -helical region and 2-3 families of the very 5' tip of the α -helical region. Together these form 6 combinations of the 3' and 5' families. This approach therefore should permit us to identify a panel of *pspAs* with 3' and 5' helical sequences representative of the greatest number of different *pspAs*. See Fig 13.

Table 29 Relationship of Capsular type and RFLP family.

| RELATIONSHIP BETWEEN CAPSULAR TYPE AND RFLP FAMILY | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|--|--------------|---|---|---|---|---|----|----|---|---|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|--|--|--|--|---|---|---|--|
| | Capsule Type | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| pspA Family | 1 | 2 | 3 | 4 | 5 | 6 | 6A | 6B | 7 | 8 | 9A | 9L | 9N | 9V | 10 | 11 | 12 | 13 | 14 | 15 | 19 | 22 | 23 | 31 | 33 | 35 | ND | | | | | | | | |
| A | | | 3 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| B | | | 1 | 1 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| C | | | | | | 2 | 1 | 2 | | | | | | | | | | | | | | 2 | | | | | | | | | | | | | |
| D | | | 1 | | | | 1 | | | | | | | | | | | | | | | | | | | | | | | | | | | 1 | |
| DD | | | | 2 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| E | | | 1 | 2 | | 1 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| F | | | | | | 1 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| FF | | | 1 | | | 1 | | | | | | | | | 1 | 1 | | 1 | | | | | 3 | | | | | | | | | | 1 | | |
| G | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| H | | | 1 | | | 1 | | | 2 | 1 | | | | | | 1 | | | 1 | 1 | 1 | 1 | | | | | | | | | | | | | |
| I | | | | | | | | | | | 2 | | | 2 | 4 | | | | | | | | | | | | | | | | | | | | |
| II | | | | | 1 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| J | 2 | | | | | 2 | | | | | | 1 | 1 | | | | | 1 | 2 | | 2 | | | | | | | | | | | | 1 | | |
| K | | 1 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| KK | 1 | | | | | 1 | | 1 | | | | | | | | | | | 1 | | | | | | | | | | | | | 1 | | | |
| L | | | | | | | | | | | | | | | | | | | | | | 1 | | | | | | | | | | | | | |
| M | | | | | | | | | | | | | | | | | | | | | | | | 1 | | | | | | | | | | 1 | |
| MM | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

Table 24. Oligonucleotides used in this study.

| Designation | Sequence 5'-3' | Nucleotide position |
|-------------|---|---------------------|
| LSM2 | GCG CGT CGA CGG CTT AAA CCC ATT CAC CAT TGG | 1990 to 1967 |
| LSM1 | CCG GAT CCA GCT CCT GCA CCA AAA AC | 1312 to 1331 |
| LSM13 | GCA AGC TTA TGA TAT Aga AAT TTG TAA C | 1 to 28 |
| SKH2 | CCA CAT ACC GTT TTC TTG TTT CCA GCC | 1333 to 1355 |

Table 25 Amplification of *pspA* from a panel of 72 independent isolates* of *S. pneumoniae*.

| CAPSULE TYPE | NUMBER OF STRAINS EXAMINED | LSM13 AND LSM2 | LSM13 AND SKH2 |
|--------------|----------------------------|------------------------|------------------------|
| | | % OF STRAINS AMPLIFIED | % OF STRAINS AMPLIFIED |
| 1 | 3 | 100 | 100 |
| 2 | 1 | 100 | 100 |
| 3 | 8 | 50 | 87 |
| 4 | 6 | 67 | 100 |
| 5 | 1 | 100 | 100 |
| 6 | 7 | 29 | 86 |
| 6A | 2 | 100 | 100 |
| 6B | 6 | 100 | 100 |
| 7 | 2 | 50 | 100 |
| 8 | 1 | 100 | 100 |
| 9V | 3 | 100 | 100 |
| 9A | 2 | 100 | 100 |
| 9L | 1 | 100 | 100 |
| 9N | 3 | 100 | 100 |
| 10 | 1 | 100 | 100 |
| 11 | 2 | 50 | 100 |
| 12 | 2 | 0 | 100 |
| 13 | 1 | 100 | 100 |
| 14 | 4 | 0 | 75 |
| 15 | 2 | 50 | 50 |
| 19 | 5 | 100 | 100 |
| 22 | 3 | 33 | 100 |
| 23 | 1 | 100 | 100 |
| 33 | 1 | 0 | 100 |
| 35 | 1 | 0 | 100 |
| nd | 3 | 100 | 100 |

*Our strain collection contains several groups of isolates known to be previously to be clonal and collected for that purpose. The data reported in the table includes only one representative isolate from such clonal groups.

Table 36. Rx1-D39 derivatives

| ISOLATE | SIZE OF <i>Hha</i> I DIGESTS (Kb) | SIZE OF <i>Sau</i> 3A I DIGESTS (Kb) |
|---------|--------------------------------------|---|
| D39 | .76, .47, .39, .35, .12 | .83, .58, .36, .27 |
| Rx1 | .76, .47, .39, .35, .12 | .83, .58, .36, .27 |
| R800 | .76, .47, .39, .35, .12 | .83, .58, .36, .27 |
| R6 | .76, .47, .39, .35, .12 | .83, .58, .36, .27 |
| R61 | .76, .47, .39, .35, .12 | .83, .58, .36, .27 |
| R6X | .76, .47, .39, .35, .12 | .83, .58, .36, .27 |
| R36NC | .76, .47, .39, .35, .12 | .83, .58, .36, .27 |
| R36A | .76, .47, .39, .35, .12 | .83, .58, .36, .27 |

TABLE 33. Strain information and family designation of independent isolates.

| STRAIN | CAPSULE TYPE | PspA TYPE | FAMILY | SIZE OF <i>Hha</i> I FRAGMENTS | SIZE OF <i>Sau</i> 3A I FRAGMENTS |
|---------|-----------------|--------------|--------|-----------------------------------|--------------------------------------|
| BG9163 | 6B | 21 | C | 1.55, .35 | 1.05, .35, .22 |
| EF6796 | 6A | 1 | C | 1.5, .35 | 1.05, .35, .22 |
| EF5668 | 4 | 12 | DD | 1.25, .49, .32 | 1.0, .80, .35 |
| EF8616A | 4 | ND | DD | 1.25, .49, .32 | 1.0, .80, .35 |
| EF3296 | 4 | 20 | E | 1.0, .40, .33 | 1.15, .50, .34 |
| EF4135 | 4 | ND | E | 1.0, .40, .33 | 1.15, .50, .34 |
| BG7619 | 10 | ND | F | 1.3, .40, .29, .10 | .82, .76, .35 |
| BG7941 | 11 | ND | F | 1.3, .40, .29, .10 | .82, .76, .35 |
| BG7813 | 14 | 8 | H | 1.05, .70, .36 | .90, .77, .35 |
| BG7736 | 8 | ND | H | 1.05, .70, .36 | .90, .77, .35 |
| AC113 | 9A | ND | I | 1.4, .34, .28 | 1.2, .80 |
| AC99 | 9V | 5 | I | 1.4, .34, .28 | 1.2, .80 |

Table 2. Relationship of RFLP family and PspA type.

| RELATIONSHIP BETWEEN PSPA TYPE AND RFLP FAMILY | |
|--|---|
| | PspA Type |
| pspA FAMILY | 0 1 3 5 8 12 13 16 18 19 20 21 24 25 26 30 33 34 36 37 ND |
| A | 1 |
| B | 1 |
| C | 2 |
| D | 1 1 |
| DD | 1 1 |
| E | 2 |
| F | 1 |
| FF | 1 |
| G | 1 |
| H | 1 1 |
| I | 1 1 |
| II | 1 1 |
| J | 1 1 |
| K | 1 1 |
| KK | 1 |
| L | 1 |
| M | 1 |
| MM | 1 |

EXAMPLE 7 - Ability of PspA immunogens to protect against individual challenge strains

CBA/N or BALB cJ mice were given 1 injection of 0.5 - μ g PspA in CFA, followed 2 weeks later by a boost in saline, and challenged between 7 and 14 (average 10) days post boost. Control mice were administered a similar immunization regimen, except that the immunization came from an isogeneic strain unable to make PspA. The PspA was either full length, isolated from pneumococci or cloned full length or BC100 PspA, as little statistical significance has been seen in immunogenicity between full length PspA and BC100. The challenge doses ranged from about 10^3 to 10^4 pneumocci in inoculum, but in all cases the challenge was at least 100 times LD₅₀.

The results are shown in the following Tables 30 to 60, and the conclusions set forth therein.

From the data, it appears that an antigenic, immunological or vaccine composition can contain any two to seven, preferably three to five PspA, e.g., PspAs from R36A and BG9739, alone, or combined with any or all of PspAs from Wu2, Ef5668, and DB15. Note that surprisingly WU2 PspA provided better protection against D39 than did R36a/Rx1/D39, and that also surprisingly PspA from Wu2 protected better against BG9739 than did PspA from BG9739. Combinations containing R36A, BG9739 and WU2 PspAs were most widely protective; and therefore, a preferred composition can contain any three PspA, preferably

R36A, BG9739 and WU2. The data in this Example shows that PspA from varying strains is protective, and that it is possible to formulate protective compositions using any PspA or any combination of the PspAs from the eight different PspAs employed in the tests. Similarly, one can select PspAs on the basis of the groupings in the previous Example. Note additionally that each of PspA from R36A, BG9739, EF5668 and DBL5 are, from the data, good for use in compositions.

A note about use of medians rather than averages.

Applicants have chosen to express data as median (a non-parametric parameter) rather than averages because the times to death do not follow a normal distribution. In fact there are generally two peaks. One is around day 3 or 6 when most of the mice die and the other is at > 21 for mice that live. Thus, it becomes nonsensical to average values like 21 or 22 with values like 3 or 6. One mouse that lives out of 5 has a tremendous effect on such an average but very little effect on the median. Thus, the median becomes the most robust estimator of time to death of most of the mice.

TABLE 30

| Relative ability of different PspAs to Protect against each challenge strains of <i>S. pneumoniae</i> (Summary of statistically significant protection) | | | | | | | | | | | | | |
|--|------|------|--------|------------------|---------------|-------------------|--------|--------|--------|------------------|-----------------|---------------|-----------------|
| | | | | Vaccine PspA | | | | | | | | | |
| Challenge | Caps | PspA | pspA | R36A, Rx1,D39 | JD908/ WU2 | JS1020/ BG9739 | EF3296 | EF5668 | L81905 | JS5010.3 DBL5 | JS3020 DBL6A | All immune | best protect |
| Strain | type | type | family | K | a | b | E | DD | b | II | D | - | - |
| D39 | 2 | 25 | K | ++ | +++ | | | + | | | | ++ | +++ |
| WU2 | 3 | 1 | a | +++ | +++ | +++ | | +++ | +++ | +++ | +++ | +++ | +++ |
| A66 | 3 | 13 | a | +++ | +++ | +++ | | +++ | +++ | +++ | +± | +++ | +++ |
| EF10197 | 3 | 18 | M | +++ | | +++ | | | | | | +++ | +++ |
| ATCC6303 | 3 | 7 | a | +++ | | | | | | | | +++ | +++ |
| BG9739 | 4 | 26 | b | + | +++ | + | 0+ | 0 | +± | 0 | 0 | ++ | +++ |
| EF3296 | 4 | 20 | E | +± | +± | 0+ | | | | 0 | 0 | 0 | +± |
| EF5668 | 4 | 12 | DD | + | 0 | +++ | 0+ | +++ | 0+ | + | 0+ | ++ | +++ |
| L81905 | 4 | 23 | b | + | + | ++ | ++ | 0 | + | +± | +± | ++ | ++ |
| DBL5 | 5 | 33 | II | + | | + | | + | + | ++ | 0 | ++ | ++ |
| EF6796 | 6A | 1 | C | +++ | | | | | | | | +++ | +++ |
| DBL6A | 6A | 19 | D | +++ | +± | ++ | +± | +++ | +± | +± | +++ | ++ | +++ |
| BG9163 | 6B | 21 | C | +++ | | +++ | | | | | | +++ | +++ |
| BG7322 | 6B | 24 | C | +++ | +++ | +± | 0 | +++ | +± | +++ | +± | +++ | +++ |

Note: Empty cells indicate that no experiment has been done. Bold means significant at $P < 0.05$, Small font bold (+) means $0.02 \leq P < 0.05$. Large font bold means $P < 0.02$. For this table statistical significance refers to delay in time to death except as indicate in the (+) footnote below. When "all immune" showed significant protection against death but individual data cells did not, the result for "all immune" is presented under best protection on the assumption that if more mice were done in each data cell one or more of them would have exhibited significant protection against death.

+++ = statistically significant protection against death; $\geq 50\%$ protection from death

++ = statistically significant protection against death; $< 50\%$ protection from death

+± = statistically significant delay in death; ≥ 20 protection from death

+ = statistically significant delay in death; < 20 protection from death, (or significant protection against death but not a significant delay in death)

0++ = Not statistically delay in time to death; but $\geq 50\%$ protection from death

0+ = Not statistically delay in time to death; but > 1.5 day extension in median time to death or $\geq 20\%$ protection from death.

0 = No apparent extension in time to death or protection from death.

TABLE 31

| Relative ability of different PspAs to Protect against each challenge strains of <i>S. pneumoniae</i> (Expressed as Median days Alive post challenge) | | | | | | | | | | | | | |
|--|------|------|--------|------------------|--------------|-------------------|--------|--------|--------|------------------|-----------------|---------------|----------------|
| | | | | Vaccine PspA | | | | | | | | | |
| Challenge | Caps | PspA | PspA | R36A, Rx1.D39 | JD908 WU2 | JS1020/ BG9739 | EF3296 | EF3668 | L81905 | JS5010.3 DBL5 | JS3020 DBL6A | All immune | All control |
| Strain | type | type | family | K | a | b | E | DD | b | II | D | -- | -- |
| D39 | 2 | 25 | K | 4.5 | >21 | | | 4 | | | | 5 | 2 |
| WU2 | 3 | 1 | a | >21 | >21 | >21 | | >21 | >21 | >21 | >21 | >21 | 2 |
| A66 | 3 | 13 | a | >21 | >21 | >21 | | >21 | >21 | >21 | 4 | >21 | 2 |
| EF10197 | 3 | 18 | M | >21 | | >21 | | | | | | >21 | 2 |
| ATCC3303 | 3 | 7 | a | >21 | | | | | | | | >21 | 5 |
| BG9739 | 4 | 26 | b | 3 | >21 | 6 | 3 | 3 | 5, 13 | 2 | 2 | 3 | 2 |
| EF3296 | 4 | 20 | E | 5 | 5 | 4.5 | | | | 2 | 2 | 3 | 2 |
| EF3668 | 4 | 12 | DD | 6 | 2 | >21 | 13 | >21 | 4 | >21 | 5 | 6 | 3 |
| L81905 | 4 | 23 | b | 5 | 5 | 8 | 6 | 3 | 5 | 3 | 3.5 | 5 | 2 |
| DBL5 | 5 | 33 | II | 4 | | 3 | | 3 | 3.5 | 6 | 2 | 3.5 | 2 |
| EF6796 | 6A | 1 | C | >21 | | | | | | | | >21 | 1 |
| DBL6A | 6A | 19 | D | >21 | 8.5 | 13 | 9 | >21 | 8 | 12 | >21 | 12.5 | 5.5 |
| BG9163 | 6B | 21 | C | >21 | | >21 | | | | | | >21 | 8.5 |
| BG7322 | 6B | 24 | C | >21 | >21 | 14.5 | 6 | >21 | 12.5 | >21 | 11 | >21 | 7 |

Note: Bold denotes statistically significant extension of life at $P < 0.05$. Small font denotes $0.025P < 0.05$; large font denotes $P < 0.02$. Median times to death indicated as 8, >21, are situations where the median is not within a continuum of values. In those cases the numbers shown are those closest to the median. In these cases the values give are those closest to the calculated median. Fractional values such as 3.5, indicate that the median is halfway between two numbers, in this case 3 and 4. As indicated in the original data (S103B), some experiments were terminated prior to 21 days post infection. There is little reason to assume, however, that results would have been significantly effected by the early termination's since very few mice infected with the strains used in those studies, have ever been observed to die later than 10 or 15 days post challenge. For statistical purposes all mice alive at the end of experiments were assumed to have been completely protected, and for the sake of calculations all surviving mice were assigned values of >21.

TABLE 32

| Ability of different PspAs to Protect Against Each Challenge strain of <i>S. pneumoniae</i> (Expressed as increase in survival time in days) (A denotes $\geq 50\%$ immune mice alive) | | | | | | | | | | | | | |
|---|------|------|--------|------------------|--------------|-------------------|--------|--------|--------|------------------|-----------------|---------------|----------------|
| | | | | Vaccine PspA | | | | | | | | | |
| Challenge | Caps | PspA | pspA | R36A, Rx1,D39 | JD908 WU2 | JS1020/ BG9739 | EF3296 | EF5668 | L81905 | JS5010.3 DBL5 | JS3020 DBL6A | All immune | Best Result |
| Strain | type | type | family | K | a | b | E | DD | b | II | D | | |
| D39 | 2 | 25 | K | 2.5 | A | | | 2 | | | | 3 | A |
| WU2 | 3 | 1 | a | A | A | A | | A | A | A | A | A | A |
| A66 | 3 | 13 | a | A | A | A | | A | A | A | 2 | A | A |
| EF10197 | 3 | 18 | M | A | | A | | | | | | A | A |
| ATCC6303 | 3 | 7 | a | A | | | | | | | | A | A |
| BG9739 | 4 | 26 | b | 1 | A | 4 | 1 | 1 | 3, 11 | 0 | 0 | 1 | A |
| EF3296 | 4 | 20 | E | 3 | 3 | 2.5 | | | | 0 | 0 | 1 | 3 |
| EF5668 | 4 | 12 | DD | 3 | -1 | A | 10 | A | 1 | A | 2 | 5 | A |
| L81905 | 4 | 23 | b | 3 | 3 | 6 | 4 | 1 | 3 | 1 | 1.5 | 3 | 6 |
| DBL5 | 5 | 33 | II | 2 | | 1 | | 1 | 1.5 | 4 | 0 | 1.5 | 4 |
| EF6796 | 6A | 1 | C | A | | | | | | | | A | A |
| DBL6A | 6A | 19 | D | A | 3 | 7.5 | 3.5 | A | 2.5 | 6.5 | A | 7 | A |
| BG9163 | 6B | 21 | C | A | | A | | | | | | A | A |
| BG7322 | 6B | 24 | C | A | A | 7.5 | -1 | A | 5.5 | A | 4 | A | A |
| | | | | R36A | WU2 | BG9739 | EF3296 | EF5668 | L81905 | DBL5 | DBL6A | All | Best |

Note: Bold denotes statistically significant extension of life at $P < 0.05$. Small font denotes $0.02 \leq P < 0.05$; large font denotes $P < 0.02$. Median increases in survival listed as 3, 9 or 1, A denote groups where the median does not fall within a continuum of values. In these cases the values give are those closest to the calculated median. Fractional values such as 3.5, indicate that the median is halfway between two numbers, in this case 3 and 4.

TABLE 33

| Relative ability of different PspAs to Protect against each challenge strains of <i>S. pneumoniae</i> (expressed % alive at 21 days post challenge) | | | | | | | | | | | | | |
|---|------|------|--------|------------------|--------------|-------------------|--------|--------|--------|------------------|-----------------|---------------|----------------|
| | | | | Vaccine PspA | | | | | | | | | |
| Challenge | Caps | PspA | pspA | R36A, Rx1,D39 | JD908 WU2 | JS1020/ BG9739 | EF3296 | EF5668 | L81905 | JS5010.3 DBL5 | JS3020 DBL6A | All immune | All control |
| Strain | type | type | family | K | a | b | E | DD | b | II | D | — | |
| D39 | 2 | 25 | K | 38 | 60 | | | 30 | | | | 38 | 3 |
| WU2 | 3 | 1 | a | 100 | 100 | 100 | | 100 | 100 | 100 | 100 | 100 | 1.5 |
| A66 | 3 | 13 | a | 75 | 100 | 80 | | 75 | 100 | 60 | 20 | 76 | 5 |
| EF10197 | 3 | 18 | M | 100 | | 80 | | | | | | 90 | 0 |
| ATCC6303 | 3 | 7 | a | 100 | | | | | | | | 100 | 0 |
| BG9739 | 4 | 26 | b | 11 | 60 | 13 | 25 | 0 | 25 | 0 | 0 | 12 | 0 |
| EF3296 | 4 | 20 | E | 25 | 20 | 10 | | | | 0 | 0 | 8 | 0 |
| EF5668 | 4 | 12 | DD | 22 | 25 | 60 | 40 | 100 | 40 | 60 | 0 | 41 | 9 |
| L81905 | 4 | 23 | b | 10 | 0 | 31 | 40 | 0 | 0 | 14 | 0 | 14 | 0 |
| DBL5 | 5 | 33 | II | 10 | | 14 | | 0 | 0 | 29 | 0 | 4 | 0 |
| EF6796 | 6A | 1 | C | 100 | | | | | | | | 100 | 0 |
| DBL6A | 6A | 19 | D | 67 | 25 | 33 | 0 | 60 | 25 | 0 | 80 | 35 | 4 |
| BG9163 | 6B | 21 | C | 89 | | | | 80 | | | | 86 | 20 |
| BG7322 | 6B | 24 | C | 100 | 60 | 25 | 0 | 89 | 25 | 80 | 25 | 55 | 6 |

Bold, denotes statistically significant protection against death at $P < 0.05$. Bold small font, indicates significant protection against death at $0.02 \leq P < 0.05$. Bold large font, indicates significant protection against death at $P < 0.02$.

TABLE 4

| Relative ability of different PspAs to Protect against each challenge strain of <i>S. pneumoniae</i> (% protected from death at 21 days post challenge) | | | | | | | | | | | | | |
|--|------|------|--------|--------------|--------------|------------------|--------|--------|--------|------------------|-----------------|---------------|----------------|
| | | | | Vaccine PspA | | | | | | | | | |
| Challenge | Caps | PspA | pspA | Rx1, D39 | WU2 JD908 | BG9739 JS1020 | EF3296 | EF5668 | L81905 | DBL5 JS5010.3 | DBL6A JS3020 | All immune | Best result |
| Strain | type | type | family | K | a | b | E | DD | b | II | D | - | |
| D39 | 2 | 25 | K | 36 | 59 | | | 28 | | | | 36 | 59 |
| WU2 | 3 | 1 | a | 100 | 100 | 100 | | 100 | 100 | 100 | 100 | 100 | 100 |
| A66 | 3 | 13 | a | 71 | 100 | 79 | | 74 | 100 | 58 | 16 | 75 | 100 |
| EF10197 | 3 | 18 | M | 100 | | 80 | | | | | | 90 | 100 |
| ATCC6303 | 3 | 7 | a | 100 | | | | | | | | 100 | 100 |
| BG9739 | 4 | 26 | b | 11 | 60 | 13 | 25 | 0 | 25 | 0 | 0 | 12 | 60 |
| EF3296 | 4 | 20 | E | 25 | 20 | 10 | | | | 0 | 0 | 8 | 23 |
| EF5668 | 4 | 12 | DD | 14 | 18 | 36 | 34 | 100 | 34 | 56 | -10 | 35 | 100 |
| L81905 | 4 | 23 | b | 10 | 0 | 31 | 40 | 0 | 0 | 14 | 0 | 14 | 40 |
| DBL5 | 5 | 33 | II | 10 | | 14 | | 0 | 0 | 29 | 0 | 4 | 29 |
| EF6796 | 6A | 1 | C | 100 | | | | | | | | 100 | 100 |
| DBL6A | 6A | 19 | D | 66 | 22 | 30 | -4 | 58 | 22 | -4 | 79 | 33 | 79 |
| BG9163 | 6B | 21 | C | 86 | | 75 | | | | | | 83 | 86 |
| BG7322 | 6B | 24 | C | 100 | 57 | 22 | 0 | 88 | 22 | 79 | 22 | 52 | 100 |

Bold, denotes statistically significant protection against death at $P < 0.05$. Bold small font, indicates significant protection against death at $0.02 \leq P < 0.05$. Bold large font, indicates significant protection against death at $P < 0.02$.

% protected has been corrected for any survivors in the control mice.

% protected = $100 \times (\% \text{ alive in immune} - \% \text{ alive in control}) / (100 - \% \text{ alive in control})$. Thus, if there were any mice alive in the control animals, the calculated "% protected" is less than the observed "% alive" listed in the previous table. The only exceptions to this are if 100% of immunized mice lived. Negative numbers mean that less immunized mice lived than did control mice. Please note that none of these negative numbers are significant even though we are using a one tailed test.

TABLE 35

| Recommended Immunogens to Protection against the indicated challenge strains of <i>S. pneumoniae</i> Based on Protection Score Based on median days alive and percent protected (numbers refer to preference as a vaccine strain with respect to the indicated challenge strain, 1= best) | | | | | | | | | | | |
|---|------|------|--------|--------------|-----------|---------------|--------|--------|--------|---------------|--------------|
| | | | | Vaccine PspA | | | | | | | |
| Challenge | Caps | PspA | pspA | Rx1.D39 | WU2 JD908 | BG9739 JS1020 | EF3296 | EF5668 | L81905 | DBL5 JS3010.3 | DBL6A JS3020 |
| Strain | type | type | family | K | a | b | E | DD | b | II | D |
| D39 | 2 | 25 | K | 2 | 1 | | | 3 | | | |
| WU2 | 3 | 1 | a | 1 | 1 | 1 | | 1 | 1 | 1 | 1 |
| A66 | 3 | 13 | a | 2 | 1 | 2 | | 2 | 1 | 3 | 0 |
| EF10197 | 3 | 18 | M | 1 | | 2 | | | | | |
| ATCC6303 | 3 | 7 | a | 1 | | | | | | | |
| BG9739 | 4 | 26 | b | 3 | 1 | 2 | 3 | 3 | 2 | 0 | 0 |
| EF3296 | 4 | 20 | E | 1 | 1 | 2 | | | | 0 | 0 |
| EF5668 | 4 | 12 | DD | 0 | 0 | 2 | 3 | 1 | 0 | 2 | 0 |
| L81905 | 4 | 23 | b | 2 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| DBL5 | 5 | 33 | II | 2 | | 3 | | 0 | 3 | 1 | 0 |
| EF6796 | 6A | 1 | C | 1 | | | | | | | |
| DBL6A | 6A | 19 | D | 2 | 0 | 3 | 0 | 2 | 0 | 0 | 1 |
| BG9163 | 6B | 21 | C | 1 | | 1 | | | | | |
| BG7322 | 6B | 24 | C | 1 | 2 | 3 | | 1 | 3 | 1 | 3 |
| Number of #1's | | | | 7 | 5 | 3 | 1 | 3 | 2 | 3 | 2 |

Bold, denotes statistically significant protection against death at $P < 0.05$. Where more than one PspA were equally protective, the same values were given to each. Recommendations are based on days to death with % protection dividing ties, especially among those where greater than 50% of mice lived to 21 days. "0" indicates test were conducted but compared to the other PspAs this one is not recommended.

Conclusions:

Statistically significant protection against death with >50% protection; 11/14 of the strains = 79%

Statistically significant protection against death; 13 / 14 strains = 93%

Statistically significant extension of life in 14/14 or 100% of strains.

TABLE 36

| Best Choice for Vaccine Components as of 95/8/27 | | | | | | |
|---|--|------------------------|------------------------|------------------------|----------------------|------------------------|
| | Vaccine Component (cumulative strains protected) % maximally protected | | | | | |
| Criterion | 1 | 2 | 3 | 4 | 5 | 6 |
| ≥ #1 PspA for each challenge strain | R36A (7) 50% | WU2 (10) 71% | BG9739* (11) 79% | EF5668 (12) 86% | DBL5 (13) 93% | DBL6A (14) 100% |
| ≥ #2 PspA for each challenge strain | R36A (12) 86% | BG9739 (12) 100% | | | | |
| Max score (+) type score | R36A (9) 64% | WU2 (11) 79% | BG9739 (13) 92% | DBL5 (14) 100% | | |
| Max Increase in Days alive | R36A (9) 64% | WU2 (11) 79% | BG9739 (13) 92% | DBL5 (14) 100% | | |
| % protected | R36A (7) 50% | WU2 (10) 64% | DBL5 (11) 79% | EF5668 (12) 86% | DBL6A (13) 92% | EF3296 (14) 100% |
| Theoretical mixture based on a few testable assumptions (see below) | R36A (10) 64% | BG9739 (12) 86% | DBL5 (13) 92% | EF3296 (14) 100% | | |

* This is not a unique combination. See table below.

TABLE 37

| Combinations where all Challenge Strains have a Vaccine strain with a score of ≥ #2 | | | | |
|---|---------------------|----------------------|-----------|-------------------|
| Number of PspAs in Combination | Combination | Number of #1 strains | Total #1s | Total #1s and #2s |
| 2 | R36A + BG9739 | 8 | 10 | 20 |
| 3 | R36A + BG9739 + WU2 | 11 | 15 | 25 |
| 3 | R36A + WU2 + DBL5 | 11 | 15 | 21 |
| 3 | R36A + WU2 + EF5668 | 11 | 15 | 23 |
| 3 | R36A + WU2 + DBL5 | 11 | 15 | 22 |

Pooled Data for Protection against D39 by various PspAs;
Days alive for each mouse

TABLE 38

| Exp. | Log CFU D39 | Mice | Days to Death/ immunogen | | | | |
|----------------|-------------|--------|------------------------------------|-------------|-----------------------|---|------------------------------------|
| | | | Rx1/R36A D39 | JD908 (WU2) | EF5668 | All Immune | control |
| I43 | 4.5 | CBA/N | | | 1,1,2,2,2 | | 1,1,2,2,3 |
| E145 | 4.0 | CBA/N | 2,3,3,3,4 | | | | 1,1,2,3,4 |
| E028 BCG | 5.93 | BALB/c | 3, 3x >21 | | | | 2,2,2,4 |
| E143 | 3.0 | CBA/N | | | 2,6,3x>10 | | 3,3,3,5,5 |
| E140 BC100 | 2.81 | CBA/N | 4,4,5,7,15 | | | | 2,2,2 |
| E169 | 2.7 | CBA/N | 2, 4x >21 | 2,5,3x >21 | | | 1,2,2,2,3 |
| E154 | 2.6 | CBA/N | 2,2,3,2x >21 | | | | 4x 2, 5, >21 |
| All ≤ 3.0 | | | 2,3,3,3,4,4,4,5,7,15 | | 1,1,2,2,2,2 | | 4x 1, 6x 2, 3,3,4 |
| All | | | 4x 2, 5x 3, 3x 4, 5, 7, 15, 9x >21 | 2,5,3x >21 | 1,1,2,2,2,2,6, 3x >21 | 1,1,9x 2, 5x 3, 3x 4, 5,5,6,7,15, 15x >21 | 5x 1, 16x 2, 6x 3, 4, 4, 5,5,5,>21 |

Pooled Data for Protection against D39 by various PspAs
Median Days Alive & alive: dead
with corresponding P values.

TABLE 39

| Exp. | Log CFU D39 | Mice | Rx1/R36A D39 | | JD908 (WU2) | | EF5668 | | All Immune | | Control | |
|----------------|-------------|--------|--------------|------|-------------|-------|---------|------|------------|-------|----------|------|
| | | | med | a:d | med | a:d | med | a:d | med | a:d | med | a:d |
| I43 | 4.5 | CBA/N | | | | | 2 | 0:5 | | | 2 | 0:5 |
| E145 | 4.0 | CBA/N | 3 | 0:5 | | | | | | | 2 | 0:5 |
| E028 BCG | 5.93 | BALB/c | >21 | 3:1 | | | | | | | 2 | 0:4 |
| E143 | 3.0 | CBA/N | | | | | >21 | 3:2 | | | 3 | 0:5 |
| E140 BC100 | 2.81 | CBA/N | 5 | 0:5 | | | | | | | 2 | 0:3 |
| E169 | 2.7 | CBA/N | >21 | 4:1 | >21 | 3:2 | | | | | 2 | 0:5 |
| E154 | 2.6 | CBA/N | 3 | 2:3 | | | | | | | 2 | 1:5 |
| All ≤ 3.0 | | | 4 | 0:10 | | | 2 | 0:5 | | | 2 | 0:13 |
| All | | | 4.5 | 9:15 | >21 | 3:2 | 4 (2.4) | 3:7 | 5 | 15:24 | 2 | 1:32 |
| | | | .0057 | .001 | .006 | .0045 | n.s. | .034 | .0001 | .0002 | | |
| | | | ++ | ++ | +++ | +++ | + | + | + | ++ | | |
| % alive | | | 38 | | 60 | | 30 | | 38 | | 3 | |
| | | | 36 | | 59 | | 28 | | 36 | | | |
| | | | Rx1/D39 | | WU2 | | EF5668 | | All immune | | controls | |

TABLE 40

| Pooled Data for Protection against WU2. by various PspAs | | | | | | | | | | | | |
|--|-------------------|--------|--------------------------|--------------|----------------|--------------------|-----------------|--------|-----------------|---------------|-------------------|--|
| Exp. | CFU WU2 | Mice | Days to Death/ immunogen | | | | | | | | | |
| | | | FL-R36A | Rx1 BC100 | JD108 (WU2) | JS1020 (BG9739) | BG9739 bc100 | EF5668 | L81905 bc100 | DBL5 bc100 | JS3020 (DBL6A) | control |
| Dr. Ed. expt. | | | | | | | | | | | +++ | |
| lots of prior expts. | | | +++ | | | | | | | | | |
| E012 | 3.0 | CBA/N | 15x >21 | | | | | | | | | 1,1, 11x 2, 7x 3, 4 |
| E028 | 6.01 | BALB/c | 4x >21 0.05/n.s. | | | | | | | | | 4, 6, 6, >21 |
| E084 | 3.75 ¹ | CBA/N | | | | 3x >15 | | | | | | 1,2,2,2,3,3, >15 |
| E125 bc100 | 3.57 | CBA/N | | | | | 4x >21 | | 4x >21 | 4x >21 | | 2,2,3,3,3, >21 |
| E129 | 3.18 | CBA/N | | | | 5x >23 | | | | | | 2,2,2,2,3 |
| E140 BC100 | 3.43 | CBA/N | | 4x >21 | | | | | | | | 1, 5x 2, 3, 4 |
| E143 | 3.0 | CBA/N | | | | | | 8x >10 | | | | 1,1,2,2,2,3 |
| E144 | 3.9 | CBA/N | | | | | | | | | 5x >21 | 5x 2 |
| E172 | 3.98 | CBA/N | | | 5x >21 | | | | | | | 5x 3 |
| All | | | 19x >21 | 4x >21 | 5x >21 | 8x >21 | 4x >21 | 8x >21 | 4x >21 | 4x >21 | 5x >21 | 6x 1, 3, 3 x 2, 20x 3, 4,4,4,6,6, >21 |
| All Immune | | | 61x >21 | | | | | | | | | |

| Pooled Data for Protection against WU2 by various PspAs | | | | | | | | | | | | |
|---|-------------------|--------|--|-----------------------------|-----------------------------|-----------------------------|-----------------------------|-----------------------------|-----------------------------|-----------------------------|-----------------------------|------------------------|
| Exp. | CFU | Mice | Median days Alive Alive : Dead P value based on Alive : Dead P value calculated compared to pooled controls (in this case 65 control mice) Score | | | | | | | | | |
| | | | FL-R36A | Rx1 BC100 | JD108 (WU2) | JS1020 (BG9739) | BG9739 bc100 | EF3668 | L81905 bc100 | DBL5 bc100 | JS3020 (DBL6A) | control |
| | | | | | | | | | | | | |
| | | | | | | | | | | | | |
| WU2 | | | | | | | | | | | | |
| Dr. Ed. expt. | | | | | | | | | | | +++ | |
| lots of prior expts. | | | +++ | | | | | | | | | |
| E012 | 3.0 | CBA/N | >21 15:0 | | | | | | | | | 1,1, 11x 2, 7x 3, 4 |
| E028 | 6.01 | BALB/c | 4x >21 | | | | | | | | | 4, 6, 6, >21 |
| E084 | 3.75 ¹ | CBA/N | | | | 3x >15 | | | | | | 1,2,2,2,3,3, >15 |
| E125 bc100 | 3.57 | CBA/N | | | | | 4x >21 | | 4x >21 | 4x >21 | | 2,2,3,3,3, >21 |
| E129 | 3.18 | CBA/N | | | | 5x >23 | | | | | | 2,2,2,2,3 |
| E140 BC100 | 3.43 | CBA/N | | 4x >21 | | | | | | | | 1, 5x 2, 3, 4 |
| E143 | 3.0 | CBA/N | | | | | | 8x >10 | | | | 1,1,2,2,2,3 |
| E144 | 3.9 | CBA/N | | | | | | | | | 5x >21 | 5x 2 |
| E172 | 3.98 | CBA/N | | | 5x >21 | | | | | | | 5x 3 |
| All | | | >21 19:0 <.0001 +++ | >21 4:0 <.0001 +++ | >21 5:0 <.0001 +++ | >21 8:0 <.0001 +++ | >21 4:0 <.0001 +++ | >21 8:0 <.0001 +++ | >21 4:0 <.0001 +++ | >21 4:0 <.0001 +++ | >21 4:0 <.0001 +++ | 2 1:64 |
| % alive | | | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 2 |
| | | | | | | | | | | | | |
| | | | FL-R36A | Rx1 BC100 | JD108 (WU2) | JS1020 (BG9739) | BG9739 bc100 | EF5668 | L81905 bc100 | DBL5 bc100 | JS3020 (DBL6A) | control |

TABLE 41

| 7/12 Challenge | days of death | median days of death | alive : dead | P value based on days to death | P value based on alive : dead | Score | % alive | % prot. |
|-------------------|--|-------------------------|-----------------|--------------------------------------|-------------------------------------|-------|------------|------------|
| All immune | 61x >21 | >21 | 61:0 | <.0001 | <.0001 | +++ | 100 | 100 |
| All controls | 6x 1, 33x 2, 20x 3, 4, 4, 6, 6, >21 | 2 | 1:64 | | | | 2 | 2 |

TABLE 42

| Pooled Data for Protection against A66. by various PspAs | | | | | | | | | | | | | | |
|--|------------|-------|----------------------------|------------------|--------------------|--------------------|-------------------|-----------------|--------------------|------------------|------------------------|------------------|-------------------|--|
| Exp. | CFU A66 | Mice | Days to Death/ immunogen | | | | | | | | | | | |
| | | | FL-R36A/ D39 | Rx1 BC100 | JD908 (WU2) | JS1020 (BG9739) | BG9739 bc100 | EF5668 | L81905 FL | L81905 bc100 | JS5010.3 FL (DBL5) | DBL5 bc100 | JS3020 (DBL6A) | control |
| E169 | 2.60 | CBA/N | 5x >21 | | 5x >21 | | | | | | | | | 1,1,2,2,6 |
| E152 bc100 | 2.78 | CBA/N | | | | | 4x >21 | | | 4x >21 | | 4x >21 | | 3x 2, 3, 6, 6 >21 |
| E104 | 3.0 | CBA/N | | | | 2,8,3x >21 | | | | | 3,4,4,2x >21 | | 2,4,4,5,>21 | 2,2,2,2,3 |
| E143 | 3.0 | CBA/N | | | | | | 4, 4x >10 | | | | | | 2,2,3,3 |
| E140 | 3.43 | CBA/N | | 4x >21 | | | | | | | | | | 1,1,1 |
| E172 | 3.94 | CBA/N | | | | | | | 5x >21 | | | | | |
| E145 | 3.97 | CBA/N | 13, 4x >21 | | | | | | | | | | | 1, 2, 2, 2, 4 |
| E121 | 4.16 | CBA/N | 3x 3, 2x 4, 5x >21 | | | | | | | | | | | 1, 8x 2, >21 |
| All | | | 3x 3, 2x 4, 13, 14x >21 | 4x >21 | 5x >21 | 2,8,3x >21 | 4x >21 | 4, 4x >21 | 5x >21 | 4x >21 | 3,4,4,2x >21 | 4x >21 | 2,4,4,5,>21 | 7x 1, 22x 2, 3x 3, 4, 3x 6, 2x >21 |
| median; A: D | | | >21 14:6 | >21 4:0 | >21 5:0 | >21 3:2 | >21 5:0 | >21 4:1 | >21 5:0 | >21 4:0 | 4 2:3 | >21 4:0 | 4 1:4 | 2 2:36 |
| P values | | | <0.0001 <0.0001 | 0.0002 0.0001 | <0.0001 <0.0001 | 0.004 0.0075 | 0.0002 <0.0001 | 0.0006 0.006 | <0.0001 <0.0001 | 0.0002 0.0001 | 0.0025 n.s. | 0.0002 0.0001 | 0.015 n.s. | |
| Mini Pools | | | R36A/Rx1/WG44.1 | JD908 | | BG9739 | EF5668 | | L81905 | | DBL5 3,4,4,4,6x >21 | | DBL6A | Control |
| | | | >21 18:6 | >21 5:0 | | >21 8:2 | >21 4:1 | | >21 9:0 | | >21 6:4 | | 4 1:4 | 2 2:36 |
| P values rank/a:d | | | <0.0001 | <0.0001 | <0.0001 | <0.0001 | 0.0006 0.006 | | <0.0001 | | 0.0004 | | 0.015 n.s. | |
| Score | | | +++ | +++ | +++ | +++ | +++ | +++ | +++ | +++ | +++ | +++ | ++ | |
| % alive | | | 72 71 | 100 100 | | 80 79 | 75 74 | 100 100 | | 60 58 | | 20 16 | | 5 0 |
| A66 challenge | | | R36A/Rx1/WG44.1 | JD908 | | BG9739 | EF5668 | | L81905 | | DBL5 | | DBL6A | |

| A66 challenge | days of death | median days alive | alive : dead | P - days to death | P - alive : dead | Score | % alive | % protected |
|---------------|--|----------------------|-----------------|----------------------|---------------------|-------|------------|----------------|
| All immune | 2, 2, 4x 3, 7x 4, 5, 8, 13, 50x >21 | >21 | 50:16 | <0.0001 | <0.0001 | +++ | 76 | 75 |
| All controls | 7x 1, 22x 2, 3x 3, 4, 3x 6, 2x >21 | 2 | 2:36 | | | | 5 | 0 |

TABLE 43

| Pooled Data for Protection against EF10197. by various PspAs | | | | | | | | | |
|--|--------------------|-------|--------------------------|--------------------|--------|-------------------|--------|------------------------|-----------|
| Exp. | CFU EF 10197 | Mice | Days to Death/ immunogen | | | | | | |
| | | | Rx1 BC100 | JS1020 (BG9739) | L81905 | JS3020 (DBL6A) | EF5668 | JS5010.3 FL (DBL5)0 | control |
| E140 | 3.00 | CBA/N | 5x >21 | | | | | | 2,2,2 |
| MI BCG | 2.70 | CBA/N | * | | | | | | 2,2,2,2,2 |
| E129 | 3.34 | CBA/N | | 8, 4x >23 | | | | | 2,2,2,2,9 |

* This was a passive protection study. Its controls have been included to increase the numbers of control mice.

TABLE 44

| Pool of Pools for protection against EF10197 | | | | | |
|--|-------------------------------|--------------------------------|-----------------------------|-------------|----------------------------|
| line | Group Description | Delay in death and/or survival | | Survival | |
| | | days to death (median) | P values etc. | alive: dead | P values etc. |
| 1a | Rx1 (E140) | 5x >21 | 0.017 vs 1b 0.0013 vs 4b | 5:0 | 0.018 vs 1b 0.008 vs 4b |
| 3a | JS1020 (E129) | 8, 4x >23 | 0.0007 vs 3b | 4:1 | 0.024 vs 3b |
| 4a | all immune | 8, 9x >21 | <0.0001 vs 4b | 9:1 | 0.0002 vs 4b |
| 1b | Rx1 controls (E140) | 2,2,2 | | 0:3 | |
| 2b | MI BCG | 2,2,2,2,2 | | 0:5 | |
| 3b | JS1020 cont. (E129) | 2,2,2,2,9 | | 0:5 | |
| 4b | all controls (without MI BCG) | 2,2,2,2,2,2,9 | | 0:8 | |

TABLE 45

| Summary of protection against EF10197 | | | | | | | |
|---------------------------------------|-------------|---------|-------------|------------|--------------|---------------|--------|
| Immunogen | alive: dead | % alive | % protected | median DOD | P time alive | P alive: dead | Score* |
| Rx1 | 5:0 | 100 | 100 | >21 | 0.017 | 0.018 | +++ |
| JS1020 | 4:1 | 80 | 80 | >21 | 0.0007 | 0.024 | +++ |
| all immune | 9:1 | 90 | 90 | >21 | <0.0001 | 0.0002 | +++ |
| all controls | 0:8 | 0 | 0 | 2 | -- | -- | -- |

* +++ = statistically significant protection against death with ≥50% protected.

TABLE 46

| Pooled Data for Protection against ATCC6303, by various PspAs | | | | | | | | |
|---|---------------|-------|--------------------------|-----------------|--------|----------------|--------|-----------------------------|
| Exp. | CFU ATCC 6303 | Mice | Days to Death/ immunogen | | | | | |
| | | | Rx1 BC100 | JS1020 (BC9739) | LS1905 | JS3020 (DBL6A) | EF3668 | JS3010.3 FL (DBL5)0 control |
| E140 | 2.30 | CBA/N | 5x >21 | | | | | 4, 4x 5 |
| E129 | 3.80 | CBA/N | | n.v. | | | | |

TABLE 47

| Pool of Pools for protection against ATCC6303 | | | | | |
|---|---------------------|--------------------------------|---------------|-------------|---------------|
| line | Group Description | Delay in death and/or survival | | Survival | |
| | | days to death (median) | P values etc. | alive: dead | P values etc. |
| 1a | Rx1 (E140) | 5x >21 (>21) | 0.0040 | 5:0 | 0.004 |
| 1b | RX1 controls (E140) | 4, 4x 5 | 5 | 0:5 | -- |

| Summary of protection against ATCC6303 | | | | | | | |
|--|-------------|---------|-------------|------------|--------------|---------------|--------|
| Immunogen | alive: dead | % alive | % protected | median DOD | P time alive | P alive: dead | Score* |
| Rx1 | 5:0 | 100 | 100 | >21 | 0.004 | 0.004 | +++ |
| Rx1 controls | 0:5 | 0 | 0 | 5 | -- | -- | -- |

* +++ = statistically significant protection against death with ≥50% protected.

TABLE 48

TABLE 49

| Pooled Data for Protection against BG9739. by various FL PspAs | | | | | | | | | | | | | |
|--|---------------|--------|--------------------------|----------------|----------------|----------------------------------|-------------------|-----------------|--------------|-------------------|--------------------------|-----------------|---|
| Exp. | CFU BG9739 | Mice | Days to Death/ immunogen | | | | | | | | | | |
| | | | R36A FL | BC100 (Rx1) | JD908 (WU2) | JS1020 (BG9739) | bc100 (BG9739) | EF3296 FL | EF5668 FL | bc100 (L81905) | JS5010.3 FL (DBL5) | bc100 (DBL5) | JS3020 (DBL6 A) |
| E140 | 2.76 | CBA/N | | 3,3,10,11 | | | | | | | | | 2,2,3 |
| E104 | 2.89 | Xid | | | | 6,6,7,8,8 | | | | 2,2,2,3,4 | | 2,2,2,2,3 | 2,2,3,5,5 |
| E125 | 3.56 | CBA/N | | | | | 5,5,5,7 | | | 4,5,13, >21 | | 2,2,2,4 | 3,3,4,4,5 6 |
| E172 | 3.71 | CBA/N | | | 6,7,3x >21 | | | | | | | | 3,4,6,6,7 |
| E124 | 3.76 | Xid | | | | | | | | 2,2,2,2,3 | | 2,2,2,2,9 | 2,2,2,2,2 |
| E084 | 4.05 | BALB/c | | | | 4x2, 2x >14 | | | | | | | 9x2 |
| E144 | 4.09 | Xid | 2,3,3,6,>2 1 | | | | | 2,3,3,7, >10 | 2,3,3,3,4 | | | | 2,2,2,3,3 |
| | | | | | | | | | | | | | |
| | | | | | | | | | | | | | |
| All | | | 2,3,3,6,>2 1 | 3,3,10,11 | 6,7,3x >21 | 4x2, 6, 6, 7, 8, 8, 2x >21 | 5,5,5,7 | 2,3,3,7 >21 | 2,3,3,3,4 | | 7x2, 3, 3, 4 | | 8x2, 3, 9 21x2, 7x3, 3x 4, 3x5, 3x6, 7 |
| median | | | 3 | 3, 10 | >21 | 6 | 5 | 3 | 3 | 5, 13 | 2 | 2 | 2 |
| a:d | | | 1:4 | 0:4 | 3:2 | 2:9 | 0:4 | 1:4 | 0:5 | 1:3 | 0:10 | 0:4 | 0:10 |
| P rank | | | | | | | | | | | | | |
| P a:d | | | | | | | | | | | | | |

| Pooled Data for Protection against BG9739. by bc100s and FL PspAs | | | | | | | | | | | | | |
|---|---------------|--------|--------------------------|-----------------|-----------------------------------|--------------------|-------------------|-----------------|----------------|-------------------|--------------------------|-----------------|---------------------------------------|
| Exp. | CFU BG9739 | Mice | Days to Death/ immunogen | | | | | | | | | | |
| | | | R36A FL | BC100 (Rx1) | JD908 (WU2) | JS1020 (BG9739) | bc100 (BG9739) | EF3296 FL | EF5668 FL | bc100 (L81905) | JS5010.3 FL (DBL5) | bc100 (DBL5) | JS3020 (DBL6 A) |
| E140 | 2.76 | CBA/N | | 3,3,10,11 | | | | | | | | | 2,2,3 |
| E104 | 2.89 | Xid | | | | 6,6,7,8,8 | | | | 2,2,2,3,4 | | 2,2,2,2, 3 | 2,2,3,5,5 |
| E125 | 3.56 | CBA/N | | | | | 5,5,5,7 | | | 4,5,13, >21 | 2,2,2,4 | | 3,3,4,4,5 6 |
| E172 | 3.71 | CBA/N | | | 6, 7, 3x >21 | | | | | | | | 3,4,6,6,7 |
| E124 | 3.76 | Xid | | | | | | | | 2,2,2,2,3 | | 2,2,2,2, 9 | 2,2,2,2,2 |
| E084 | 4.05 | BALB/c | | | | 4x2, 2x >14 | | | | | | | 9x2 |
| E144 | 4.09 | Xid | 2,3,3,6, >21 | | | | | 2,3,3,7, >10 | 2,3,3,3,4 | | | | 2,2,2,3,3 |
| FL + bc100 BG9739 | | | R36A/Rx1/D39 | WU2 | BG9739 | | EF3296 | EF5668 | L81905 | DBL5 | | DBL6A | Cont. |
| All | | | 2, 4x 3, 6, 10, 11, >21 | 6, 7, 3x >21 | 4x2, 3x5, 2x6, 2x7, 2x8, 2x>21 | | 2,3,3,7, >21 | 2,3x3,4 | 4,5,13, >21 | 10x2, 3, 3, 4, 4 | | 8x2, 3, 9 | 21x2, 7x3, 3x 4, 3x5, 3x6, 7 |
| median days alive | | | 3 | >21 | 6 | | 3 | 3 | 5, 13 | 2 | | 2 | 2 |
| alive : dead | | | 1:8 | 3:2 | 2:13 | | 1:4 | 0:5 | 1:3 | 0:14 | | 0:10 | 0:38 |
| P - days alive | | | 0.0096 | <0.0001 | 0.0013 | | n.s. | n.s. | 0.0022 | n.s. | | n.s. | |
| P- alive : dead | | | n.s. | 0.0008 | n.s. | | n.s. | n.s. | n.s. | n.s. | | n.s. | |
| Score | | | + | +++ | + | | 0+ | 0 | ++ | 0 | | 0 | |
| % alive | | | 11 | 60 | 13 | | 25 | 0 | 25 | 0 | | 0 | 0 |
| % protected | | | 11 | 60 | 13 | | 25 | 0 | 25 | 0 | | 0 | 0 |
| BG9739 challenge | | | R36A/Rx1/D39 | WU2 | BG9739 | | EF3296 | EF5668 | L81905 | DBL5 | | DBL6A | Cont. |

TABLE 50

110 K

| BC9739 | days of death | median days of death | alive/dead | P value based on days to death | P value based on alive : dead | Score | % Alive | % | |
|--------------|---------------|----------------------|------------|--------------------------------|-------------------------------|-------|---------|----|--|
| All immune | | 3 | 8:59 | 0.009 | 0.023 | ++ | 12 | 12 | |
| All controls | | 2 | 0:38 | | | | | | |

TABLE 51

Pooled Data for Protection against EF3296. by various PspAs

| Exp. | CFU EF3296 | Mice | Days to Death/ immunogen | | | | | |
|----------------------|------------|--------|--------------------------|-------------|---------------------|--------------------|------------------|-------------------------|
| | | | Rx1 BC100 | JD908 WU2 | JS1020 (BG9739) | JS5010.3 FL (DBL5) | JS3020 (DBL6A) | control |
| E84 ¹ | 3.99 | BALB/c | | | 4x2, >14 | | | 9x2 |
| E140 | 2.92 | CBA/N | 3, 4, 6, >21 | | | | | 3,3,3 |
| E104 | 3.11 | CBA/N | | | 4,5,5,5,6 | 2,2,2,3,3 | 2,2,3,4,5 | 2,2,2,3,4 |
| E124 | 3.94 | CBA/N | | | | 1,1,2,2,2 | 1,1,2,2,2 | 1,1,2,2,2 |
| E172 | 4.06 | CBA/N | | | | | | 3, 4x6 |
| All | | | 3, 4, 6, >21 | 3,3,5,5,>21 | 4x2, 4, 3x5, 6, >21 | 1,1,5x2, 3,3 | 1, 1,5x2, 3, 4,5 | 1, 1, 15x2, 3x3, 4, 4x6 |
| median days to death | | | 5 | 5 | 4.5 | 2 | 2 | 2 |
| alive : dead | | | 1:3 | 1:4 | 1:9 | 0:9 | 0:10 | 0:27 |
| P - days to death | | | 0.0077 | 0.0094 | n.s. | n.s. | n.s. | |
| P - alive : dead | | | n.s. | n.s. | n.s. | n.s. | n.s. | n.s. |
| Score | | | ++ | ++ | 0+ | 0 | 0 | |
| % alive | | | 25 | 20 | 10 | 0 | 0 | 0 |
| % prot. | | | 25 | 20 | 10 | 0 | 0 | 0 |
| Best | | | | | | | | |
| EF3296 challenge | | | Rx1 BC100 | JD908 WU2 | JS1020 (BG9739) | JS5010.3 FL (DBL5) | JS3020 (DBL6A) | control |

| EF3296 challenge | median days alive | alive : dead | P - days to death | P - alive : dead | Score | % alive | % prot |
|------------------|-------------------|--------------|-------------------|------------------|-------|---------|--------|
| All immune | 3 | 3:35 | n.s. | n.s. | 0 | 8 | 8 |
| All control | 2 | 0:27 | | | | | |

TABLE 52

Pooled Data for Protection against EF5668. by various FL-PspAs and bc100s

| Exp. | CFU EF5668 | Mice | Days to Death/ immunogen | | | | | | | | | |
|-------------------|------------|-------|--------------------------|------------|-------------|-----------------|----------------|--------|---------------|------------------|--------------|-----------------------------------|
| | | | R36A | Rx1 BC100 | JD908 (WU2) | JS1020 (BG9739) | EF3296 | EF5668 | L81905 | JS5010.3 FL DBL5 | JS3020 DBL6A | control |
| E143 | 3.0 | CBA/N | | | | | | 5x >10 | | | | 1,1,2,2, >10 |
| E140 | 3.59 | CBA/N | | 4,6,12,>21 | | | | | | | | 2,4,6 |
| E171 | 3.69 | CBA/N | | | 2,2,2,3,>21 | | | | 3,3,4, 2x >21 | | | 1,3,6,6,7 |
| E124 | 3.90 | CBA/N | | | | | | | | 3,3,3x >15 | 3,4,5,6,6 | 3,3,3,4,9 |
| E145 | 3.94 | CBA/N | 3,4,4, 16, >19 | | | 2,10, 3x >19 | 2,4,13, 2x >19 | | | | | 2,3,3,4, >21 |
| Pool | | | 3,3x4,6,12,16, 2x >21 | | 2,2,2,3,>21 | 2,10, 3x >21 | 2,4,13, 2x >21 | 5x >21 | 3,3,4, 2x >21 | 3,3,3x >21 | 3,4,5,6,6 | 3x1,4x2, 6x3,3x4, 3x6,7,9, 2x >21 |
| median days alive | | | 6 | | 2 | >21 | 13 | >21 | 4 | >21 | 5 | 3 |
| alive : dead | | | 2:7 | | 1:4 | 3:2 | 2:3 | 5:0 | 2:3 | 3:2 | 0:5 | 2:21 |
| P - days alive | | | 0.013 | | n.s. | 0.0187 | n.s. | 0.001 | n.s. | n.s. | n.s. | |
| P - alive : dead | | | n.s. | | n.s. | 0.027 | n.s. | 0.0002 | n.s. | 0.027 | n.s. | |
| Score | | | + | | 0 | +++ | 0+ | +++ | 0+ | + | 0+ | |
| % alive | | | 22 | | 25 | 60 | 40 | 100 | 40 | 60 | 0 | 9 |
| % prot | | | 14 | | 18 | 56 | 34 | 100 | 34 | 56 | -10 | 9 |
| EF5668 | | | R36A/Rx1/D39 | | WU2 | BG9739 | EF3296 | EF5668 | L81905 | DBL5 | DBL6A | control |

| Summary of protection against EF6796 | | | | | | |
|--------------------------------------|--------------|---------|-------------|------------|---------------|-----------------|
| Immunogen | alive : dead | % alive | % protected | median DOD | P -time alive | P alive vs dead |
| Rx1 | 4:0 | 100 | 100 | >21 | 0.029 | 0.029 |
| controls | 0:3 | 0 | 0 | 1 | -- | -- |

* +++ = statistically significant protection from death with $\geq 50\%$ protected;

TABLE 53

| Pooled Data for Protection against DBL6A. by various FL PspAs and bc100 PspAs | | | | | | | | | | | | | | | |
|--|------------------|-------|--------------------------|---------------------|---------------------------|------------------|--------------------|--------------------|-------------------------|-----------------|---------------------------|------------------|------------------|--|---------------|
| Exp. | CFU DBL6 A | Mice | Days to Death/ immunogen | | | | | | | | | | | | |
| | | | BC100 Rx1 | R36A | JD908 WU2 | JS1020 BG9739 | bc100 BG9739 | EF3296 | EF5668 | L81905 FL | bc100 L81905 | JS5010.3 DBL5 | bc100 DBL5 | JS3020 DBL6A | control |
| E171 | 2.69 | CBA/N | | | 6,7,8,9, >21 | | | | | 3,3,7,9, >21 | | | | | 2,3,4,6,6 |
| E152 | 3.24 | CBA/N | | | | | 15, 3x >21 | | | | 7,16, 2x >21 | | 8, 10, 13, 21 | | 3x 3, 4, 3x 6 |
| E140 | 3.25 | CBA/N | 4x >21 | | | | | | | | | | | | 4, 7, 7 |
| E146 | 3.57 | CBA/N | | 7, 8, 10, 2x >21 | | | | 6, 8, 9, 10, 10 | 10, 13, 3x >21 | | | 7,8,12, 13,13 | | 9, 4x >21 | 4,4,5,5,18 |
| E129 | 4.14 | CBA/N | | | | 3,6,8,10, 13 | | | | | | | | | 4,5,6,8,>23 |
| Total | | | | | | | | | | | | | | | |
| Name of Pools | | | R36A/Rx1/D39 | WU2 | BG9739 | | EF3296 | EF5668 | L81905 | | DBL5 | | DBL6A | controls | |
| Pooled data | | | 7, 8, 10, 6x >21 | 6,8,9, >21 | 3,6,8,10,13,15, 3x >21 | | 6, 8, 9, 10, 10 | 10, 13, 3x >21 | 3,3,7,7,9,16, 2x >21 | | 7,8,8,10,12, 3x 13, 21 | | 9, 4x >21 | 2, 4x 3, 6x 4, 3x 5, 6x 6, 7, 7, 8, 18, >21 | |
| median days alive | | | >21 | 8.5 | 13 | | 9 | >21 | 8 | | 12 | | >21 | 5 | |
| alive : dead | | | 6:3 | 1:3 | 3:6 | | 0:5 | 3:2 | 2:6 | | 0:9 | | 4:1 | 1:24 | |
| P - days alive | | | <0.0001 | 0.0082 | 0.0025 | | 0.0036 | 0.0001 | 0.037 | | 0.002 | | <0.0001 | | |
| P - alive : dead | | | 0.0019 | n.s. | 0.048 | | n.s. | 0.0093 | n.s. | | n.s. | | 0.0009 | | |
| Score | | | +++ | +± | ++ | | +± | +++ | +± | | +± | | +++ | | |
| | | | 67 | 25 | 33 | | 0 | 60 | 25 | | 0 | | 80 | 4 | |
| | | | 66 | 22 | 30 | | -4 | 58 | 22 | | -4 | | 79 | 0 | |
| DBL6A challenge | | | R36A/Rx1/D39 | WU2 | BG9739 | | EF3296 | EF5668 | L81905 | | DBL5 | | DBL6A | controls | |

| DBL6A challenge | days of death | median days of death | alive : dead | P value based on days to death | P value based on alive : dead | Score | % alive | % prot. |
|-----------------|---------------|----------------------|--------------|--------------------------------|-------------------------------|-------|---------|---------|
| All immune | | 12.5 | 19:35 | <0.0001 | 0.0019 | ++ | 35 | 33 |
| All control | | 5 | 1:24 | | | | | |

TABLE 54

| Pooled Data for Protection against BG9163 by various PspAs | | | | | | |
|--|---------------|-------|--------------------------|------------|--------------------|-----------------------------|
| Exp. | CFU BG9163 | Mice | Days to Death/ immunogen | | | |
| | | | Rx1 | Rx1.BCG | JS1020 (BG9739) | all immune control |
| E169 | 2.67 | CBA/N | 5x >24 | | | 4,5,8,8,>24 |
| E140 | 3.14 | CBA/N | n.v. | | | |
| E129 | 4.0 | CBA/N | | | 12, 4x >23 | 7, 9, 9, 13, >23 |
| E028 | 6.217 | CBA/N | | 6, 3x >21 | | 5,6,8,10 |
| Immunogens | | | Rx1/R36A/D39 | BC9739 | all immune | control |
| Pooled Data | | | 6, 8x >21 | 12, 4x >21 | 6, 12, 12x >21 | 4,5,7,8,8,9, 9,12,2x >21 |
| median days alive | | | >21 | >21 | >21 | 8.5 |
| alive : dead | | | 8:1 | 4:1 | 12:2 | 2:8 |
| P - days alive | | | 0.0086 | 0.0097 | 0.0027 | |
| P - alive : dead | | | 0.0045 | 0.047 | 0.0022 | |
| % alive | | | 89 | 80 | 86 | 20 |
| % prot. | | | 86 | 75 | 83 | 0 |
| score | | | +++ | +++ | +++ | |
| BG9163 Challenge | | | Rx1/R36A/D39 | BC9739 | all immune | control |

| EF5668 | days of death | median days of death | alive : dead | P value based on days to death | P value based on alive : dead | Score | % alive | % prot. |
|-------------|---------------|----------------------|--------------|--------------------------------|-------------------------------|-------|---------|---------|
| All immune | | 8 | 18:26 | 0.0015 | 0.005 | ++ | 41 | 35 |
| All control | | 3 | 2:21 | | | | | |

TABLE 55

Pooled Data for Protection against L81905. by various FL-PspAs

| Exp. | CFU L81905 | Mice | Days to Death/ immunogen | | | | | | | | | | | |
|----------------|---------------|-------|--------------------------|----------------|----------------|--------------------------------|-----------------|------------------|-----------|-----------------|-----------------------|-----------------|------------------------------|------------------------------------|
| | | | R36A | BC100 (Rx1) | JD908 (WU2) | JS1020 (BG9739) | bc100 BG9739 | EF3296 | EF5668 | bc100 L81905 | JS3010.3 (DBL5) | bc100 (DBL5) | JS3020 (DBL6A) | control |
| E172 | 2.45 | CBA/N | | | 3,4,5,6,6 | | | | | | | | | 3,3,4,4,4 |
| E140 | 3.11 | CBA/N | | 2,5,5,6,8 | | | | | | | | | | 2,2,2,3,3 |
| E084 | 3.86 | BALB | | | | 2, 2, 5x >14 | | | | | | | | 1, 8x 2 |
| E104 | -3.5 | CBA/N | | | | 3,7,8,8,11 | | | | | 3,3,3,2x >22 | | 3,4,5,5,6 | 2, 4, 4, 4, 5 |
| E124 | -3.5 | CBA/N | | | | | | | | | 2, 2, 2, 2, 3 | | 2,2,2,3,5 | 1,2,2,2,2 |
| E125 | 3.6 | CBA/N | | | | | 5,6,8,8 | | | 3,4,6,8 | | 4,5,5,5 | | 2,2,3,5,5,5 |
| E144 | 4.11 | CBA/N | 3,3,5,6, >10 | | | | | 6,6,6, 2x >10 | 2,2,3,3,3 | | | | | 2, 2, 3x 3 |
| All | | | 3,3,5,6, >21 | | 3,4,5,6,6 | 2,2,3,7,8,8 , 11, 3x >21 | 5,6,8,8 | 6,6,6, 2x >10 | 2,2,3,3,3 | 3,4,6,8 | 4x 2, 4x 3, 2x >21 | 4,5,5,5 | 3x 2, 3, 3, 4, 3x 5, 6 | 1, 1, 20x 2 8x 3, 6x 4, 4x 5 |
| median | | | 5 | 5 | 5 | >21 | 7 | 6 | 3 | 5 | 3 | 5 | 3.5 | 2 |
| alive: dead | | | 1:4 | 0:5 | 0:5 | 5:7 | 0:4 | 2:3 | 0:5 | 0:4 | 2:8 | 0:4 | 0:10 | 0:40 |
| P rank | | | | | | | | | | | | | | |
| P a:d | | | | | | | | | | | | | | |
| score | | | | | | | | | | | | | | |

TABLE 56

Protection against L81905. by various bc100s & FL-PspAs pooled together

| Exp. | CFU L81905 | Mice | Days to Death/ immunogen | | | | | | | | | | | |
|-----------------------|---------------|-------|--------------------------|----------------|----------------|-------------------------------|-----------------|------------------|-----------|-----------------|-------------------------------|-----------------|---------------------------|-----------------------------------|
| | | | R36A | BC100 (Rx1) | JD908 (WU2) | JS1020 (BG9739) | bc100 BG9739 | EF3296 | EF5668 | bc100 L81905 | JS5010.3 (DBL5) | bc100 (DBL5) | JS3020 (DBL6A) | control |
| E172 | 2.45 | CBA/N | | | 3,4,5,6,6 | | | | | | | | | 3,3,4,4,4 |
| E140 | 3.11 | CBA/N | | 2,5,5,6,8 | | | | | | | | | | 2,2,2,3,3 |
| E084 | 3.86 | BALB | | | | 2, 2, 5x >14 | | | | | | | | 1, 8x2 |
| E104 | -3.5 | CBA/N | | | | 3,7,8,8,11 | | | | | 3,3,3,2x >22 | | 3,4,5,5,6 | 2, 4, 4, 4, 5 |
| E124 | -3.5 | CBA/N | | | | | | | | | 2, 2, 2, 2, 3 | | 2,2,2,3,5 | 1,2,2,2,2 |
| E125 | 3.6 | CBA/N | | | | | 5,6,8,8 | | | 3,4,6,8 | | 4,5,5,5 | | 2,2,3,5,5,5 |
| E144 | 4.11 | CBA/N | 3,3,5,6, >10 | | | | | 6,6,6, 2x >10 | 2,2,3,3,3 | | | | | 2, 2, 3x3 |
| Pooled | | | 2,3,3, 3x 5, 6,6,8,>21 | | 3,4,5,6,6 | 2,2,3,5,6,7,8x 8,11, 5x>21 | | 6,6,6, 2x >10 | 2,2,3,3,3 | 3,4,6,8 | 4x 2, 4x 3, 4,5,5,5, 2x>21 | | 3x 2, 3, 3, 4, 3x 5, 6 | 1, 1, 20x2 8x 3, 6x 4, 4x 5 |
| median days alive | | | 5 | | 5 | 8 | | 6 | 3 | 5 | 3 | | 3.5 | 2 |
| alive: dead | | | 1:9 | | 0:5 | 5:11 | | 2:3 | 0:5 | 0:4 | 2:12 | | 0:10 | 0:40 |
| P - days alive | | | 0.0005 | | 0.0035 | <0.0001 | | 0.0002 | n.s. | 0.01 | 0.035 | | 0.044 | |
| P - alive : dead - | | | n.s. | | n.s. | 0.0001 | | 0.01 | n.s. | n.s. | n.s. | | n.s. | |
| score | | | + | | + | ++ | | ++ | 0 | + | + | | + | |
| % alive | | | 10 | | 0 | 31 | | 40 | 0 | 0 | 14 | | 0 | 0 |
| % protected | | | | | | | | | | | | | | |
| challenge with L81905 | | | R36A/Rx1/D39 | | WU2 | BG9739 | | EF3296 | EF5668 | L81905 | DBL5 | | DBL6A | controle |

| L21905 challenge | days of death | median days of death | alive : dead | P value based on days to death | P value based on alive : dead | Score | % alive | % prot. |
|------------------|---------------|----------------------|--------------|--------------------------------|-------------------------------|-------|---------|---------|
| All immune | | 5 | 10:59 | <0.0001 | 0.008 | ++ | 14 | 14 |
| All control | | 2 | 0:40 | | | | | |

TABLE 57

| Pooled Data for Protection against DBL5 by various FL-PspAs & bc100s | | | | | | | | | | | |
|--|------|--------|--------------------------|-----------|-------------------------|--------------|-----------|--------------|------------------|-------------------------|--------------|
| Exp | CFU | Mice | Days to Death/ immunogen | | | | | | | | |
| | | | R36A | BC100 Rx1 | JS1020 BC9739 | bc100 JS1020 | EF5668 | bc100 L81905 | JS5010.3 DBL5 | bc100 DBL5 | JS3020 DBL6A |
| E84 ¹ | 3.90 | BALB/c | | | 6x2 | | | | | | 9x2 |
| E140 | 3.27 | CBA/N | | 4,4,5,5,5 | | | | | | | 2,2,2 |
| E104 | 3.39 | Xid | | | 3,3,6, >22, >22 | | | | 7,7,15, >22, >22 | | 2,2,4,5,5 |
| E124 | 3.76 | Xid | | | | | | | 2,2,2,5, >15 | | 5x2 |
| E125 | 3.81 | CBA/N | | | | 3,3,4,5 | | 3,3,4,4 | | 2,2,2,>21 | 5x2,5 |
| E144 | 4.13 | XID | 3,3,3,3, >10 | | | | 2,2,3,4,4 | | | | 5x2 |
| total | | | | | | | | | | | |
| name of pool | | | R36A/Rx1/D39 | | BC9739 | | EF5668 | L81905 | | DBL5 | DBL6A |
| pooled data | | | 4x3, 2x4, 3x5, >21 | | 6x2, 4x3, 4,5, >21, >21 | | 2,2,3,4,4 | 3,3,4,4 | | 6x2, 5, 7, 7, 15, 4x>21 | 7x2, 4, 5, 5 |
| median days alive | | | 4 | | 3 | | 3 | 3.5 | | 6 | 2 |
| alive : dead | | | 1:9 | | 2:12 | | 0:4 | 0:4 | | 4:10 | 0:10 |
| P - days alive | | | <0.0001 | | 0.0063 | | .041 | 0.001 | | 0.0025 | n.s. |
| P - alive : dead | | | n.s. | | n.s. | | n.s. | n.s. | | 0.0056 | n.s. |
| Score | | | + | | + | | ++ | + | | ++ | 0 |
| % alive | | | 10 | | 14 | | 0 | 0 | | 29 | 0 |
| % protected | | | 10 | | 14 | | 0 | 0 | | 29 | 0 |
| DBL5 challenge | | | R36A/Rx1/D39 | | BC9739 | | EF5668 | L81905 | | DBL5 | DBL6A |

¹ This immunization was with cell eluted PspA. Note BALB/c mice were used. Also note 10⁴ Challenge CFU.

| DBL5 challenge | days of death | median days of death | alive : dead | P value based on days to death | P value based on alive : dead | Score | % alive | % prot. |
|----------------|---------------|----------------------|--------------|--------------------------------|-------------------------------|-------|---------|---------|
| All immune | | 3.5 | 7:49 | <0.0001 | 0.034 | ++ | 3.6 | 3.6 |
| All control | | 2 | 0:33 | | | | | |

TABLE 58

| Pooled Data for Protection against EF6796 by various PspAs | | | | | | | | |
|--|------|-------|--------------------------|-----------------|--------|----------------|--------------------|---------|
| Exp. | CFU | Mice | Days to Death/ immunogen | | | | | |
| | | | Rx1 BC100 | JS1020 (BC9739) | L81905 | JS3020 (DBL6A) | JS5010.3 FL (DBL5) | control |
| E140 | 3.75 | CBA/N | 4x>21 | | | | | 1,1,1 |
| E28 | ? | BALB | n.v. | | | | | |

TABLE 59

| Pool of Pools for protection against EF6796 | | | | | |
|---|-------------------|--|---------------|--------------------------|---------------|
| line | Group Description | Delay in time to death and/or survival | | Protection against death | |
| | | days to death (median DOD) | P values etc. | alive : dead | P values etc. |
| 1a | Rx1 | 4x>21 (>21) | 0.029 | 4:0 | 0.029 |
| 1b | Rx1 controls | 1,1,1 (1) | -- | 0:3 | -- |

TABLE 60

| Pooled Data for Protection against BG7322. by various FL- PspAs and bc100s | | | | | | | | | | | | | |
|--|-------------|-------|--------------------------|-----------|----------------|-----------------|-------------|-----------|-----------------|--------------------------------------|--------------------|-------------------|--|
| Exp. | CFU BG 7322 | Mice | Days to Death/ immunogen | | | | | | | | | | |
| | | | D39/ R36A | Rx1 BC100 | JD908 (WU2) | bc100 BG9739 | EF3296 | EF5668 | bv100 L81905 | JSS0103 DBL5 | bc100 DBL5 | JSS020 DBL6A | control |
| E171 | 2.78 | CBA/N | | | 10, 15, 3x >21 | | | | | | | | 1, 3, 6, 6, 7 |
| S143 | 3.0 | CBA/N | | | | | | 7, 8x >10 | | | | | 2, 2, 4, 5, 7, 7, 8, 8 |
| E140 BC100 | 3.14 | CBA/N | | 4x >21 | | | | | | | | | 0, 6, 6, >21 |
| E152 | 3.11 | CBA/N | | | | 12, 13, 16, >21 | | | 10, 12, 13, >21 | | >21, >21, >21, >21 | | 5, 7, 7, 8, 8, 9, 14 |
| E146 | 3.57 | CBA/N | 18, 20, 3x >21 | | | | 5, 3x 6, 10 | | | 6, 10, 11, 11, 19 | | 4, 8, 11, 18, >21 | 4, 5, 5, 6, >21 |
| E169 | 3.94 | CBA/N | 5x >21 | | | | | | | | | | 2, 5, 5, 6, 7 |
| Immunogens | | | R36A/Rx1/D39 | | JD908 | BG9739 | EF3296 | EF5668 | L81905 | DBL5 | | DBL6A | Cont. |
| Pools | | | 18, 20, 12x >21 | | 10, 15, 3x >21 | 12, 13, 16, >21 | 5, 3x 6, 10 | 7, 8x >21 | 10, 12, 13, >21 | 6, 10, 11, 11, 19 >21, >21, >21, >21 | | 4, 8, 11, 18, >21 | 1, 3x 2, 3, 3, 4, 4, 5x 5, 7x 6, 6x 7, 4x 8, 9, 14, 2x >21 |
| median day alive | | | >21 | | >21 | 14.5 | 6 | >21 | 12.5 | >21 | | 11 | 6 |
| alive : dead | | | 9:0 | | 3:2 | 1:3 | 0:5 | 8:1 | 1:3 | 4:5 | | 1:3 | 2:32 |
| P - days alive | | | <0.0001 | | 0.0007 | 0.001 | n.s. | <0.0001 | 0.0013 | 0.0002 | | 0.028 | |
| P - alive : dead | | | <0.0001 | | 0.004 | n.s. | n.s. | <0.0001 | n.s. | 0.0076 | | n.s. | |
| % alive | | | 100 | | 60 | 25 | 0 | 89 | 25 | 80 | | 25 | 6 |
| % protected | | | 100 | | 57 | 22 | 0 | 88 | 22 | 79 | | 22 | 6 |
| Score | | | +++ | | +++ | ++ | 0 | +++ | ++ | +++ | | ++ | |
| BG7322 Challenge | | | R36A/Rx1/D39 | | JD908 | BG9739 | EF3296 | EF5668 | L81905 | DBL5 | | DBL6A | Cont. |

| BG7322 Challenge | median days of death | alive : dead | P value based on days to death | P value based on alive : dead | Score | % alive | % prot. |
|------------------|----------------------|--------------|--------------------------------|-------------------------------|-------|---------|---------|
| All immune | >21 | 30:25 | <0.0001 | <0.0001 | +++ | 55 | 52 |
| All controls | 6 | 2:32 | | | | | |

EXAMPLE 8 - Ability of PspA immunogens to protect against individual challenge strains

In example 7 some of the capsular type 2, 4, and 5 strains were not completely protected from death by immunization. In these studies the BALB/cByJ mouse was used instead of the hypersusceptible, immunodeficient CBA/N mouse used for the Example 7 studies. With the BALB/cJ mouse it was observed that immunization with PspA was in fact able to protect against death with capsular type 2, 4, and 5 pneumococci. This result is shown in the table below.

The data from Table 60A also demonstrates that a mixture of 4 - 5 full length PspAs was as effective, or more effective than immunization with a single PspA.

Table 60A. Days of death of BALB/cBYJ mice after immunization with monovalent and polyvalent vaccine.

| Challenge Strains | | | | | Immunogen | | | | |
|-------------------|------|------|----------------|----------------|-----------------|---|-----------------------|-----------------------------|--|
| strain | caps | PapA | ppvA | Log | Days to Death | | | | |
| name | type | type | B region clade | Challenge dose | 1 mg R36A + CFA | 4 - 5 valent mixture (0.5µg each) + CFA | JY2141 + CFA | None | |
| D39 | 2 | 25 | 2 | 4.76 | 3, 4x >21 | 3, 4x >21 | 3, 4, 5, 11, >21 | 3, 3, 4, 4, 8 | |
| WU2 | 3 | 1 | 2 | 4.8 | 4x >21 | 4x >21 | 6, 3x >21 | 3, 4, 2x >21 | |
| A66 | 3 | 13 | ? | 4.7 | 3, 3, >21, >21 | 2, 3x >21 | 2, 2, 3, 4 | 2, 3, 4, 4 | |
| BG9739 | 4 | 26 | 1 | 4.07 - 4.4 | 7, 8x >21 | 3, 8x >21 | 1, 5, 6, 6, 9, 4x >21 | 3, 3, 3, 4, 6, 7, 7, 2x >21 | |

| | | | | | | | | |
|--------|----|----|---|----------------|--------------------------------|------------------------|--------------------------------------|--------------------------|
| L81905 | 4 | 23 | 1 | 6.90 - 6.96 | 2, 2, 2, 2, 5, 5, 4x >21 | 2, 6, 8, 9, 6x >21 | 1, 1, 1, 1, 2, 3, 4, 5, 2x >21 | 1, 4x 2, 3x 3, 4, >21 |
| EF5668 | 4 | 12 | 4 | 6.10- 6.93 | 3, 3, 4, 7x >21 | 3x 3, 6x >21 | 4x 3, 4, 4, 6, 6, >21 | 3, 5x 4, 6, >21 |
| DBL5 | 4 | 33 | 2 | 3.30 | 7, 14, 3x >21 | 3, 5, 5, 2x >21 | 2, 2, 2, 4, 6 | 4, 5, 5, 6, 9 |
| DBL6A | 6A | 19 | 1 | 4.34 | 6, 9, 10, 11, >21 | 10, 11, 12, 13, >21 | 3, 11, 11, 13, 16 | 8, 9, 11, 21, >21 |
| BG7322 | 6B | 21 | ? | 3.9 | 8, 8, 3x >21 | 5x >21 | 6, 6, 7, 8, 10 | 2, 5, 6, 8, 8 |

Note, JY2141 is a preparation from a strain that lacks PspA. None = no immunization.

Note, mice were given two immunizations with PspA two weeks apart and challenged intravenously 2 weeks after the last immunization. The first immunization was given with complete Freund's adjuvant (CFA) subcutaneously, the second immunization was given intraperitoneally in saline.

¹ 4 valent vaccine mixture R36A, BG9739, EF5668, and DBL5 -- all E180

² 4 valent vaccine mixture R36A, BG9739, DBL5, EF3296 D39 and DBL6A

³ 5 valent vaccine mixture R36A, BG9739, DBL5, EF3296, EF5668

EXAMPLE 9 - CHARACTERIZATION OF PspA EPITOPES WITHIN PNEUMOCOCCAL STRAINS MC25-28

The strains examined came from a group of 13 capsular serotype 6B strains which have been identified that are members of a multiresistant clone, having resistance to penicillin, chloramphenicol, tetracycline, and some have acquired resistance to erythromycin. The pneumococcal isolates described in the following studies (MC25-28) are members of this 6B clone. Although previously thought to be geographically restricted to Spain (unlike the widespread multiresistant Spanish serotype 23F clone), members of this clone have been shown to be responsible for an increase in resistance to penicillin in Iceland (Soares, S., et al., J. Infect. Dis. 1993; 168: 158-163).

The following techniques were used to characterize the location of difference PspA epitopes:

Bacterial cell culture. Bacteria were grown in Todd-Hewitt broth with 0.5% yeast extract or on blood agar plates overnight at 37°C in a candle jar. Capsular serotype was confirmed by cell agglutination using Danish antisera (Statens Seruminstitut, Copenhagen, Denmark). The isolates were subtyped as 6B by Quellung reaction, utilizing rabbit antisera against 6A or 6B capsule antigen.

Bacterial lysates. Cell lysates were prepared by incubating the bacterial cell pellet with 0.1% sodium deoxycholate, 0.01% sodium dodecylsulfate (SDS), and 0.15 M sodium citrate, and then diluting

the lysate in 0.5M Tris hydrochloride (pH 6.8). Total pneumococcal protein in the lysates was quantitated by the bicinchoninic acid method (BCA Protein Assay Reagent; Pierce Chemical Company, Rockford, IL).

PspA serotyping. Pneumococcal cell lysates were subjected to SDS-PAGE, transferred to nitrocellulose membranes, and developed as Western blots using a panel of seven MAbs to PspA. PspA serotypes were assigned based on the particular combination of MAbs with which each PspA was reactive.

Colony immunoblotting. A ten mL tube of Todd-Hewitt broth with 0.5% yeast extract was inoculated with overnight growth of MC25 from a blood agar plate. The isolate was allowed to grow to a concentration of 10^7 cells/mL as determined by an O.D. of 0.07 at 590nm. MC25 was serially diluted and spread-plated on blood agar plates to give approximately 100 cells per plate. The plates were allowed to grow overnight in a candle jar, and a single blood agar plate with well-defined colonies was selected. Four nitrocellulose membranes were consecutively placed on the plate. Each membrane was lightly weighted and left in place for 5 min. In order to investigate the possibility of phase-variation between the two proteins detected on Western blots a single colony was picked from the plate, resuspended in ringer's solution, and spreadplated onto a blood agar plate. The membranes were developed as Western blots according to PspA serotyping methods.

When the strains MC25-28 were examined with the panel of seven MAbs specific for different PspA epitopes, all four demonstrated the same patterns of reactivity (Fig. 14). The MAbs XiR278 and 2A4 detected a PspA molecule with an apparent molecular weight of 190 kDa in each isolate. In accordance with the PspA serotyping system, the 190 kDa molecule was designated as PspA type 6 because of its reactivity with XiR278 and 2A4, but none of the five other MAbs in the typing system. Each isolate also produced a second PspA molecule with an apparent molecular weight of 82 kDa. The 82 kDa PspA of each isolate was detected only with the MAb 7D2 and was designated as type 34. No reactivity was detected with MAbs Xi126, Xi64, 1A4, or SR4Wr. Results from the colony immunoblotting showed that both PspAs were present simultaneously in these isolates under *in vitro* growth conditions. All colonies on the plate, as well as all of the progeny form a single colony, reacted with MAbs XiR278, 2A4, and 7D2.

EXAMPLE 10 - SOUTHERN BLOT ANALYSIS OF CHROMOSOMAL DNA ISOLATED FROM PNEUMOCOCCAL STRAINS MC25-28

Pneumococcal chromosomal DNA was prepared by the Youderian method (Sheffield, J.S., et al., Biotechniques, 1992; 12: 836-839). Briefly, for a 500 ml culture in THY or THY with 1% choline, cells were centrifuged at 8000 rpm in GSA rotor for 30 minutes at 4°C. The supernatant was decanted, and the cells were washed with 1 to 2 volumes of sterile water to remove

choline, if used. This step was only necessary when sodium deoxycholate was used. The washed cells were centrifuged twice at 8000 rpm in GSA rotor for 10 minutes. Cells were resuspended in 3.5 ml TE buffer, containing 1% SDS or 1% sodium deoxycholate, and incubated at 37°C for 15 minutes if sodium deoxycholate was used. If SDS was used, incubation at 37°C was not necessary. The cells were incubated at 65°C for 15 minutes, and 1/5 volume of 5.0 M potassium acetate was added, and the cell suspension was incubated for 30 minutes at 65°C.

The cells were placed on ice for 60 minutes, and centrifuged at 12,000 rpm in an SS-34 rotor for 10 minutes. The supernatant was transferred to a clean centrifuge tube, and 2 volumes of cold 95% ethanol was added. After mixing, DNA was spooled on to a glass pasteur pipet, and air dried. The DNA was resuspended in 4 ml TE, and 4.0 g cesium chloride was added. The solution was split into two aliquots in ultracentrifuge tubes, and the tubes were filled to their maximum capacity using 1.0 g/ml cesium chloride in TE. Before closing the tubes, 300 µl of 10 µg/ml ethidium bromide was added.

The solution was centrifuged at 45,000 rpm overnight, or for 6 hours at 55,000 rpm. The chromosomal band was extracted using a gradient, at least 6 times with 1 volume each salt-saturated isopropanol. The aqueous phase was extracted by adding 2 volumes 95% ethanol. The DNA came out of solution immediately, and it was spooled on to a pasteur pipet. The DNA pellet was

washed by dipping the spooled DNA in 5 ml 70% ethanol. The DNA was air dried, and resuspended in the desired volume of TE, e.g., 500 μ l.

The cells were harvested, washed, lysed, and digested with 0.5% (st/vol) SDS and 100 μ g/mL proteinase K at 37°C for 1 h. The cell wall debris, proteins, and polysaccharides were complexed with 1% hexadecyl trimethyl ammonium bromide (CTAB) and 0.7M sodium chloride at 65°C for 20 min., and then extracted with chloroform/isoamyl alcohol. DNA was precipitated with 0.6 volumes isopropanol, washed, and resuspended in 10mM Tris-HCl, 1mM EDTA, pH 8.0. DNA concentration was determined by spectrophotometric analysis at 260 nm (Meade, H.M. et al., J. Bacteriol 1982; 149: 114-122; Silhavy, T.J. et al., Experiments with Gene Fusion, Cold Spring Harbor: Cold Spring Harbor Laboratory, 1984; and Murray, M.G., et al., Nucleic Acids Res. 1980; 8 4321-4325).

Probe preparation. 5' and 3' oligonucleotide primers homologous with nucleotides to 26 and 1967 to 1990 of Rx1 *pspA* (LSM13 and LSM2, respectively) were used to amplify the full length *pspA* and construct probe LSM*pspA*13/2 from Rx1 genomic DNA. 5' and 3' oligonucleotide primers homologous to nucleotides 161 to 187 and nucleotides 1093 to 1117 (LSM12 and LSM6, respectively) were used to amplify the variable α -helical region to construct probe LSM*pspA*12/6. PCR generated DNA was purified by Gene Clean (Bio101 Inc., Vista, CA) and random prime-labeled with

digoxigenin-11-dUTP using the Genius 1 Nonradioactive DNA Labeling and Detection Kit as described by the manufacturer (Boehringer Mannheim, Indianapolis, IN).

DNA electrophoresis. For Southern blot analysis, approximately 10 μ g of chromosomal DNA was digested to completion with a single restriction endonuclease (Hind III, Kpn I, EcoRI, Dra I, or Pst I), then electrophoresed on a 0.7% agarose gel for 16-48 h at 35 volts. For PCR analysis, 5 μ L of product were incubated with a single restriction endonuclease (Bcl I, BamH I, Bst I, Pst I, Sac I, EcoR I, Sma I, and Kpn I), then electrophoresed on a 1.3% agarose gel for 2-3 h at 90 volts. In both cases, 1 kb DNA ladder was used for molecular weight markers (BRL, Gaithersburg, MD), and gels were stained with ethidium bromide for 10 min and photographed with a ruler.

Southern blot hybridization. The DNA in the gel was depurinated in 0.25N HCl for 10 min, denatured in 0.5M NaOH and 1.5M NaCl for 30 min, and neutralized in 0.5M Tris-HCl (pH 7.2), 1.5M NaCl and 1mM disodium EDTA for 30 min. DNA was transferred to a nylon membrane (Micron Separations INC, MA) using a POSIBLOT pressure blotter (Stratagene, LaJolla, CA) for 45 min and fixed by UV irradiation. The membranes were prehybridized for 3 h at 42°C in 50% formamide, 5X SSC, 5X Denhardt solution, 25mM sodium phosphate (pH 6.5), 0.5% SDS, 3% (wt/vol) dextran sulfate and 500 μ g/mL of denatured salmon sperm DNA. The membranes were then hybridized at 42°C for 18 h in a solution containing 45%

formamide, 5X SSC, 1X Denhardt solution, 20mM sodium phosphate (pH 6.5), 0.5% SDS, 3% dextran sulfate, 250 μ g/mL denatured sheared salmon sperm DNA, and about 20ng of heat-denatured digoxigenin-labeled probe DNA. After hybridization, the membranes were washed twice in 0.1% SDS and 2X SSC for 3 min at room temperature. The membranes were washed twice to a final stringency of 0.1% SDS in 0.3X SSC at 65°C for 15 min. This procedure yielded a stringency greater than 95 percent. The membranes were developed using the Genius 1 Nonradioactive DNA Labeling and Detection Kit as described by the manufacturer (Boehringer Mannheim, Indianapolis, IN). To perform additional hybridization with other probes, the membranes were stripped in 0.2N NaOH/0.1% SDS at 40°C for 30 min and then washed twice in 2X SSC.

PCR. 5' and 3' primers homologous with the DNA encoding the N- and C-terminal ends of PspA (LSM13 and LSM2, respectively) were used. Reactions were conducted in 50 μ L volumes containing 0.2mM of each dNTP, and 1 μ L of each primer at a working concentration of 50mM. MgCl₂ was used at an optimal concentration of 1.75mM with 0.25 units of Taq DNA polymerase. Ten to thirty ng of genomic DNA was added to each reaction tube. The amplification reactions were performed in a thermal cycler (M.J. Research, Inc.) using the following three step program: Step 1 consisted of a denaturing temperature of 94°C for 2 min; Step 2 consisted of 9 complete cycles of a denaturing temperature of 94°C for 1

min, an annealing temperature of 50°C for 2 min, and an extension temperature of 72°C for 3 min; Step 3 cycled for 19 times with a denaturing temperature 94°C for 1 min, an annealing temperature of 60°C for 2 min, and an extension temperature of 72°C for 3 min; and at the end of the last cycle, the samples were held at 72°C for 5 min to ensure complete extension.

Band size estimation. Fragment sizes in the molecular weight standard and in the Southern blot hybridization patterns were calculated from migration distances. The standard molecular sizes were fitted to a logarithmic regression model using Cricket Graph (Cricket Software, Malvern, PA). The molecular weights of the detected bands were estimated by entering the logarithmic line equation obtained by Cricket Graph into Microsoft Excel (Microsoft Corporation, Redmond, WA) in order to calculate molecular weights based on migration distances observed in the Southern blot.

Since most strains contain a *pspA* gene and a *pspC* gene, it was expected that if an extra gene were present one might observe at least three *pspA* homologous loci in isolates MC25-28. In Hind III digests of MC25-28 each strain revealed 7.7 and 3.6 kb bands when probed with LSM*pspA*13/2 (Figure 15A and 15C). In comparison, when Rx1 DNA was digested with Hind III and hybridized with LSM*pspA*13/2, homologous sequences were detected on 9.1 and 4.2 kb fragments, as expected from previous studies

with PspA (Figure 15A). Results consistent with two *pspA*-homologous genes in MC25-28 were obtained with two *pspA*-homologous genes in MC25-28 digested using four additional enzymes (Table 61).

Table 61. Chromosomal RFLPs with probe LSM*pspA*13/2 for isolates MC25-28 and Rx1

| Restriction Enzyme | Strains Examined | | | | | Restriction Fragments (sizes in kilobases) | |
|--------------------|------------------|------|------|------|-----|--|-----------|
| | MC25 | MC26 | MC27 | MC28 | Rx1 | MC25-28 | Rx1 |
| Hind III | + | + | + | + | + | 7.7, 3.6 | 9.1, 4.2 |
| Kpn I | + | + | + | + | + | 11.6, 10.6 | 10.6, 9.8 |
| EcoR I | + | | | | + | 8.4, 7.6 | 7.8, 6.6 |
| Dra I | + | | | | + | 2.1, 1.1 | 1.9, 0.9 |
| Pst I | + | | | | + | >14, 6.1 | 10.0, 4.0 |

The four isolates examined are all members of a single clone of capsular type 6B pneumococci isolated from Spain. These four isolates are the first in which two PspAs have been observed, i.e., PspA and PspC, based on the observation that bands of different molecular weights were detected by different MAbs to PspA. Mutation and immunochemistry studies have demonstrated that all of the different sized PspA bands from Rx1 are made of a single gene capable of encoding a 69kDa protein, supporting the assertion that two PspAs have been observed, i.e., PspA and PspC.

It has been observed that probes for the 5' half of *pspA* (encoding the α -helical half of the protein) bind the *pspC* sequence of most strains only at a stringency of around 90%.

With chromosomal digests of MC25-28, it was observed that the 5' Rx1 probe LSM ψ spA12/6 (Figure 15D) bound two *pspA* homologous bands at even higher stringency. The same probe bound only the *pspA* containing fragment of Rx1 at the higher stringency (Figure 15B).

Further characterization of the *pspA* gene was done by RFLP analysis of PCR amplified *pspA* from each strain. Since previous studies indicated that individual strains yielded only one product, and since the amplification was conducted with primers based on a known *pspA* sequence, it was assumed that the product amplified from each strain represented the *pspA* rather than the *pspC* gene. When MC25-28 were subjected to this procedure, an amplified *pspA* product of 2.1 kb was obtained from each of the four strains. When digested with Hha I, this fragment yielded bands of 1.1, 0.46, 0.21 and 0.19 kb for each of the four isolates. A single isolate, MC25, was analyzed with eight additional enzymes. Using each restriction enzyme, the sum of the fragments was always approximately equal to the size of whole *pspA* (Figure 16). These results suggested that the 2.1 kb amplified DNA represents the amplified product of only a single *pspA* gene. Rx1 produced an amplified product of 2.0 kb and five fragments of 0.76, 0.468, 0.390, 0.349 and 0.120 kb when digested with Hha I as expected from its known *pspA* sequence.

There are several possible explanations for the observation of PspA and PspC in these strains but not in other

strains. All isolates might make PspA and PspC in culture, but MAbS generally recognize only PspA (perhaps, in this isolate there has been a recombination between *pspC* DNA and the *pspC* locus, allowing that locus to make a product detected by MAb to PspA). All isolates can have PspA and PspC, but the expression of one of them generally does not occur under *in vitro* growth conditions. The *pspC* locus is normally a nonfunctional pseudogene sequence that, for an unexplained reason, has become functional in these isolates. Results from the colony immunoblotting of these isolates failed to show a detectable *in vitro* phase shift between either PspA type 6 (XIR278 and 2A4) or PspA type 34 (7D2) protein. This strengthens the second explanation, and suggests that the second PspA in these isolates is due to the *pspC* gene not being turned off during *in vitro* growth conditions.

Presumably, in these four strains, the second PspA protein is provided by the *pspC* DNA sequence. At high stringency, the probe comprising the coding region of the α -helical half of PspA recognized both *pspA* homologous sequences of MC25-18, but not the *pspC* sequence of Rx1. The finding indicated that the *pspC* sequence of MC25-28 is more similar to the Rx1 *pspA* sequence than the Rx1 *pspC* sequence. If the *pspC* sequence of these strains is more similar to *pspA* than most *pspC* sequences, it could explain why the products of *pspC* genes cannot generally be identified by MAbS.

Example 11 - Identification of conserved and variable regions
of pspA and pspC sequences of S. pneumoniae

The *S. pneumoniae* strains used in this study are listed in Table 62. The strains are human clinical isolates representing 12 capsular and 9 PspA serotypes. All strains were grown at 37°C in 100ml of Todd-Hewitt broth supplemented with 0.5% yeast extract to an approximate density of 5×10^8 cells/ml. After harvesting of the cells by centrifugation (2900 g, 10min), the DNA was isolated, and stored at 4°C in TE (10 mM Tris, 1mM EDTA, pH8.0).

Table 62. *Streptococcus pneumoniae* strains used.

| Strain | Relevant phenotype | Reference |
|---------|---|---------------------------|
| WU2 | Capsular type 3, PspA type 1 | Briles et al., 1981 |
| D39 | Capsular type 2, PspA type 25 | Avery et al., 1944 |
| R36A | Nonencapsulated mutant of D39, pspA type 25 | Avery et al., 1944 |
| Rx1 | Derivative of R36A, PspA type 25 | Shoemaker and Guild, 1974 |
| DBL5 | Capsular type 5, PspA type 33 | Yother et al., 1986 |
| DBL6A | Capsular type 6A, PspA type 19 | Yother et al., 1986 |
| A66 | Capsular type 3, PspA type 13 | Avery et al., 1944 |
| AC94 | Capsular type 9L, PspA type 0 | Waltman et al., 1992 |
| AC17 | Capsular type 9L, PspA type 0 | Waltman et al., 1992 |
| AC40 | Capsular type 9L, PspA type 0 | Waltman et al., 1992 |
| AC107 | Capsular type 9V, PspA type 0 | Waltman et al., 1992 |
| AC100 | Capsular type 9V, PspA type 0 | Waltman et al., 1992 |
| AC140 | Capsular type 9N, PspA type 18 | Waltman et al., 1992 |
| D109-1B | Capsular type 23, PspA type 12 | McDaniel et al., 1992 |
| BG9709 | Capsular type 9, PspA type 0 | McDaniel et al., 1992 |
| L81905 | Capsular type 4, PspA type 25 | McDaniel et al., 1992 |
| L82233 | Capsular type 14, PspA type 0 | McDaniel et al., 1992 |
| L82006 | Capsular type 1, PspA type 0 | McDaniel et al., 1992 |

Approximately 5µg of chromosomal DNA was digested with *Hind*III according to the manufacturer's instructions (Promega, Inc., Madison, WI). The digested DNA was subjected to electrophoresis at 35 mV overnight in 0.8% agarose gels and then vacuum-blotted onto Nytran® membranes (Schleicher & Schuell, Keene, NH).

The oligonucleotides used were based on the previously determined sequence of Rx1 *pspA*. Their position and orientation relative to the structural domains of Rx1 *pspA* are shown in Figure 17. Labeling of oligonucleotides and detection of probe-target hybrids were both performed with the Genius System® according to manufacturer's instructions (Boehringer-Mannheim, Indianapolis, IN). All hybridizations were done for 18 hours at 42°C without formamide. By assuming that 1% base-pair mismatching results in a 1°C decrease in T_m , arbitrary designations of "high" and "low" stringency were defined by salt concentration and temperature of post-hybridization washes. Homology between probe and target sequences was derived using calculated T_m by established methods. High stringency is defined as $\geq 90\%$, and low stringency is $\leq 85\%$ base-pair matching.

PCR primers, which were also used as oligonucleotide probes in Southern blotting and hybridizations, were designed based on the sequence of *pspA* from pneumococcal strain Rx1. These oligonucleotides were synthesized by Oligos, Etc. (Wilson, OR), and are listed in Table 63.

Table 63. Oligonucleotide sequences.

| Primer | 5' -> 3' |
|--------|----------------------------------|
| LSM111 | CCGGATCCAGCTCCTGCACCAAAC |
| LSM2 | GCGGTCGACGCTTAAACCCATTACCATTTGG |
| LSM3 | CCGGATCCTGAGCCAGAGCAGTTGGCTG |
| LSM4 | CCGGATCCGCTCAAAGAGATTGATGAGTCTG |
| LSM5 | GCGGATCCCGTAGCCAGTCAGTCTAAAGCTG |
| LSM6 | CTGAGTCGACTGGAGTTTCTGGAGCTGGAGC |
| LSM7 | CCGGATCCAGCTCCAGCTCCAGAACTCCAG |
| LSM9 | GTTTTTGGTGCAGGAGCTGG |
| LSM10 | GCTATGGCTACAGGTTG |
| LSM12 | CCGGATCCAGCGTGCCTATCTTAGGGGCTGGT |
| LSM112 | GCGGATCCTTGACCAATARRRACGGAGGAGGC |

PCR was done with an MJ Research, Inc., Programmable Thermal Cycler (Watertown, MA), using approximately 10 ng of genomic pneumococcal DNA as template with designated 5' and 3' primer pairs. The sample was brought to a total volume of 50 μ l containing a final concentration of 50 mM KCl, 10 mM Tris-HCl (pH 8.3), 1.5mM MgCl₂, 0.01% gelatin, 0.5 μ M of each primer, 200 μ M of each deoxynucleoside triphosphate, and 2.5 U of Taq DNA polymerase. The samples were denatured at 94°C for 2 minutes and subjected to 10 cycles consisting of: 1 min at 94°C, 2 min at 50°C, and 3 min at 72°C, followed by 20 cycles of: 1 min at 94°C, 2 min at 60°C, and 3 min at 72°C. After 30 total cycles, the samples were held at 72°C for an additional 5 min prior to

cooling to 4°C. The amplicons were then analyzed by agarose gel electrophoresis.

Oligonucleosides were used to probe *Hind*III digests of DNA from 18 strains of *S. pneumoniae* under conditions of low and high stringency. Each strain was also screened using a full-length *pspA* probe. Table 64 summarizes the results for each strain under conditions of high stringency. Strain Rx1 is a laboratory derivative of the clinical isolate D39 and consequently, both strains showed identical hybridization patterns and are a single column in Table 64.

Table 64. Summary of hybridization of oligonucleotides with *HINDIII* chromosomal restriction fragments.

| Probe | Rsl/ 139 | WU 2 | DBLS | DMSA A | AC6 | ACM | AC17 | AC40 | AC107 | AC100 | AC14 0 | DIII0 9 | BG9709 | BG58C | L8190 5 | L8223 3 | L82006 |
|-------------------------|-------------|---------|---------|-----------|---------|---------|---------|---------|---------|--------|-----------|------------|---------|----------------|----------------|------------|---------|
| | | | | | | | | | | | | | | | | | |
| PL- Rsl ^a | 40.9.1 6 | 3.8 | 3.7.5.8 | 30.3.4 | 3.6.4.3 | 3.6.6.3 | 3.6.6.3 | 3.2.3.6 | 3.2.3.6 | 40.8.0 | 30.4.0 | 3.3.4.7 | 3.3.4.7 | 1.4.3.2 3.6 | 3.6.5.2 | 8.2.3.7 | 4.3.6.4 |
| LSM12 | 40.9.1 | 3.8 | 3.7.5.8 | 30.3.4 | 4.3 | . | 3.6.6.3 | 3.2.3.6 | . | 40.8.0 | 4.0 | 3.3.4.7 | 2.2.9.6 | 1.4.3.2 3.6 | 3.6 | 1.3.3.7 | . |
| LSM5 | 4.0 | . | . | . | . | . | 3.6.6.3 | . | . | . | . | . | 2.2.9.6 | 3.6 | 1.2.2.3 3.6 | . | . |
| LSM3 | 4.0 | 3.8 | . | . | . | . | 6.3 | . | . | . | . | . | 2.2 | 3.6 | 3.6 | . | . |
| LSM4 | 4.0 | . | . | . | . | . | . | . | . | . | . | . | 2.2 | 3.6 | 3.6 | 3.7 | . |
| LSM7 | 40.9.1 | 3.8 | 3.7 | 30.3.4 | 3.6 | . | . | 3.2.3.6 | . | . | 30.4.0 | 3.3.4.7 | 2.2.9.6 | 3.6 | 3.6.2.3 | 3.7 | . |
| LSM11 | 40.9.1 | 3.8 | 3.7.5.8 | 3.4 | . | 6.3 | . | 3.2 | 3.6 | 4.0 | 4.0 | . | 2.2 | . | 5.2 | . | . |
| LSM10 | 40.9.1 | 3.8 | 3.7 | 3.4 | 3.6.4.3 | . | 3.6.6.3 | 3.2 | 3.6.3.3 | 4.0 | 4.0 | 3.3.4.7 | 2.2.9.6 | 3.6.3.2 | 3.6.5.2 | 1.3.3.7 | 4.3.6.4 |
| LSM2 | 4.0 | 0 | 3.7 | . | . | 3.6 | 3.6 | . | 3.6.6.3 | 4.0 | 30.4.0 | 4.7 | . | . | . | . | 4.3 |

^aFull-length pspA of strain Rsl.
^bNumbers are size in kilobase pairs.
^cNo hybridization observed with corresponding probe.

The only strain which did not have more than one *pspA*-homologous *Hind*III fragment was WU2, which was previously shown using a full-length *pspA* probe. Even at high stringency, six of the eight probes detected more than one fragment in at least one of the 18 strains (Table 64). LSM7, 10 and 12 hybridized with two fragments in more than one-half of the strains, and the fragments detected by the oligonucleotide probes were identical in size to those detected by the full-length *pspA* probe. Moreover, the same pairs of fragments were frequently detected by probes derived from the 3' as well as the 5' region of Rx1 *pspA*. These results suggested that the *Hind*III fragments from different isolates include two separate but homologous sequences, rather than fragments of a single *pspA* gene. Based on the diversity of the hybridization patterns and the size of restriction fragments, it is clear that *pspA* and *pspC* sequences are highly diverse and that these loci have considerable sequence variability as determined by location of *Hind*III recognition sites.

Oligonucleotides which hybridize with a single restriction fragments in each strain were assumed to be specific for *pspA*. At high stringency, LSM3 and LSM4 detected only a single *Hind*III fragment in the strains with which they reacted. Restriction fragments containing homology to LSM3 or LSM4 were the same as those which hybridize with all of the other homologous probes. This suggested that LSM3 and LSM4 specifically detect *pspA* rather than the *pspC* sequence. That

LSM3 hybridizes with a single restriction fragment of WU2 further confirmed that this oligonucleotide is specific for *pspA*. Sequences from the portion of the gene encoding the second proline region (LSM111) and the C-terminus (LSM2) appeared to be relatively specific for *pspA* since they generally detect only one of the *HindIII* fragments of each strain.

Oligonucleotides LSM12 and LSM10 were able to detect the most conserved epitopes of *pspA* and generally hybridize with multiple restriction fragments of each strain (Table 65). LSM7 was not as broadly cross-reactive, but detected two *pspA*s in 41% of strains including almost 60% of the strains with which it reacts. Thus, sequences representing the leader, first proline region, and the repeat region appear to be relatively conserved not only within *pspA* but between the *pspA* and *pspC* sequences. LSM3, 4, and 5 hybridize with the smallest number of strains of any oligonucleotides (29-35 percent), suggesting that the α -helical domain is the least conserved region within *pspA*. In strains BG58C and L81905 oligonucleotides detect more than two *HindIII* fragments containing sequences with homology to *pspA*. Because of the absence of *HindIII* restriction sites within any of the oligonucleotides it was unlikely that these multiple fragments result from the digestion of chromosomal DNA within the target regions. Also, the additional restriction fragments were detected at high stringency by more than one oligonucleotide. Possibly, in these two strains, there are three or four sequences

with DNA homology to some portions of *pspA*. The probes most consistently reactive with these additional sequences are those for the leader, the alpha-helical region, and the proline-rich region.

The oligonucleotides used as hybridization probes were also tested for their utility as primers in the polymerase chain reaction (PCR). Amplification of *pspA* from 14 strains of *S. pneumoniae* comprising 12 different capsular types was attempted with the primers listed in Table 63. LSM2, derived from the 3' end of *pspA*, were able to amplify an apparent *pspA* sequence from each of 14 pneumococcal strains when used in combination with LSM111, which is within the sequence of *pspA* encoding the proline-rich region. Combinations of LSM2 with primers upstream in *pspA* were variably successful in amplifying sequences (Table 65). The lowest frequency of amplification was observed with LSM112 which was derived from the *Rx1* sequence 5' to the *pspA* start site. This oligonucleotide was not used in the hybridization studies. DNA fragments generated by PCR were blotted and hybridized with a full-length *pspA* probe to confirm homology to *pspA*.

Further evidence for variability at the *pspA* locus comes from the differences in the sizes of the amplified *pspA* gene. When PCR primers LSM12 and LSM2 were used to amplify the entire coding region of *PspA*, PCR products from different pneumococcal isolates ranged in size from 1.9 to 2.3 kbp. The

regions of *pspA* which encode the α -helical, proline-rich, and repeat domains were amplified from corresponding strains and variation in *pspA* appears to come from sequences within the α -helical coding region.

Table 65. Amplification of *pspA* by PCR using the indicated oligonucleotides as 5' primers in combination with the 3' - primer LSM2.

| 5' - primer | Domain | Amplified/ Tested | Percent Amplified |
|-------------|-------------------|----------------------|----------------------|
| LSM112 | -35 (upstream) | 2/14 | 14 |
| LSM12 | leader | 8/14 | 57 |
| LSM3 | α -helical | 3/14 | 21 |
| LSM7 | proline | 12/14 | 86 |
| LSM111 | proline | 14/14 | 100 |

These studies have provided a finer resolution map of the location of conserved and variable sequences within *pspA*. Additionally, regions of divergence and identity between *pspA* and the *pspC* sequences have been identified. This data confirmed serological studies, and demonstrated that *pspA* and *pspC* sequences are highly variable at the DNA sequence level. The diversity of *HindIII* restriction fragment polymorphisms contained *pspA* and the *pspC* sequence supported earlier data using larger probes that detected extensive variability of the DNA in and around these sequences.

A useful *pspA*-specific DNA probe would identify Rx1 and WU2 *pspA* genes, in which restriction maps are known, and would identify only a single restriction fragment in most strains. Two probes, LSM3 and LSM4, do not hybridize with more than one *HindIII* restriction fragment in any strain of pneumococcus. Both of these oligonucleotides hybridize with Rx1 *pspA* and LSM3 hybridizes with WU2 *pspA*. However, each of these probes hybridize with only four of the other 15 strains. When these probes identify a fragment, however, it is generally also detected by all other Rx1-derived probes. Oligonucleotides from the second proline-rich region (LSM111) and the C-terminus of *pspA* (LSM2) generally identify only one *pspA*-homologous sequence at high stringency. Collectively, LSM111, 2, 3 and 4 react with 16 of the 17 isolates and in each case revealed a consensus DNA fragment recognized by most or all of the oligonucleotide probes.

When an oligonucleotide probe detected only a single DNA fragment it was presumed to be *pspA*. If the probe detected multiple fragments, it was presumed to hybridize with *pspA*. If the probe detected multiple fragments, it was presumed to hybridize with *pspA* and the *pspC* sequence. Based on these assumptions the most variable portion between *pspA* and *pspC* is the region immediately upstream from the -35 promoter region and that portion encoding the α -helical region. The most conserved portion between *pspA* and *pspC* was found to be the repeat region, the leader and the proline-rich region sequences. Although only

one probe from within the repeat region was used, the high degree of conservation among the 10 repeats in the Rx1 sequence makes it likely that other probes within the repeat sequences would give similar results.

The portion of Rx1 *pspA* most similar to the *pspC* sequence was that encoding the leader peptide, the upstream portion of the proline-rich region, and the repeat region. The repeat region of PspA has been shown to be involved in the attachment of this protein to the pneumococcal cell surface. The conservation of the repeat region within *pspC* sequences suggests that if these loci encode a protein, it may have a similar functional attachment domain. The conservation of the leader sequence between *pspA* and the *pspC* sequence was also not surprising since similar conservation has been reported for the leader sequence of other proteins from gram positive organisms, such as M protein of group A streptococci (Haanes-Fritz, E. et al., Nucl. Acids Res. 1988; 16: 4667-4677).

In two strain, some oligonucleotide probes identified more than two *pspA*-homologous sequences. In these strains, there was a predominant sequence recognized by almost all of the probes, and two or three additional sequences share homology with DNA encoding the leader, α -helical, and proline region, and they have no homology with sequences encoding the repeat region in the C-terminus of PspA. These sequences might serve as cassettes which can recombine with *pspA* and/or the *pspC* sequences to

generate antigenic diversity. Alternatively, the sequences might encode proteins with very different C-terminal regions and might not be surface attached by the mechanism of PspA.

Oligonucleotides which hybridize with a single chromosomal DNA fragment were used as primers in PCR to examine the variability of domains within *pspA*. These results demonstrate that full-length *pspA* varies in size among strains of pneumococci, and that this variability is almost exclusively the result of sequences in the alpha-helix coding region.

Example 12 - Cloning of PspC

Chromosomal DNA from *S. pneumoniae* EF6796, serotype 6A clinical isolate, was isolated by methods including purification through a cesium chloride gradient, as described in Example 8. The *Hind*III-*Eco*RI fragment of EF6796 was cloned in modified pZero vector (Invitrogen, San Diego, CA) in which the Zeocin-resistance cassette was replaced by a kanamycin cassette (shown in Figure 18). Recombinant plasmids were electroporated into *Escherichia coli* TOP10F' cells [F' {*lacI*^qTet^R} *mcrA* Δ (*mrr*-*hsdRMS*-*mcrBC*) ϕ 80*lacZ* Δ M15 Δ *lacX74* *deoR* *recA1* *araD139* Δ (*ara-leu*)7967 *galU* *galK* *rpsL* *endA1* *nupG*] (Invitrogen).

The 5' region of *pspA*.Rx1 does not hybridize to *pspC* sequence at high stringencies by Southern analysis. Utilizing both the full-length Rx1 *pspA* probe, and a probe containing the sequence encoding α -helical region of PspA, it was possible to identify which DNA fragment contained *pspA* and which fragment

contained the *pspC* locus. The *pspC* locus and the *pspA* gene of EF6796 were mapped using restriction enzymes. After digestion of chromosomal DNA with *HindIII*, the *pspC* locus was localized to a fragment of approximately 6.8 kb. Following a double digest with *HindIII* and *EcoRI*, the *pspC* locus was located in a 3.5 kb fragment. To obtain the intact *pspC* gene of EF6796, chromosomal DNA was digested with *HindIII*, separated by agarose gel electrophoresis, the region between 6 and 7.5 kb purified, and subsequently digested with *EcoRI*. This digested DNA was analyzed by electrophoresis, and DNA fragments of 3.0 to 4.0 kb were purified (GeneClean, Bio101, Inc., Vista, CA). The size-fractionated DNA was then ligated in *HindIII*-*EcoRI*-digested pZero, and electroplated into *E. coli* TOP10F' cells. Kanamycin-resistant transformants were screened by colony blots and probed with full-length *pspA*. A transformant, LXS200, contained a vector with a 3.5 kb insert which hybridized to *pspA*.

Sequencing of *pspC* in pLXS200 was completed using automated DNA sequencing on an ABI 377 (Applied Biosystems, Inc., PLACE). Sequence analyses were performed using the University Of Wisconsin Genetics Computer Group (GCG) programs supported by the Center for AIDS Research (P30 AI27767), MacVector 5.0, Sequencer 2.1, and DNA Strider programs. Sequence similarities of *pspC* were determined using the NCBI BLAST server. The coiled-coil structure predicted by *pspC* sequence was analyzed using Matcher.

A gene probe for cloning the *pspC* locus. Two oligonucleotide primers, N192 and C558 (shown in Figure 19), have been used previously to clone fragments homologous to the region of Rx1 *pspA* encoding amino acids 192-588 from various pneumococcal strains. These primers are modifications (altered restriction sites) of LSM4 and LSM2 which were previously shown to amplify DNA encoding the C-terminal 396 amino acids of PspA.Rx1 (Figure 17); this includes approximately 100 amino acids of the α -helical region, the proline rich region, and the C-terminal choline-binding repeat region. Using primers N192 and C558, a 1.2 kb fragment from strain EF6796 was amplified by PCR, and subsequently cloned in pET-9A (designated PRCT135). This insert was then partially sequenced.

Independently, a larger *pspA* fragment from strain EF6796 was made using primers LSM13 and SKH2 (shown in Figure 19) for the purpose of direct sequencing of serologically diverse *pspA* genes.

The LSM13 and SKH2 primer pair result in the amplification of the 5' end of most *pspA* gene(s) encoding the upstream promoter, the leader peptide, the α -helical, and the proline-rich regions (amino acid -15 to 450) (Figure 20). From the strain EF6796, the LSM13 and SKH2 primers amplified a 1.3 kb fragment (*pspA.EF6796*), which was sequenced. The sequence from PRCT135 and the LSM13/SKH2 PCR-generated fragment *pspA.EF6796* was not identical. The fragment obtained by PCR using primers LSM13

and SKH2 was designated *pspA* based on its location within the same chromosomal location as *pspA.Rx1*. The cloned fragment in pRCT135 was assumed to represent the sequence of the second gene locus, *pspC*, known to be present from Southern analysis. Both genes have significant similarity to the corresponding regions of the prototype *pspA* gene from strain Rx1. The second gene locus was called *pspC*, in recognition of its distinct chromosomal location, not sequence differences from the prototype *pspA* gene.

Analysis of the nucleotide and amino acid sequence of pspC

EF6796. To test the hypothesis that pRCT135 represented *pspC* of EF6796, and to further investigate *pspC*, the entire EF6796 *pspC* gene was cloned as a 3.4 kb *HindIII-EcoRI* fragment forming pLXS200. DNA sequence of the *pspC*-containing clone pLXS200 revealed an open reading frame of 2782 nucleotides based on the analysis of putative transcriptional and translation start and stop sites (Figure 21). The predicted open reading frame encodes a 105 kDa protein which has an estimated pI of 6.09.

PspA.Rx1 and PspC.EF6796 are similar in that they both contain an α -helical region followed by a proline-rich domain and repeat region (Figure 20). However, there are several features of the amino acid sequence of PspC which are quite distinct from PspA. From comparisons at the nucleotide as well as the predicted amino acid sequence, it is apparent that the region of strong homology between PspC and PspA begins at amino acid 458 of

PspC (amino acid 147 of PspA) and extends to the C-terminus of both proteins (positions 899 and 588 respectively). The predicted amino acid sequence of PspC.EF6796 and PspA.Rx1 are 76% similar and 68% identical based on GCG Bestfit program for this region (Figure 22). The nucleotide sequence identity between *pspC* and *pspA* is 87% for the same region. Eight bases upstream of the ATG start site is putative ribosomal binding site, TAGAAGGA. The proposed transcriptional start -35 (TATACA) and -10 (TATAGT) regions are located between 258 to 263 and 280 to 285, respectively (Figure 21). A potential transcriptional terminator occurs at a stem loop between nucleotides 3237 through 3287. The putative signal sequence of PspC is typical of other gram positive bacteria. This region consists of a charged region followed by a hydrophobic core of amino acids. A potential cleavage site of the signal peptide occurs at amino acid 37 following the Val-His-Ala. The first amino acid of the mature protein is a Glu residue.

Other than features similar to all signal sequences, there is no homology in this region between *pspA* and *pspC*. This confirms that *pspC* is present in a separate chromosomal locus from that of *pspA*. The signal sequence and upstream region have striking similarity to the similar regions of *S. agalactiae* β antigen (accession number X59771). The β antigen of Group B streptococci is a cell surface receptor that binds IgA. Similarity to the *bac* gene ends with the start of the mature

protein of PspC, and the nucleotides are 75% identical in this region. Thus, although *pspC* is in a very similar chromosomal locus to the β antigen, it is clearly a distinct protein.

The N-terminus of PspC is quite different from the N-terminus of PspA. Prediction of the secondary structure utilizing Chou-Fausman analysis (Chao, P.Y. et al., Adv. Enzymol. Relat Areas Mol. Biol. 1978: 47: 45-148), suggests that the structure of amino acids 16 to 589 of PspC is predominately α -helical. The Matcher program was used to examine periodicity in the α -helical region of PspA. The characteristic seven residue periodicity is maintained by having hydrophobic residues at the first and fourth positions (a and d) and hydrophobic residues at the remaining positions. The coiled-coil region of the α -helix of PspC (between amino acid 32 to 600) has three breaks in the heptad repeat (Figure 23). These disturbances in the 7 residue periodicity occur at amino acids 99 to 104, 224 to 267 and 346 to 350. The α -helical region of PspA has seven breaks in the motif, each break ranging from a few amino acids to 23 amino acids each. In contrast, the three breaks in the coiled-coil motif of PspC involve 5, 43 and 4 amino acids, respectively.

The sequence encoding the α -helical region of PspC contains two direct repeats 483 nucleotides (160 amino acids) long which are 88% percent identical at the nucleotide level. These repeats, which occur between nucleotides 562 to 1045 and nucleotides 1312 to 1795, are conserved both at the nucleotide

and amino acid level (amino acids 188 to 348 and 438 to 598) (Figure 24). PspA lacks evidence for any repeats this prominent within the α -helical region. These repeat regions could provide a mechanism for recombination that could alter the N-terminal half of the PspC molecule. Although repeat motifs are common in bacterial surface proteins, a direct repeat this large or separated by a large spacer region is novel. The evolutionary significance of this region is not known. A Blast search of the repeat region and the 267 nucleotide bases between them revealed no sequence with significant homology at the nucleotide or amino acid level. However, one of the structural breaks in the coil-coiled region of PspC is the region between the two repeats. Perhaps some deviation from coiled-coil structure between the two repeats is critical to maintain the α -helical structure.

Previous studies have shown that a major cross-protective region of PspA comprises the C-terminal 1/3 of the α -helical region (between residues 192 and 260 of PspA.Rx1). This region accounts for the binding of 4 of 5 cross-protective immunity in mice. Homology between PspC and PspA begins at amino acid 148 of PspA, thus including the region from 192 - 299. The homology between PspA and the PspC includes the entire PspC sequence C-terminal of amino acid 486. Based on the fact that PspA and PspC are so similar in this region known to be protection-eliciting, PspC is also likely to be a protection-eliciting molecule. Because of close sequence and conformational

similarity of the proteins in this region, antibodies specific for the region of PspA between amino acid 148 and 299 should cross-react with PspC and thus afford protection by reacting with PspC and PspA. Likewise, immunization with the PspC would be expected to elicit antibodies cross-protective against PspA. The differences between PspC of strain EF6796 and PspA of strain Rx1 is no greater than the differences between many additional PspAs, which have been shown to be highly cross-protective.

A proline-rich domain exists between amino acid 590 to 652. The sequence, PAPAPEK, is repeated six times in this region. This region is very similar to the proline-rich region of PspA.Rx1 which contains the sequence PAPAP repeated eight times in two proline-rich regions. These two regions of PspA.Rx1 are separated by 27 charge amino acids; no such spacer region is present in PspC.

Many cell surface proteins of other gram positive bacteria contain proline-rich regions. These are often associated with a domain of protein that is predicted to be near the cell wall murein layer when the protein is cell-associated. For example, in M proteins of *S. pyogenes* this domain contains both a Pro- and Gly-rich regions. The fibronectin-binding protein of *S. pyogenes*, *S. dysgalactiae*, and *Staphylococcus aureus* contains a proline-rich region with a three-residue periodicity (pro-charged-uncharged) that is not found in PspA or PspC. An M-like protein of *S. equi* contains a proline-rich

region that is comprised of the tetrapeptide PEPK. This region lacks glycine normally found in the proline regions of M-proteins. The last proline repeat region of this molecule is PAPAK, which is more similar to the proline-region of PspA and PspC than it is to M-proteins.

Proline-rich regions of gram positive bacterial proteins have been reported previously to transit the cell wall. The differences in proline-rich regions of proteins from diverse bacteria may reflect differences in protein function or possibly subtle differences in cell wall function. Proline-rich regions are thought to be responsible for aberrant migration of these proteins through SDS-polyacrylamide gels.

The repeat region of PspC is a common motif found among several proteins in gram positive organisms. Autolysin of *S. pneumoniae*, toxins A and B of *Clostridium difficile*, glucosyltransferases from *S. downei* and *S. mutans*, and CspA of *C. acetobilyticum* all contain similar regions. In PspA these repeats are responsible for binding to the phosphatidylcholine of teichoic acid and lipoteichoic acid in cell wall of pneumococci. However, bacterial proteins containing C-terminal repeats are secreted, which may imply either a lost or gained function. Although all of these proteins have similar repeat regions the similarity of the repeat regions of PspA and PspC is much greater than that of PspC to the other proteins (Table 66).

Interestingly, PspC like PspA has a 17 amino acid partially hydrophobic tail. The function of this 17 amino acid region is unknown. In the case of PspA it has been shown that mutants lacking the tail bind the surface of pneumococci as well as PspAs in which the tail is expressed. Presently, it is now known whether PspC is attached to the cell surface or secreted.

PspA and PspC proteins both have α -helical coiled-coil regions, proline-rich central regions, repeat regions, with a choline binding motifs, and the C-terminal 17 amino acid tail. PspA and PspC share three regions of high sequence identity. One of these is a protection-eliciting region present within the α -helical domain. The other two regions are the proline-rich domain and a repeat domain shared with other choline binding proteins and thought to play a role in cell surface association. The similarity throughout most of the structure of the PspA and the PspC molecules raises the possibility that the two molecules may play at least slightly redundant functions. However, the fact that the N-terminal half of the protein is not homologous to any of the α -helical sequence of PspA suggests the PspC and PspA may have evolved for at least somewhat different roles on the cell surface. One of the most striking differences between the two molecules is the single repeat in the α -helical region of PspC. Although neither the exact function of PspA nor of PspC are known, the observation that a major cross-protective region of PspA is highly homologous with a similar region of PspC,

raises the possibility that both molecules are protection-eliciting and elicit cross-protective antibodies.

The sequence similarity between the promoter region of the *pspC* gene and the *bac* gene from group B streptococci is very intriguing. It implies that an interspecies recombination event has occurred and, this interspecies recombination has contributed to the evolution of the *pspC*. The *pspC* gene thus has a chimeric structure, being partially like *pspA* and partially like the δ antigen. In the latter case, all protein similarity is limited to the signal sequence. Similar interspecies recombination events have contributed to the evolution of the genes encoding penicillin binding protein.

Using analogous procedures, a second PspC sequence was isolated from strain D39 of *S. pneumoniae*. Figures 25 to 29 show the sequence data of PspC from strain D39, complete from upstream of the promoter through the proline-rich region. Strain D39 has the same genetic background as strains Rx1, from which *pspA* was sequenced. D39 and Rx1 have the same *pspC* gene based on Southern blot analysis.

The alpha-helical encoding region of the D39 *pspC* gene is one third of the size of the homologous region from the EF6796 *pspC* gene. The proline-rich region of the D39 *pspC* gene was more similar to Rx1 *pspA* than to EF6796 *pspC*. Even so, the two *pspC* genes were 86% identical at the nucleotide sequence, and 67% identical at the amino acid level.

In the alpha-helical sequence of EF6797 *pspC* a strong repeat was observed. This was absent in the *pspC* sequence of D39. The D39 *pspC* sequence also lacks a leader sequence, found in the EF6797 *pspC* sequence.

This data strongly indicates that there is variability in the structure of *pspC*, similar to previous observations for *pspA*. In the case of *pspC*, however, the extent of variability appears to be even greater than that which has been observed for *pspA*.

Table 66.

| PERCENT HOMOLOGY OF CHOLINE BINDING REGIONS | | | |
|---|--------------------------|--------------------------------|---------|
| | | Percent similarity/identity | |
| Protein | Organism | PspA | PspC |
| PspC | <i>S. pneumoniae</i> | 86/60 | 100/100 |
| Bacteriophage Cp-1 | <i>S. pneumoniae</i> | 56/30 | 56/28 |
| LytA | <i>S. pneumoniae</i> | 57/33 | 61/32 |
| PspA | <i>C. perfringens</i> | 64/45 | 59/42 |
| alpha toxin | <i>C. novyi</i> | 54/29 | 57/33 |
| CspB | <i>C. acetobutylicum</i> | 58/36 | 61/45 |

Having thus described in detail certain preferred embodiments of the present invention, it is to be understood that the invention defined by the appended claims is not to be limited by particular details set forth in the above description, as many apparent variations thereof are possible without departing from the spirit or scope thereof.

REFERENCES

Mufson MA. *Streptococcus pneumoniae*. In: Mandell GL, Douglas RG, Jr, Bennett JE, (eds.) Principles and Practice of Infectious Diseases. New York: Churchill Livingstone, 1990:1539-50.

Cohen C, Parry DAD. alpha-helical coiled coils: more facts and better predictions. Science 1994;236:488-9.

Shapiro ED, Berg AT, Austrian R, et al. Protective efficacy of polyvalent pneumococcal polysaccharide vaccine. N Engl J Med 1991;325:1453-60.

Feldman C, Munro NC, Jeffery PK, et al. Pneumolysin induces the salient histologic features of pneumococcal infection in the rat lung in vivo. Am J Respir Cell Mol Biol 1992;5:416-23.

Lock RA, Paton JC, Hansman D. Comparative efficacy of pneumococcal neuraminidase and pneumolysin as immunogens protective against *Streptococcus pneumoniae*. Microb Pathog 1988;5:461-7.

Sampson JS, O'Connor SP, Stinson AR, Tharpe JA, Russell H. Cloning and nucleotide sequence analysis of *psaA*, the *Streptococcus pneumoniae* gene encoding a 37-kilodalton protein

homologus to previously reported *Streptococcus* sp. adhesins.
Infect Immun 1994;62:319-24.

McDaniel LS, Sheffield JS, Delucchi P, Briles DE. PspA, a surface protein of *Streptococcus pneumoniae*, is capable of eliciting protection against pneumococci of more than one capsular type. *Infect Immun* 1991;59:222-8.

Berry AM, Lock RA, Hansman D, Paton JC. Contribution of autolysin to virulence of *Streptococcus pneumoniae*. *Infect Immun* 1989;57:2324-30.

Berry AM, Yother J, Briles DE, Hansman D, Paton JC. Reduced virulence of a defined pneumolysin-negative mutant of *Streptococcus pneumoniae*. *Infect Immun* 1989;57:2037-42.

McDaniel LS, Yother J, Vijayakumar M, McGarry L, Guild WR, Briles DE. Use of insertional inactivation to facilitate studies of biological properties of pneumococcal surface protein A (PspA). *J Exp Med* 1987;165:381-94.

Waltman WD II, McDaniel LS, Gray BM, Briles DE. Variation in the molecular weight of PspA (Pneumococcal Surface Protein A) among *Streptococcus pneumoniae*. *Microb Pathog* 1990;8:61-9.

Crain MJ, Waltman WD II, Turner JS, et al. Pneumococcal surface protein A (PspA) is serologically highly variable and is expressed by all clinically important capsular serotypes of *Streptococcus pneumoniae*. *Infect Immun* 1990;58:3293-9.

Yother J, Briles DE. Structural properties and evolutionary relationships of PspA, a surface protein of *Streptococcus pneumoniae*, as revealed by sequence analysis. *J Bact* 1992;174:601-9.

McDaniel LS, Scott G, Kearney JF, Briles DE. Monoclonal antibodies against protease sensitive pneumococcal antigens can protect mice from fatal infection with *Streptococcus pneumoniae*. *J Exp Med* 1984;160:386-97.

McDaniel LS, Ralph BA, McDaniel DO, Briles DE. Localization of protection-eliciting epitopes on PspA of *Streptococcus pneumoniae* between amino acids residues 192 and 260. *Microb Path* 1994;17:323-37.

Yother J, Forman C, Gray BM, Briles DE. Protection of mice from infection with *Streptococcus pneumoniae* by anti-phosphocholine antibody. *Infect Immun* 1982;36:184-8.

Briles DE, Crain MJ, Gray BM, Forman C, Yother J. Strong association between capsular type and virulence for mice among human isolates of *Streptococcus pneumoniae*. Infect Immun 1992;60:111-6.

McDaniel LS, Sheffield JS, Swiatlo E, Yother J, Crain MJ, Briles DE. Molecular localization of variable and conserved regions of *pspA*, and identification of additional *pspA* homologous sequences in *Streptococcus pneumoniae*. Microb Pathog 1992;13:261-9.

McDaniel LS, McDaniel DO. Analysis of the gene encoding type 12 PspA of *S. pneumoniae* EF5668. In: Ferretti JJ, Gilmore MS, Klaenhammer TR, Brown F ed. Genetics of Streptococci, Enterococci and Lactococci. Basel: Karger, 1995:283-6.

Briles DE, Forman C, Crain M. Mouse antibody to phosphocholine can protect mice from infection with mouse-virulent human isolates of *Streptococcus pneumoniae*. Infect Immun 1992;60:1957-62.

Davis RW, Boststein D, Roth JR. A manual for genetic engineering: advanced bacterial genetics. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press, 1980.

Studier FW, Moffatt BA. Use of bacteriophage T7 RNA polymerase to direct selective high-level expression of cloned genes. *J Mol Biol* 1986;189:113-30.

Hanahan D. Studies on transformation of *Escherichia coli* with plasmids. *J Mol Biol* 1983;166:557-80.

Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970;227:680-5.

Towbin H, Staehelin T, Gordon J. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *PNAS* 1979;76:4350-4.

Amsbaugh DF, Hansen CT, Prescott B, Stashak PW, Barthold DR, Baker PJ. Genetic control of the antibody response to type III pneumococcal polysaccharide in mice. I. evidence that an X-linked gene plays a decisive role in determining responsiveness. *J Exp Med* 1972;136:931-49.

Briles DE, Nahm M, Schroer K, et al. Antiphosphocholine antibodies found in normal mouse serum are protective against intravenous infection with type 3 *Streptococcus pneumoniae*. *J Exp Med* 1981;153:694-705.

Zar JH. Biostatistical Analysis. 2nd Ed. Englewood Cliffs, N. J.: Prentice-Hall, Inc., 1984:718.

Yother J, Handsome GL, Briles DE. Truncated forms of PspA that are secreted from *Streptococcus pneumoniae* and their use in functional studies and cloning of the *pspA* gene. J Bact 1992;174:610-8.

Talkington DF, Voellinger DC, McDaniel LS, Briles DE. Analysis of pneumococcal PspA microheterogeneity in SDS polyacrylamide gels and the association of PspA with the cell membrane. Microb Pathog 1992;13:343-55.

Talkington DF, Crimmins DL, Voellinger DC, Yother J, Briles DE. A 43-kilodalton pneumococcal surface protein, PspA: isolation, protective abilities, and structural analysis of the amino-terminal sequence. Infect Immun 1991;59:1285-9.

Schneewind O, Model P, Fischetti VA. Sorting of protein A to the staphylococcal cell wall. Cell 1992;70:267-81.

Yother J, White JM. Novel surface attachment mechanism for the streptococcus pneumoniae protein PspA. J Bact 1994;176:2976-85.

Gray BM. Pneumococcal infection in an era of multiple antibiotic resistance. *Adv Ped Inf Dis* 1995;In press.

Filice G. A., L. L. Van Etta, C. P. Darby and D. W. Fraser. 1986. Bacteremia in Charleston County, South Carolina. *Am. J. Epidemiol.* 123:128.

Gillespie S. H. 1989. Aspects of pneumococcal infection including bacterial virulence, host response and vaccination. *J. Med. Microbiol.* 28:237.

Musher D. M. 1992. Infections caused by *Streptococcus pneumoniae*: Clinical spectrum, pathogenesis, immunity, and treatment. *Clin. Infect. Dis.* 14:801.

Nordenstam G., B. Anderson, D. E. Briles, J. Brooks, A. Oden, A. Svanborg and C. S. Eden. 1990. High anti-phosphorylcholine antibody levels and mortality associated with pneumonia. *Scand. J. Infect. Dis.* 22:187.

Giebink G. S. 1989. The microbiology of otitis media. *Pediatr. Infect. Dis. J.* 8:S18.

Giebink G. S. 1985. Preventing pneumococcal disease in children: recommendations for using pneumococcal vaccine. *Pediatr. Infect Dis.* 4:343.

Siber G. R. 1994. Pneumococcal disease: prospects for a new generation of vaccines. *Science* 265:1385.

Cadoz M., J. Armand, F. Arminjon, J.-P. Michel, M. Michel, F. Denis and G. Schiffman. 1985. A new 23 valent pneumococcal vaccine: immunogenicity and reactogenicity in adults. *J. Biol. Stand.* 13:261.

Robbins J. B., R. Austrian, C.-J. Lee, S. C. Rastogi, G. Schiffman, J. Henrichsen, P. H. Makela, C. V. Broome, R. R. Facklam, R. H. Tiesjema and J. C. Parke Jr. 1983. Considerations for formulating the second-generation pneumococcal capsular polysaccharide vaccine with emphasis on the cross-reactive types within groups. *J. Infect. Dis.* 148:1136.

Forrester H. L., D. W. Jahigen and F. M. LaForce. 1987. Inefficacy of pneumococcal vaccine in a high-risk population. *Am. J. Med.* 83:425.

Douglas R. M. and H. B. Miles. 1984. Vaccination against *Streptococcus pneumoniae* in childhood: lack of demonstrable benefit in young Australian children. *J. Infect. Dis.* 149:861.

Douglas R. M., J. C. Paton, S. J. Duncan and D. J. Hansman. 1983. Antibody response to pneumococcal vaccination in children younger than five years of age. *J. Infect. Dis.* 148:131.

Leinonen M., A. Sakkinen, R. Kallioikoski, J. Luotenen, M. Timonen and P. H. Mekela. 1986. Antibody response to 14-valent pneumococcal capsular polysaccharide vaccine in preschool age children. *Pediatr. Infect. Dis.* 5:39.

Makela P. H., M. Leinonen, J. Pukander and P. Karma. 1981. A study of the pneumococcal vaccine in prevention of clinically acute attacks of recurrent otitis media. *Rev. Infect. Dis.* 3:S124.

Riley I. D. and R. M. Douglas. 1981. An epidemiologic approach to pneumococcal disease. *Rev. Infect. Dis.* 3:233.

Wright P. F., S. H. Sell, W. K. Vaughn, C. Andrews, K. B. McConnell and G. Schiffman. 1981. Clinical studies of pneumococcal vaccines in infants. II. Efficacy and effect on nasopharyngeal carriage. *Rev. Infect. Dis.* 3:S108.

Lock R. A., J. C. Paton and D. Hansman. 1988. Comparative efficacy of pneumococcal neuraminidase and pneumolysin as immunogens protective against *Streptococcus pneumoniae*. *Microbial Pathogenesis* 5:461.

McDaniel L. S. and D. E. Briles. 1986. Monoclonal antibodies against bacteria. Orlando, Fla: Academic Press, Inc., 143.

Paton J. C., R. A. Lock and D. J. Hansman. 1983. Effect of immunization with pneumolysin on survival time of mice challenged with *Streptococcus pneumoniae*. *Infect. Immun.* 40:548.

Talkington D. F., D. L. Crimmins, D. C. Voellinger, J. Yother and D. E. Briles. 1991. A 43-kilodalton pneumococcal surface protein, PspA: isolation, protective abilities, and structural analysis of the amino-terminal sequence. *Infect. Immun.* 59:1285.

Yother J., C. Forman, B. M. Gray and D. E. Briles. 1982. Protection of mice from infection with *Streptococcus pneumoniae* by anti-phosphocholine antibody. *Infect. Immun.* 36:184.

Crain M. J., W. D. Waltman, J. S. Turner, J. Yother, D. F. Talkington, L. S. McDaniel, B. M. Gray and D. E. Briles. 1990. Pneumococcal surface protein A (PspA) is serologically highly variable and is expressed by all clinically important capsular serotypes of *Streptococcus pneumoniae*. *Infect. Immun.* 58:3293.

Briles D. E., J. Yother and L. S. McDaniel. 1988. Role of pneumococcal surface protein A in the virulence of *Streptococcus pneumoniae*. *Rev. Infect. Dis.* 10:S372.

McDaniel L. S., J. Yother, M. Vijayakumar, L. McGarry, W. R. Guild and D. E. Briles. 1987. Use of insertional inactivation to facilitate studies of biological properties of pneumococcal surface protein A (PspA). *J. Exp. Med.* 165:381.

Waltman W. D., L. S. McDaniel, B. M. Gray and D. E. Briles. 1990. Variation in the molecular weight of PspA (pneumococcal surface protein A) among *Streptococcus pneumoniae*. *Microbial Pathogenesis* 8:61.

Yother J. and D. E. Briles. 1992. Structural properties and evolutionary relationships of PspA, a surface protein of *Streptococcus pneumoniae*, as revealed by sequence analysis. *J. Bacteriol.* 174:601.

Yother J. and J. M. White. 1994. Novel surface attachment mechanism for the *Streptococcus pneumoniae* protein PspA. *J. Bact.* 176:2976.

Yother J., G. L. Handsome and D. E. Briles. 1992. Truncated forms of PspA that are secreted from *Streptococcus pneumoniae* and their use in functional studies and cloning of the *pspA* gene. *J. Bacteriol.* 174:610.

McDaniel L. S., J. S. Sheffield, P. Delucchi and D. E. Briles. 1991. PspA, a surface protein of *Streptococcus pneumoniae*, is capable of eliciting protection against pneumococci of more than one capsular type. *Infect. Immun.* 59:222.

McDaniel L. S., B. A. Ralph, D. O. McDaniel and D. E. Briles. 1994. Localization of protection-eliciting epitopes on PspA of *Streptococcus pneumoniae* between amino acid residues 192 and 260. *Micro. Pathogenesis* 17:323.

McDaniel L. S., K. Scott, J. F. Kearney and D. E. Briles. 1984. Monoclonal antibodies against protease sensitive pneumococcal antigens can protect mice from fatal infection with *Streptococcus pneumoniae*. *J. Exp. Med.* 160:386.

Davis R. W., W. D. Boststein and J. R. Roth. 1980. A manual for genetic engineering: Advanced bacterial genetics. . Cold Spring Harbor, New York: Cold Spring Harbor Laboratory, 201.

Hanahan D. 1983. Studies on transformation of *Escherichia coli* with plasmids. *J. Mol. Biol.* 166:557.

Birnboim H. C. and J. Doly. 1979. A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nuc. Acids Res.* 7:1513.

Osborn M. J. and J. Munson. 1974. Separation of the inner (cytoplasmic) and outer membranes of gram negative bacteria. *Methods Enzymol.* 31A:642.

Wicker L. S. and I. Scher. 1986. X-linked immune deficiency (*Xid*) of CBA/N mice . New York: Apringer-Verlag, 86.

Amsbaugh D. F., C. T. Hansen, B. Prescott, P. W. Stashak, D. R. Barthold and P. J. Baker. 1972. Genetic control of the antibody response to type III pneumococcal polysaccharide in mice. I. Evidence that an X-linked gene plays a decisive role in determining responsiveness. *J. Exp. Med.* 136:931.

Briles D. E., M. Nahm, K. Schoer, J. Davie, P. Baker, J. F. Kearney and R. Barletta. 1981. Anti-phosphocholine antibodies found in normal mouse serum are protective against intravenous infection with type 3 *S. pneumoniae*. *J. Exp. Med.* 153:694.

McDaniel L. S., J. S. Sheffield, E. Swiatlo, J. Yother, M. J. Crain and D. E. Briles. 1992. Molecular localization of variable and conserved regions of *pspA* and identification of additional *pspA* homologous sequences in *Streptococcus pneumoniae*. *Microbial Pathogenesis* 13:261.

Alexander, J.E., Lock, R.A., Peeters, C.C.A.M., Poolman, J.T., Andrew, P.W., Mitchell, T.J., Hansman, D., and Paton, J.C. (1994) Immunization of mice with pneumolysin toxoid confers a significant degree of protection against at least nine serotypes of *Streptococcus pneumoniae*. *Infect Immun* 62: 5683-5688.

Avery, O.T., McLeod, C.M., and McCarty, M. (1944) Studies on the chemical nature of the substance inducing transformation of pneumococcal types. Induction of transformation by a desoxyribonucleic acid fraction isolated from pneumococcus type III. *J Exp Med* 79: 137-158.

Briles, D.E., Nahm, M., Schroer, K., Davie, J., Baker, P., Kearney, J.F., and Barletta, R. (1981) Anti-phosphocholine

antibodies found in normal mouse serum are protective against intravenous infection with type 3 *S. pneumoniae*. *J Exp Med* 153: 694-705.

Crain, M.J., Waltman, W.D. II, Turner, J.S., Yother, J., Talkington, D.E., McDaniel, L.M., Gray, B.M., and Briles, D.E. (1990) Pneumococcal surface protein A (PspA) is serologically highly variable and is expressed by all clinically important capsular serotypes of *Streptococcus pneumoniae*. *Infect Immun* 58: 3293-3299.

Haanes-Fritz, E., Kraus, W., Burdett, V., Dale, J.B., Beachey, E.H., and Cleary, P. (1988) Comparison of the leader sequences of four group A streptococcal M protein genes. *Nucl Acids Res* 16: 4667-4677.

McDaniel, L.S., Yother, J., Vijayakumar, M., McGarry, L., Guild, W.R., and Briles, D.E. (1987) Use of insertional inactivation to facilitate studies of biological properties of pneumococcal surface protein A (PspA). *J Exp Med* 165: 381-394.

McDaniel, L.S., Sheffield, J.S., Delucchi, P., and Briles, D.E. (1991) PspA, a surface protein of *Streptococcus pneumoniae*, is capable of eliciting protection against pneumococci of more than one capsular type. *Infect Immun* 59: 222-228.

McDaniel, L.S., Sheffield, J.S., Swiatlo, E., Yother, J. Crain, M.J., and Briles, D.E. (1992) Molecular localization of variable and conserved regions of *pspA*, and identification of additional *pspA*-homologous sequences in *Streptococcus pneumoniae*. *Microbial Pathogenesis* 13: 261-269.

McDaniel, L.S., Ralph, B.A., McDaniel, D.O., and Briles, D.E. (1994) Localization of protection-eliciting epitopes on PspA of *Streptococcus pneumoniae* between amino acid residues 192 and 260. *Microbial Pathogenesis* 17: 323-337.

Meinkoth, J., and Wahl, G. (1984) Hybridization of nucleic acids immobilized on solid supports. *Anal Biochem* 138: 267-284.

Sampson, J.S., O'Connor, S.P., Stinson, A.R., Tharpe, J. A., and Russell, H. (1994) Cloning and nucleotide sequence analysis of *psaA*, the *Streptococcus pneumoniae* gene encoding a 37-kilodalton protein homologous to previously reported *Streptococcus* sp. adhesins. *Infect Immun* 62: 319-324.

Shoemaker, N.B., and Guild, W.R. (1974) Destruction of low efficiency markers is a slow process occurring at a heteroduplex stage of transformation. *Mol Gen Genet* 128: 283-290.

Siber, G.R. (1994) Pneumococcal disease: prospects for a new generation of vaccines. *Science* 265: 1385-1387.

Talkington, D.F., Crimmins, D.L., Voellinger, D.C., Yother, J., and Briles, D.E. (1991) A 43-kilodalton pneumococcal surface protein, PspA: isolation, protective abilities, and structural analysis of the amino-terminal sequence. *Infect Immun* 59: 1285-1289.

Waltman, W.D. II, McDaniel, L.S., Gray, B.M., and Briles, D.E. (1990) Variation in the molecular weight of PspA (pneumococcal surface protein A) among *Streptococcus pneumoniae*. *Microbial Pathogenesis* 8: 61-69.

Yother, J., McDaniel, L.S., and Briles, D.E. (1986) Transformation of encapsulated *Streptococcus pneumoniae*. *J Bacteriol* 168: 1463-1465.

Yother, J., and Briles, D.E. (1992) Structural properties and evolutionary relationships of PspA, a surface protein of *Streptococcus pneumoniae*, as revealed by sequence analysis. *J Bacteriol* 174: 601-609.

Yother, J., Handsome, G.L., and Briles, D.E. (1992)
Truncated forms of PspA that are secreted from *Streptococcus pneumoniae* and their use in functional studies and cloning of the *pspA* gene. *J Bacteriol* 174: 610-618.

Yother, J., and White, J.M. (1994) Novel surface attachment mechanism of the *Streptococcus pneumoniae* protein PspA. *J Bacteriol* 176: 2976-2985.

Anonymous. Pneumococcal polysaccharide vaccine. *MMWR* 1981, 30, 410-419

Farley, J.J., King, J.C., Nair, P., Hines, S.E., Tressler, R.L., Vink, P.E. Invasive pneumococcal disease among infected and uninfected children of mothers with immunodeficiency virus infection. *J. Pediatr.* 1994, 124, 853-858

Schwartz, B., Gove, S., Lob-Lovit, J., Kirkwood, B.R.
Potential interactions for the prevention of childhood pneumonia in developing countries: etiology of acute lower respiratory infections among young children in developing countries. *Ped. Infect. Dis.* in Press,

Avery, O.T., Goebel, W.F. Chemoimmunological studies of the soluble specific substance of pneumococcus. I. The isolation and

properties of the acetyl polysaccharide of pneumococcus type 1.

J. Exp. Med. 1933, 58, 731 - 755

Austrian, R. Pneumococcal Vaccine: Development and Prospects. *Am. J. Med* 1979, 67, 547-549

Shapiro, E.D., Berg, A.T., Austrian, R., Schroeder, D.,
Parcells, V., Margolis, A., Adair, R.K., Clemmens, J.D.
Protective efficacy of polyvalent pneumococcal polysaccharide
vaccine. *N. Engl. J. Med* 1991, 325, 1453-1460

Fedson, D.S. Pneumococcal vaccination in the prevention of
community-acquired pneumonia: an optimistic view of cost-
effectiveness. *Sem. Resp. Infect.* 1993, 8, 285-293

Gotschlich, E.C., Goldschneider, I., Lepow, M.L., Gold, R.
The immune response to bacterial polysaccharides in man. In:
Antibodies in human diagnosis and therapy, (Ed. Haber, E.,
Krause, R.M.) Raven, New York, 1977, 391-402.

Cowan, M.J., Ammann, A.J., Wara, D.W., Howie, V.M., Schultz,
L., Doyle, N., Kaplan, M. Pneumococcal polysaccharide
immunization in infants and children. *Pediatrics* 1978, 62, 721-727

Mond, J.J., Lees, A., Snapper, C.M. T cell-independent antigens type 2. *Ann. Rev. Immunol.* 1995, 13, 655-692

Stein, K.E. Thymus-independent and thymus-dependent responses to polysaccharide antigens. *J. Infect. Dis.* 1992, 162, S49

Chiu, S.S., Greenberg, P.D., Marcy, S.M., Wong, V.K., Chang, S.J., Chiu, C.Y., Ward, J.I. Mucosal antibody responses in infants following immunization with *Haemophilus influenzae*. *Pediatr. Res. Abstracts* 1994, 35, 10A

Kauppi, M., Eskola, J., Kathy, H. H. influenzae type b (Hib) conjugate vaccines induce mucosal IgA1 and IgA2 antibody responses in infants and children. *ICAAC Abstracts* 1993, 33, 174

Dagen, R., Melamed, R., Abramson, O., Piglansky, L., Greenberg, D., Mendelman, P.M., Bohidar, N., Ter-Minassian, D., Cvanovich, N., Lov, D., Rusk, C., Donnelly, J., Yagupsky, P. Effect of heptavalent pneumococcal-OMPC conjugate vaccine on nasopharyngeal carriage when administered during the 2nd year of life. *Pediatr. Res.* 1995, 37, 172A

Fattom, A., Vann, W.F., Szu, S.C., Sutton, A., Bryla, D., Shiffman, G., Robbins, J.B., Schneerson, R. Synthesis and physiochemical and immunological characterization of pneumococcus type 12F polysaccharide-diphtheria toxoid conjugates. *Infect. Immun.* 1988, 56, 2292-2298

Kennedy, D., Derousse, C., E., A. Immunologic response of 12 -18 month old children to licensed pneumococcal polysaccharide vaccine primed with *Streptococcus pneumoniae* 19F conjugate vaccine. *ICAAC* 1994, 34th annual meeting, 236

McDaniel, L.S., Ralph, B.A., McDaniel, D.O., Briles, D.E. Localization of protection-eliciting epitopes on PspA of *Streptococcus pneumoniae* between amino acid residues 192 and 260. *Microb. Pathog.* 1994, 17, 323-337

Langermann, S., Palaszynski, S.R., Burlein, J.E., Koenig, S., Hanson, M.S., Briles, D.E., Stover, C.K. Protective humoral response against pneumococcal infection in mice elicited by recombinant Bacille Calmette-Guérin vaccines expressing PspA. *J. Exp. Med.* 1994, 180, 2277-2286

Siber, G.R. Pneumococcal disease: prospects for a new generation of vaccines. *Science* 1994, 265, 1385-1387

Lock, R.A., Hansman, D., Paton, J.C. Comparative efficacy of autolysin and pneumolysin as immunogens protecting mice against infection by *Streptococcus pneumoniae*. *Microb. Pathog.* 1992, 12, 137-143

Sampson, J.S., O'Connor, S.P., Stinson, A.R., Tharpe, J.A., Russell, H. Cloning and nucleotide sequence analysis of *psaA*, the *Streptococcus pneumoniae* gene encoding a 37-kilodalton protein homologous to previously reported *Streptococcus* sp. adhesins. *Infect. Immun.* 1994, 62, 319

Paton, J.C., Lock, R.A., Lee, C.-J., Li, J.P., Berry, A.M., Mitchell. Purification and immunogenicity of genetically obtained pneumolysin toxoids and their conjugation to *Streptococcus pneumoniae* type 19F polysaccharide. *Infect. Immun.* 1991, 59, 2297-2304

McDaniel, L.S., Scott, G., Kearney, J.F., Briles, D.E. Monoclonal antibodies against protease sensitive pneumococcal antigens can protect mice from fatal infection with *Streptococcus pneumoniae*. *J. Exp. Med.* 1984, 160, 386-397

Briles, D.E., Forman, C., Horowitz, J.C., Volanakis, J.E., Benjamin, W.H., Jr., McDaniel, L.S., Eldridge, J., Brooks, J. Antipneumococcal effects of C-reactive protein and monoclonal antibodies to pneumococcal cell wall and capsular antigens. *Infect. Immun.* 1989, 57, 1457 - 1464

McDaniel, L.S., Sheffield, J.S., Delucchi, P., Briles, D.E. PspA, a surface protein of *Streptococcus pneumoniae*, is capable of eliciting protection against pneumococci of more than one capsular type. *Infect. Immun.* 1991, 59, 222-228

McDaniel, L.S., Yother, J., Vijayakumar, M., McGarry, L., Guild, W.R., Briles, D.E. Use of insertional inactivation to facilitate studies of biological properties of pneumococcal surface protein A (PspA). *J. Exp. Med.* 1987, 165, 381-394

Yother, J., McDaniel, L.S., Crain, M.J., Talkington, D.F., Briles, D.E. Pneumococcal surface protein A: Structural analysis and biological significance In: *Genetics and Molecular Biology of Streptococci, Lactococci, and Enterococci*, (Ed. Dunny, G.M., Cleary, P.P., McKay, L.L.) American Society for Microbiology, Washington, DC, 1991, 88-91.

Waltman, W.D., II, McDaniel, L.S., Gray, B.M., Briles, D.E. Variation in the molecular weight of PspA (Pneumococcal Surface Protein A) among *Streptococcus pneumoniae*. *Microb. Pathog.* 1990, 8, 61-69

Crain, M.J., Waltman, W.D., II, Turner, J.S., Yother, J., Talkington, D.E., McDaniel, L.M., Gray, B.M., Briles, D.E. Pneumococcal surface protein A (PspA) is

serologically highly variable and is expressed by all clinically important capsular serotypes of *Streptococcus pneumoniae*. *Infect. Immun.* 1990, 58, 3293-3299

McDaniel, L.S., Scott, G., Widenhofer, K., Carroll, Briles, D.E. Analysis of a surface protein of *Streptococcus pneumoniae* recognized by protective monoclonal antibodies. *Microb. Pathog.* 1986, 1, 519-531

Tart, R.C., McDaniel, L.S., Ralph, B.A., Briles, D.E. Truncated *Streptococcus pneumoniae* PspA molecules elicit cross-protective immunity against pneumococcal challenge in mice. *J. Infect. Dis.* 1995, In Press,

Yother, J., Briles, D.E. Structural properties and evolutionary relationships of PspA, a surface protein of *Streptococcus pneumoniae*, as revealed by sequence analysis. *J. Bact.* 1992, 174, 601-609

Talkington, D.F., Crimmins, D.L., Voellinger, D.C., Jother, J., Briles, D.E. A 43-kilodalton pneumococcal surface protein, PspA: isolation, protective abilities, and structural analysis of the amino-terminal sequence. *Infect. Immun.* 1991, 59:, 1285-1289

McDaniel, L.S., McDaniel, D.O. Genetic analysis of the gene encoding type 12 PspA of *Streptococcus pneumoniae* strain EF5668 In: *Genetics of the streptococci*,

enterococci, and lactococci, (Ed. Feretti, J.J., Gilmore, M.S., Khenhammer, T.R., Brown, F.) Dev. Biol. Stand. Basel Krager, Basel, 1995, 283-286.

Fischetti, V.A., Pancholi, V., Schneewind, O. Conservation of a hexapeptide sequence in the anchor region of surface proteins from gram-positive cocci. *Mol. Microbiol.* 1990, 4, 1603-1605

Schneewind, O., Fowler, A., Faull, K.F. Structure of cell wall anchor of cell surface proteins in *Staphylococcus aureus*. *Science* 1995, 268, 103-106

Yother, J., White, J.M. Novel surface attachment mechanism for the *Streptococcus pneumoniae* protein PspA. *J. Bact.* 1994, 176, 2976-2985

McDaniel, L.S., Brooks-Walter, A., Briles, D.E., Swiatlo, E. Oligonucleotides identify conserved and variable regions of *pspA* and *pspA*-like sequences of *Streptococcus pneumoniae*. *Mol. Microbiol.* Submitted,

Yother, J., Handsome, G.L., Briles, D.E. Truncated forms of PspA that are secreted from *Streptococcus pneumoniae* and their use in functional studies and cloning of the *pspA* gene. *J. Bact.* 1992, 174, 610-618

Talkington, D.F., Voellinger, D.C., McDaniel, L.S., Briles, D.E. Analysis of pneumococcal PspA microheterogeneity in SDS polyacrylamide gels and the association of PspA with the cell membrane. *Microb. Pathog.* 1992, 13, 343-355

Smith, M.D., Guild, W.R. A plasmid in *Streptococcus pneumoniae*. *J. Bacteriol.* 1979, 137, 735-739

Shoemaker, N.B., Guild, W.R. Destruction of low efficiency markers is a slow process occurring at a heteroduplex stage of transformation. *Mol. Gen. Genet.* 1974, 128, 283-290

Raven, A.W. Reciprocal capsular transformations of pneumococci. *J. Bact.* 1959, 77, 296-309

McDaniel, L.S., Sheffield, J.S., Swiatlo, E., Yother, J., Crain, M.J., Briles, D.E. Molecular localization of variable and conserved regions of *pspA*, and identification of additional *pspA* homologous sequences in *Streptococcus pneumoniae*. *Microb. Pathog.* 1992, 13, 261-269

Brooks-Walter, A., McDaniel, L.S., Hollingshead, S.K., Briles, D.E. Restriction fragment length polymorphisms of *pspA* of *Streptococcus pneumoniae* reveal a genetic polymorphism. *Submitted*

van de Rijn, I., Kessler, R.E. Growth characteristics of Group A Streptococci in a new chemically defined medium. *Infect. Immun.* 1980, 27, 444-448

Waltman, W.D., II, McDaniel, L.S., Andersson, B., Bland, L., Gray, B.M., Svanborg-Eden, C., Briles, D.E. Protein serotyping of *Streptococcus pneumoniae* based on reactivity to six monoclonal antibodies. *Microb. Pathog.* 1988, 5, 159-167

Tomasz, A. Surface components of *Streptococcus pneumoniae*. *Rev. Infect. Dis* 1981, 3, 190-211

Garcia, J.L., Garcia, E., Lopez, R. Overproduction and rapid purification of the amidase of *Streptococcus pneumoniae*. *Arch. Microbiol.* 1987, 149, 52-56

Osborn, M.J., Munson, J. Separation of the inner (cytoplasmic) and outer membranes of gram negative bacteria. *Methods Enzymol.* 1974, 31A, 642-653

Briles, D.E., Horowitz, J., McDaniel, L.S., Benjamin, W.H., Jr., Claflin, J.L., Booker, C.L., Scott, G., Forman, C. Genetic control of susceptibility to pneumococcal infection. *Curr. Top. Microbiol. Immunol.* 1986, 124, 103-120

Briles, D.E., Crain, M.J., Gray, B.M., Forman, C., Yother, J. A strong association between capsular type and mouse virulence among human isolates of *Streptococcus pneumoniae*. *Infect. Immun.* 1992, 60, 111-116

Musher, D.M., Raizan, K.R., Weinstein, L. The effect of *Listeria monocytogenes* on resistance to pneumococcal infection. *Soc. Exp. Bio. and Med.* 1970, 135, 557-560

Roberts, P., Jeffery, P.K., Mitchell, T.J., Andrew, P.W., Boulnois, G.J., Feldman, C., Cole, P.J., Wilson, R. Effect of immunization with Freund's adjuvant and pneumolysin on histologic features of pneumococcal infection in the rat lung in vivo. *Infect. Immun.* 1992, 60, 4969-4972

Weigle, W.O. Immunological unresponsiveness In: *Adv. Immunol.*, (Ed. Dixon, J.F., Kunkel, H.G.) Academic Press, New York, New York, 1973, 61-162.

Alexander, J.E., Lock, R.A., Peeters, C.C.A.M., Poolman, J.T., Andrew, P.W., Mitchell, T.J., Hansman, D., Paton, J.C. Immunization of mice with pneumolysin toxoid confers a significant degree of protection against at least nine serotypes of *Streptococcus pneumoniae*. *Infection and Immunity* 1994, 62, 5683-5688

Berry, A.M., Lock, R.A., Hansman, D., Paton, J.C. Contribution of autolysin to virulence of *Streptococcus pneumoniae*. *Infect. Immun.* 1989, 57, 2324-2330

Lock, R.A., Paton, J.C., Hansman, D. Purification and immunologic characterization of neuraminidase produced by *Streptococcus pneumoniae*. *Microb. Pathog.* 1988, 4, 33-43

Talkington, D., Koenig, A., Russell, H. The 37 kDa protein of *Streptococcus pneumoniae* protects mice against fatal challenge. *American Society of Microbiology Abstracts* 1992, 149

Dillard, J.P., Yother, J. Genetic and molecular characterization of capsular polysaccharide biosynthesis in *Streptococcus pneumoniae* type 3. *Mol. Microbiol.* 1994, 12, 959-972

Tomasz, A. Biological consequences of the replacement of choline by ethanolamine in the cell wall of pneumococcus: chain formation, loss of transformability, and loss of autolysis. *Proc. Natl. Acad. Sci. USA* 1968, 59, 86-93

Briles, D.E., Nahm, M., Schroer, K., Davie, J., Baker, P., Kearney, J., Barletta, R. Antiphosphocholine antibodies found in normal mouse serum are protective against

intravenous infection with type 3 *Streptococcus pneumoniae*. *J. Exp. Med.* 1981, 153, 694-705

Amsbaugh, D.F., Hansen, C.T., Prescott, B., Stashak, P.W., Barthold, D.R., Baker, P.J. Genetic control of the antibody response to type III pneumococcal polysaccharide in mice. I. Evidence that an X-linked gene plays a decisive role in determining responsiveness. *J. Exp. Med.* 1972, 136, 931-949

Avery, O.T., MacLeod, C.M., McCarty, M. Studies on the chemical nature of the substance inducing transformation of pneumococcal types. Induction of transformation by a desoxyribonucleic acid fraction isolated from pneumococcus type III. *J. Exp. Med.* 1944, 79, 137-158

McCarty, M. *The transforming principle*. Norton, New York, 1985, 252.

WHAT IS CLAIMED IS:

1. An isolated amino acid molecule consisting of residues 1 to 115, 1 to 260, 192 to 588, 192 to 299, or residues 192 to 260 of pneumococcal surface protein A of *Streptococcus pneumoniae*.

2. An isolated DNA molecule consisting of a fragment of pneumococcal surface protein A gene of *Streptococcus pneumonia* encoding the isolated amino acid molecule of claim 1.

3. A PCR primer consisting essentially of the isolated DNA molecule of claim 2.

4. A hybridization probe consisting essentially of the isolated DNA molecule of claim 2.

5. An immunological composition comprising the amino acid molecule of claim 1.

6. An isolated DNA molecule consisting of nucleotides 1 to 26, 1967 to 1990, 161 to 187, 1093 to 1117 or 1312 to 1331, or 1333 to 1355 of a pneumococcal surface protein A gene of *Streptococcus pneumoniae*.

7. A PCR primer consisting essentially of the isolated DNA molecule of claim 6.

8. A hybridization probe consisting essentially of the isolated DNA molecule of claim 6.

9. An isolated DNA molecule consisting of a fragment of a pneumococcal surface protein A gene of *Streptococcus*

pneumoniae consisting of a nucleotide sequence (5' to 3')
selected from

CCGGATCCAGCTCCTGCACCAAAAAC;
GCGCGTCGACGGCTTAAACCCATTCACCATTTGG;
CCGGATCCTGAGCCAGAGCAGTTGGCTG;
CCGGATCCGCTCAAAGAGATTGATGAGTCTG;
GCGGATCCCGTAGCCAGTCAGTCTAAAGCTG;
CTGAGTCGACTGGAGTTTCTGGAGCTGGAGC;
CCGGATCCAGCTCCAGCTCCAGAAACTCCAG;
GCGGATCCTTGACCAATATTTACGGAGGAGGC;
GTTTTTGGTGCAGGAGCTGG;
GCTATGGGCTACAGGTTG;
CCACCTGTAGCCATAGC;
CCGCATCCAGCGTGCCTATCTTAGGGGCTGGTT; and
GCAAGCTTATGATATAGAAATTTGTAAC.

10. A PCR primer consisting essentially of at least one isolated DNA molecule of claim 9.

11. A hybridization probe consisting essentially of at least one isolated DNA molecule of claim 9.

12. PCR probe(s) which distinguishes between *pspA* and *pspA*-like nucleotide sequences.

13. PCR probe(s) which hybridizes to both *pspA* and *pspA*-like nucleotide sequences.

14. A *PspA* extract prepared by a process comprising growing pneumococci in a first medium containing choline chloride,

eluting live pneumococci with a choline chloride containing salt solution, and

growing the pneumococci in a second medium containing an alkanolamine and substantially no choline.

15. A *PspA* extract prepared by

growing pneumococci in a first medium containing choline chloride,

eluting live pneumococci with a choline chloride containing salt solution,

growing the pneumococci in a second medium containing an alkanolamine and substantially no choline, and purifying PspA by isolation on a choline-Sepharose affinity column.

16. An immunological composition comprising the extract of claim 14.

17. An immunological composition comprising the extract of claim 15.

18. An immunological composition comprising full length PspA.

19. A method for enhancing immunogenicity of a PspA-containing immunological composition comprising including in said composition the C-terminal portion of PspA.

20. An immunological composition comprising at least two PspAs.

21. The immunological composition of claim 20 wherein the PspAs are from different groups based on RFLP.

22. PCR amplification product from a primer as claimed in claims 3, 7, 10, 12 or 13.

23. An isolated DNA molecule consisting of a nucleotide sequence homologous to a portion of *pspA*.

24. An isolated amino acid molecule comprising pneumococcal surface protein C, PspC, of *Streptococcus pneumoniae* having alpha-helical, proline rich and repeat regions.

25. An isolated DNA molecule comprising a pneumococcal surface protein C gene of *S. pneumoniae* encoding the isolated amino acid molecule of claim 24.

26. A PCR primer consisting essentially of the isolated DNA molecule of claim 25.

27. A hybridization probe consisting essentially of the isolated DNA molecule of claim 25.

28. An immunological composition comprising the amino acid molecule of claim 24.

29. An isolated amino acid molecule of claim 24 having strong homology with pneumococcal surface protein A, PspA, of *S. pneumoniae* from amino acid 458 of PspC, corresponding to amino acid 147 of PspA, extending to a C-terminus of PspC and PspA.

30. An isolated amino acid molecule of claim 24, further comprising a signal sequence consisting essentially of a charged region followed by a hydrophobic core of amino acids.

31. An isolated amino acid molecule of claim 24, wherein the alpha-helical region further comprises a seven residue periodicity and a coiled coil region having three breaks in a heptad repeat.

32. An isolated amino acid molecule comprising pneumococcal surface protein C, PspC, of *S. pneumoniae* having

alpha-helical, proline rich and repeat regions, wherein the alpha-helical region comprises a C-terminus having substantial homology with a protection-eliciting region of PspA.

33. An isolated DNA molecule comprising a pneumococcal surface protein C gene of *S. pneumoniae* encoding the isolated amino acid molecule of claim 32.

34. A PCR primer consisting essentially of the isolated DNA molecule of claim 33.

35. A hybridization probe consisting essentially of the isolated DNA molecule of claim 33.

36. An immunological composition comprising the amino acid molecule of claim 32.

37. An isolated amino acid molecule of claim 24, further comprising a 17 amino acid, partially hydrophobic tail.

38. An isolated amino acid molecule of claim 32, further comprising a 17 amino acid, partially hydrophobic tail.

39. An isolated amino acid molecule of claim 24, further comprising an epitope of interest.

40. An isolated amino acid molecule of claim 32, further comprising an epitope of interest.

41. An immunological composition comprising the amino acid molecule of claim 39.

42. An immunological composition comprising the amino acid molecule of claim 40.




FIG. 1A




FIG. 1B




FIG. 2




FIG. 3

Fig 4




Fig 5
(Figs 5A, 5B)

| | | | | | | | | | |
|-----|------|------|------|------|-----|------|------|------|------|
| RX1 | MC25 | MC26 | MC27 | MC28 | RX1 | MC25 | MC26 | MC27 | MC28 |
|-----|------|------|------|------|-----|------|------|------|------|


9.1 Kb

7.7 Kb

4.2 Kb

3.6 Kb

| | |
|--------------|--------------|
| pLSMpspA13/2 | pLSMpspA12/6 |
| 5A | 5B |






FIG. 7




FIG. 8




Fig. 9




Figure 1




Figure 2




FIG. 13
(Sheet 1)

AMINO ACID SEQUENCES IN THE NH₂-TERMINAL END OF DIFFERENT PSPA GENES. Gap inserted to maximize alignment with related PspA sequences).

| | | | | | |
|------------|------------|------------|-------------|------------|-------------|
| ATCC6303 | MNKKKMILTS | LASVAILGTG | FVASPPTLVR | AEESPQVVER | SSLEKRYEEA |
| | KAKADTAKD | YETAKK...K | AEDAQKKYDE | DQKKTEDKAK | A.VKKVDEER |
| | QKAILAVQKA | YVEY....RE | AKDKASAERQ | IAEAKRKT.. | |
| | | | | | |
| Ac94... | MNKKKMILTS | LASVAILGAG | LVTAQPTLVR | AEEAP.VASQ | SKAEKDYDTA |
| | KRDAENAKKA | LEEAKR.... | ...AQKKYED | DQKKTEEKAK | E.EKQASEAE |
| | OKANLQYQLK | LREYIQ..KT | GDRSKIQTEM | EEAEKKHKTA | KAEFDKVRGT |
| | VIPSAARV.. | | | | |
| | | | | | |
| Bg11703pro | MNKKKMILTS | LASVAILGAG | LVTSQPTLVR | AEEAP.VASQ | SKAEKDYDAA |
| | VKKSEAAKKA | YEEAKK...K | AEDAQKKYDE | DQKKTEEKA. | ENERKAAADL |
| | TEATEVHQKA | YVRYSGSNEQ | KIKNEKILAI | | |
| | | | | | |
| | | | | | |
| Bg7322pro | MNKKKMILTS | LASVAILGAG | XVASQPTKVR | AEDAP.VANQ | SQAEKDYXAA |
| | XKKSEAAKKX | YXAKKVLAE | AEAAQKKXED | XQKKPEEKA. | EKAKAASEEI |
| | VKATEEVQKA | A..... | | | |
| | | | | | |
| | | | | | |
| Bg7561pro | MNKKKMILTS | LASVAILGAG | LVTSQPTLVR | AEEAP.GASQ | SKAEKDYXAA |
| | XKKSEAAKKA | YEEAKK...K | AEDAQKKYDE | GQKKTEEKA. | RKAEAEASKEL |
| | AKATSEVQNA | YVKYQGVORN | SRLNEKERKK | QLAEIDEEIN | KAKQIWNEKN |
| | EDFKKVREEV | IPEPTELAKD | QRKAEAEAKAE | EKVAKRKYDY | ATLKVALAKS |
| | YVEAEEAXL. | | | | |
| Bg8090pro | MNKKKMILTS | LASVAILGAG | LVTSQPTFVR | AEEAP.VASQ | PKAEKDYDPA |
| | GKKSEAATKA | YEDAKP...T | AEDAQKKYDE | AQKKPDAER. | |
| | | | | | |
| | | | | | |
| | | | | | |
| Bg8743pro | MNKKKMILTS | LASVAILGAG | LVASQPTVVR | AEEAP.VAKQ | SQAERDYDAA |
| | MKKSEAAKKE | YEEAKKOLEE | AKAAQKKYGG | DPKKTGEETK | LVPK.ADGER |
| | PKANVAVPKA | YLKLREAQEQ | LNQSPNNKKN | SAQQKLKDAL | AHIDEVTLNQ |
| | KEAEA..... | | | | |
| | | | | | |
| Bg8838pro | MNKKKMILTS | LASVAILGAG | LVTSQPTVVR | AEESP.VASQ | SKAEKDYDAA |
| | VKNATAAKKA | AEDAHRALDE | AKAAQKNYDE | DQKKPEEKAK | EVPKAPAE. |
| | | | | | |
| | | | | | |

FIG. 15
(Sheet 2)

| | | | | | |
|------------|--|--|--|--|--|
| Bg9163pro | MNKKKMILTS MKKSEAAKKE AKATEEVH.. | LASVAILGAG YEDAKKVLAE | LVASQPTLVR AEAAQKKYED | AEDAP.VANQ DQKKTEEKA. | SQAEKDYDAA ENANAASEEI |
| Bg9739pro | MNKKKMILTS KRDAENAKKA QEANKDYQLK IVVPNPQELE VVLQAXXAEI | LASVAILGAG LEEAKR.... LKKYLDGRNL MARRKSEVVK ESGGHKLEPK | LVASSPTVVR ...AQEKYAD SNSSVLKKEM ATESGLVTRV | AEEAP.VASQ YORRIEEKAA EEAEKKDKEN EEAEKNVTDA | SKAEKDYDTA K.ETQASLEQ QAEFNKIRRE RQKLVLKCNE |
| Dbl1pro | MNKKKMILTS KRDAENAKKA QKANLXYQLL | LASVAILGAG LEEAKR.... LQKYVSESDG | LVASQPTVVR ...AQKKCKED KKKKEXEXKA | AEEAP.VASQ DQKKTEEKAK DAAKKEIELK | SKAEKDYDAA X.DXQASEAE XADLXKIXQE |
| Dbl5pro | MNKKKMILTS VERSKAAEED QAATLKYHLE | LASVAILGAG LE.....E SXEFLNYFQD | LVASQPTVVR AEAAQKKYDE NHR..... | AEEAP.VASQ DQKKSEENEX | SKAEKDYDAA E.TEEASERQ |
| Dbl6aapro | MNKKKMILTS KRDAENAKKA QEANKDYQLK IVVPNPQELE EVVLQAOIA. | LASVAILGAG LEEAKR.... LKKYLDGRNL MARRKSEVVK | LVASPTTVVR ...AQEKYAD SNSSVLKKEM TKESGLVKRV | AEEAP.VASQ YORRIEEKAA EEAEKKDKEN EEAEKKVTEA | SKAEKDYDTA K.ETHASLEQ PAEFNKIRRE RPKLDAERAK |
| Dbl6apro | MNKKKMILTS KRDAENAKKA QEANKDYQLK | LASVAILGAG LEEAKR.... LKKYLDGRNL | LVASPTTVVR ...AQEKYAD SNSSVLKKEM | AEEAP.VASQ YORRIEEKAA EEAEKKDKEN | SKAEKDYDTA K.ETHASLEQ QAGL..... |
| Ef10197pro | MNKKKMILTS KRDAENAKKA QEANKDYQLK IVVPNPQELE EVVLQPTR*V | LASVAILGAG LEEAKR.... LKKYLDGRNL MARRKSEVVK ENEVHKLXQK | LVTSQPTLVR ...AQEKYAD SNSSVLKKEM AKESGLVKRV | AEEAP.VASQ YORRIEEKAA EEAEKKDKEN EEAEKKVTEA | SKAEKDYDAA K.EQQASLEQ QAEFNKIRRE RQKLDAERAK |
| Ef3296pro | MNKKKMILTS KAKADTAKKD IDVALVVQNA EQQDLQNNFN | LASVAILGAG YETAKR...K YKEY....RE EVRVAVAPDP | LVTSQPTFVR AEDAQKKYED VQNRQSKYKS TCVGKDKR.. | AEEAPQVVEK DQKRTEEKAR DADYQKKLTE | SSLEKKYEEA K.EAEASQKL VDSKIEKARK |
| Ef6796pro | MNKKKMILTS KAKYDAAKRD AKATEEVQKA IQTPFVASLT | LASVAILGAG YDEAKR...K VLDYITAIRN QMIL..... | XVTSQPTXVR AAEAQKKYEE HNDSGKTSAE | AEEAPQVVEK DQKKTEEKAE EAENKAKERD | SSLEKKYEEA K.AKAASEEI YCCAGKKFDP |
| L81905pro | MNKKKMILTS KRDAENAKKA QEANKDYQLK IVVPNPQELE | LASVAILGAG LEEAKR.... LKKYLDGRNL MA..... | LVASSPTVVR ...AQEKYAD SNSSVLKKEM | AEEAP.VASQ YORRIEEKAA EEAEKKDKEN | SKAEKDYDTA K.ETQASLEQ QAEFNKIRRE |
| Rxd1pro | MNKKKMILTS KRDARNAKKA DKAVAAVQQA | LASVAILGAG VEDAQKALDD YLAYQQATDK | FVASQPTVVR AKAAQKKYDE AAKDAADKMI | AEEAP.VASQ DQKKTEEKA. DEAKKREEEA | SKAEKDYDAA ALEKAASEEM KTRKNTVRAM |

| | | | | | |
|-----------|-------------|------------|------------|------------|------------|
| | VVPEPEQLAE | TKKKSEEAKQ | KAPELTKKLE | EAKAKLEEAE | KKATEAKQKV |
| | DA..... | | | | |
| Wu2pro | MNKKKMILTS | LASVAILGAG | LVASQPTLVR | AEESE.VASQ | SKAEKDYDAA |
| | VKKSEAAKKA | YREAKKALEE | AKVAQKKYED | DQKKTEEKA. | ELEKEASEAI |
| | AKATEEVQQA | YLAYQRASNK | A..EAAKMIE | EAQRRENEAR | AKFTTIRTIM |
| | VVPEPEQLAE | TKKKAEEAKA | KEPKLAKKAA | EAKAKLEEAE | KKATEANPOV |
| | DA..... | | | | |
| Ef5668pro | MNKKKMILTS | LASVAILGAG | FVASSPTFVR | AEEAP.VANQ | SKAEKDYDAA |
| | VKKSEAAKKA | YETAKK...K | AEDAQKKYDE | DQKKTEAKAE | K.ERKASEKI |
| | AETATKEVQQA | YLAYLOASNE | SORKEADKKI | KEATHAKMRR | TCNLTIEFEO |
| | QLYFLNQVSY | LRLRKKQKRO | QKKQKYLREN | LKRQLKRYKY | RKIKYLNKML |
| | KTKRKL.... | | | | |
| Bg6692pro | MNKKKLIVTS | LASVAILGAG | SVTSPPALVR | ADEASLIASQ | SKAEKDYDAA |
| | KKDAKNAKKA | VEDAQKALDD | AKAAQKKYDE | DQKKTEKAA | AV.KKIDEEH |
| | QAANLKSQQA | LVEFLAAQRE | GNPKKKKAAQ | ATLEBAENAE | KETK..... |
| | | | | | |
| | | | | | |
| Ac122pro | MNKKKMIKTS | LASAAIFGAX | SETSQPTRVR | PVEAPE.ARH | PKVDKYYDAE |
| | ADEY..... | | | | |
| | | | | | |
| | | | | | |
| A66pro | MNKKKMILTS | LASVAILGAG | FGCVSAYSCK | SRRISRS*SA | *SSQRL.... |
| | | | | | |
| | | | | | |
| | | | | | |
| L82013pro | MNKKKMILKS | LASAAISGAX | LVXPQPTLVR | AEESE.AASQ | SHPEQDYDXX |
| | XXLCXXLXHQ | PSXGRTLLXX | XXSXFXSPTP | XXXXXXFXSX | LTXLXPLXXX |
| | LKPFPLPX SX | PXPPXPPXSP | PSPPPRFXLY | XXPPXPPXPL | SLXLIPFLLL |
| | XLPPPXXKLP | HLXSPPXPXL | PPSPTPX... | | |
| | | | | | |

FIG. 13
(Sheet 3)

SEQUENCES IN THE CENTRAL REGION - (Includes Carboxy-terminus of alpha-heli and region and some of the proline-rich region. Gaps are inserted to maximize alignment related PspA sequences.)

30 336
0922134c

Atcc6303c

Ac122c

A66c

Ac94c

Bg8090c

Bg8743c

Bg9163c

```

.....L KEIDESDSED YLKEGLRAPL QSKLDTKKAK LSKLEELSDK
IDELDAEIAK LEVQLKDAEG NNNVE..... A.YFKEGLEK TTAERKAELE
KAEADLKAV DEPETPAPA.....PQPA PAPEKPAE.. .....K
PAPAPAP... EKPAPE.. ....K.PAEK PAEKPAEPA EKPAPEK.
.....PAPTPE .KPAPTPTPT RTGKWQENG.....
.....V LDXTIAEGKA GIAAXPPNID
KT....PKDL EDSGLGLEKV LATLDPGGET PDGLDKEASE DSNIGALPNQ
VSDLENQVSE LDREVTRLPS DLKDTGNNV GDYVKGGLK ALTDEKVGLN
NTPKALDTAP KALDTALNEL G.PDGDEEET PAPAPKPE.. .....QPA
EQP.....K. ....PAPAPK PEKTDDQQA EDYARRSEEE
YNRLPQQQPP KAEK..PAPA PKPEQVPAP .....
.....GGW SWR*ILLARP
DRLAARQAE AOKQTELGKL LDSLDPEGKT QDELDKEAGE ....AELDKK
ADGLPNKVSD LEKEISNLEI LLGGADSEDD T....AALPN KLATKKAELE
KTQKELDAAL NELG..... ..PDGDEEET PAPAPQE.. .....Q
PAPAPKPEQ. ....PTPAPK PEQPTPAPKP EQ..PAP...
.....AP KPEQ..PAPA PKPEQPAPAP KP.EQPTPGP KIE.....
.....E LLLLEKAGLG
KAGADLKEAV NEPGESAGEP SQPEEPAEEA PAPEQPTPT .....
.....QPEEP AGETPAPKPE K...PAGQPK AEKTDDQQA EDYARRSEEE
YNRLTQQQPP KAEKPAPA! PQPEQPAPAP K.....
.....L KEIDESDSED YVKEGLRVPL QSELDVKQAK LLKLEELSDK
IDELDAEIAK NLKKDVEDFQ NSGGGYS... .ALYLEAAEK DLVAKKAELE
KTEADLKAV NEPEKPAEE. ....PENPAP.. .....APK
PAPAPQ... ..EKPA... ..PAPAPK PEKSADQQA EDYARRSEEE
YNRLTQQQPP KAEKPAPAV PKPEQPAPAP KSR.....
.....VXLDRGPAEA AVKEQVDSPP QQLAD*VKEI STRGKFLGGA ATEDETSALP
NKITAKQAE AKKQTELEKL LDNLDPEGKT QDELDKEAAE ....AELDKK
ADELPNKVAD LEKBISNLEI LLGGADPEDD T....AALPN KLATKKAEFE
KTPKELDAAL NELG..... ..PDGDEEET PA.....
PAPAPKPEQ. ....PAPA.. ....PAPKP EQPAPAP...
.....AP KPEQPAPAPA PKPEQPTPAP K.....
.....L KEIDESDSED YIKEGLRAPL QSKLDAKKAK LSKLDELSK
IDELDAEIAK LEKIVGDFPN SDGEQ..... AGQYLVAEEK DLDAKEAELG
NTGADLKAV DEPETPAPA.....PAPK PAPAPAPT.. .....P
EAPAPA... PKPAPAPK.. ....PAPAPK PAPAPKPAPA PKPAPAPK..
.....PAPAPKPE RT.....
.....END

```

FIG. 13 (Sheet 5)

Bg9739c
GVQRTKRAP KRIMSLSQKV XLKXVCRAPL QSKLDAQKAE LLKLEELSGK
IEELDAEIAE LEVQLKDAEG NNNVE..... A.YFKEGLEK TTAEKKAELE
KAXADLKAV DEPETPAPA.....PAPA PAP.....A
PAPAPA.... PAPAPAPK..PAPAPK PAPAPAPAPA PKPAPAPK..
.....PAPAPAPA PKPEKPAEKP APAPKPETTK TYG.....
.....
.....L KEIDESDSED YVREGFRAPL QSELDAKQAK LSKLEELSDK
IDELDAEIAK LEKDVEDFON SDGEQ..... AGOYLAAAGE DLIAKKAELE
KAEADLKAV DEPETPAPA.....PA.. PAPAPAPT..P
EAPAPAPAPA PKPAPAPK..PAPAPK PAPAPKPAPA PKPAPAPK..
.....PAPAPAPA PKPEKPAEKP APAPKPE... ..
.....
Ef1019c
.....L KEIDESDSED YVKEGFRAPL QSELDAKQAK LSKLEELSDK
IDELDAEIAK LEDQLKAAEE NNNVE..... .DYFKEGLEK TIAAKKAELE
KTEADLKAV NEPEKPAEEP SQPEKPAEBA PAPEQPTPT QPEKPAEQPQ
PAPAPQPEKP AEETPAPKPE K...PAEQPK AEKPADQOAE EDYARRSEEE
YNRLTQQQPP KAEKPAPA.. PKTK.....
.....
Ef3296c
.....GGS ALDQEAAP HQVADLEKQI TGPEIFLGA DPEADIAARP
NELAAKQAE AOKPTGLEKL LDSLDPGGKT QDELDKAGEAELDKK
ADELPNKVAD LEKEISNLEI LLGGADSEDD T....AALPN KLAXKXAELE
KTQKELDAP NELG..... ..PDGDEET PAPAPQPE..Q
PAPAPKPEQ.PAPAPK PEQPAPAPKP EQ..PAP...
.....AP KPEQ..PAPA PKPEQPAKPE KPAEPTQPE KPATPKT...
.....
Ef6796xc
.....R
VRAL..KVAE FGVQLRDAGG SNNVG..... A.YFKEGLEE TTAEXEAGLG
KAEADLKAV DEPET..... PAP.....A
PAPAPA.... PAPAPAPK..PAPAPK PAPAPAPAPA PKPAPAPK..
.....PAPAPAPA PKPEKPAEKP APAPKPETPK T.....
.....
Db15c
.....L KDIDESDSED YAKEGLRAPL QSELDTKKAK LLKLEELSGK
IEELDAEIXE LEVQLKDAEG NNNVE..... A.YFKEGLEK TTAEKKAELE
KAEADLKAV DEPETPAPA.....PAPA PAPAPTPE..A
PAPAPA.... PKPAPAPK..PAPAPK PAPAPKPAPA PKPAPAPKPA
PAPAPAPAPK PAPAPAPAPA PKPEKPAEKP APAPKPETPK TGWKQENGM.
.....
L81905c
.....L KEIDESDSED YVKEGFRAPL QSELDAKQAK LSKLEEKSDK
XDELDAEIAK LEKDVEDFKN SDGEQ..... AGOYLAAAE DLIAKKAXLE
KAEADLKAV DEPETPAPA.....PA.. PAPAPAPT..P
EAPAPA.... ..PAPAPK..PAPAPK PAPAPKPAPA PKPAPAPK..
.....PAPAPAPA PKPEKPAA.. ..
.....
Rct115c
.....LKEIDESDVE VKKAELELVK EEAKEPRNEE KVKQAKAEVE
SKKAEATRLE KIKTDRKKAE EAKRKAAEED KVKEK.....
..PAPKPEN.PAEQPK AEKPADQOAE EDYARRSEEE
YXRLTQQQPP KTEKPAQPS PKT.....

Rct121c

```

.....
.....
.....
.....K GEARESRXEE KVNQPKXEVE
SKKXEATRLE KIKTDRKKA EAXRKAAED KVKEKP AEQF QPAPAPQPEK
PAPAPKPEN. .... PAEQPK AEKPADOQAE EDYARRSEBE
YNRLTQQOPP KTEKPAQST XK.....

```

Rct123c

```

.....
.....
.....I KEIDESXSED YLKEGLRAPL QSKLDTKKAK LSKLEELSDK
IDELDAEIAK LEVQLKDAEG NNNVE..... A.YFKEGLEK TTAEKKAELE
KAEADLKAV DEPTPAPA. .... POPA PAPEKPAE.. .....K
PAPAPAP... ..... EKPAPAPEK.
.....PAPTPE. .KPAPTPEP KTGWKQENGM WYFYNTDGS
ATGWLQNGS WYILNSNGAM ATGWHQNGS WYILNS

```

Rct129c

```

.....
.....L KEIDESDSED YLKEGLRAPL QSKLDTKKAK LSKLEELSDK
IDELDAEIAK LEVQLKDAEG NNNVE..... A.YFKEGLEK TTAEKKAELE
KAEADLKAV DEPTPAPA. .... POPA PAPEKPAE.. .....K
PAPAPAP... EKPAPAPE.. ....K.PAPA PEKPAP..AP EKPAPAPEK.
.....PAPAPE .KPAPAPEKP APAPKPETPE TRLETRKRY.

```

Rct135c

```

.....
.....L KEIDESDSED YLKEGLRAPL QSKLDTKKAK LSKLEELSDK
IDELDAEIAK LEVQLKDAEG NNNVE..... A.YFKEGLEK TTAEKKAELE
KAEADLKAV DEPTPAPA. .... POPA PAPEKPAE.. .....K
..PAPAP... EKPAPAPE.. ....K.PAPA P..... EKPAPAPEK.
.....PAPAPE .KPAPTPEP KTGWKQENGM .....

```

FIG. 13 (Sheet 6)

FIG. 13 (Sheet 4)

Rx1c

.....
.....L KEIDSESESED YAKEGFRAPL QSKLDAKRAK LSKLEELSDK
IDELDAEIAK LEDQLKAAEE NNNVE..... .DYFKEGLEK TIAAKRAELE
KTEADLKKAV NEPEKPA... ..PAPET PAPEAPAE..QPK
PAPAPQP... ..APAPKPE K...PAEQPK PEKTDDQOAE EDYARRSEEE
YNRLTQQQPP KAEKPAPA.. PKTGWKQENG MWYFYNTDGS M.....

Bg6692c

.....
.....
.....GEOA... .GOYRAAAEG DLAACKQAELE
KTEADLKKAV NEPEK..PA.PAPET PAPEAPAE..QPK
PAPAPQP... ..APAPKPE K...PAEQPK AEKTDDQOAE EDYARRSEEE
YNRLTQQQPP KAEKPAPA.. PKPEQPAPA.....

Bg8838c

.....
.....
.....PK NSKGEQA... .EQYRSAAGG DLAACKQVELE
KTEADLKKAV NEPEK..PA.PAPET PAPEAPAE..QPK
PAPAPQP... ..APAPKPE K...PAEQPK AEKPADQOAE EDYDRRSEEE
YNRLTQQQPP KAEKPAPA.. PQPEQPAPAP KS.....

Dbl6ac

.....
.....L KEIDESDSED YVKEGFRAPL QSELDKQAK LSKLEELSDK
IDELDAEIAK .LEKQVEDFK XSDGEQA... .GOYLAAAE DLIACKAELE
QTEADLKKAV NEPGKPAPA.PAPET PAPEAPAE..QPK
PAPET.P... ..APAPKPE K...PAEQPK PEKPADQOAE EDYARRSEEE
YNRLTQQQPA PAQKPEQP.. AKPEKPAEEP TOPEK.....

Dbl1c

.....
.....
.....DAEIAK .LEKNVEYFK KTDAEQT... .EQYLAAAEK DLADKKAELE
KTEADLKKAV NEPEKPAEB.TPAPA PKPEQPAE..QPK
PAPAPQP... ..APAPKP.EKTDDQOAE EDYARRSEEE
YNRLPQQQPP KAEKPAPA.. PKPEQPVP.. ..

L820131c

.....
.....
.....A EXPENPAP..APK
PAXAPOPLKP EEPAEQPKPE KPPEPAGOPE PEKPDDQOAG EDYARRSGGE
YNRFPPQQQPP KAEKPAPA.. PKPEQPVPAP KT.....

Bg11703c

.....
.....LLKA KLAGAKSKAA
TKKAELEPEL EKAEAELENL LSTLDPEGKT QDELDKEAAEAELNKK
VEALPNQVSE LEBELSKLED NLKDAETNNV EDYIKEGLEE AIATKQABLE
KT.....P KELDAALNEL G.PDGDEEET PPPEAPAE..QPK
PEK.PAEET.PAPAPK PEKSADQOAE EDYARRSEEE
YNRLTQQQPP KAEKPAPAPA PKPEQPAPAP KSR.....

Bg7817c

.....
.....GLATKKKL NLAEARIELL
LKKLGLEPGL EKAGAGLGNL LSTLDPEGKT QDELDKEAAEAELNKK
VEALPNQVAE LEEELSKLED NLKDAETNHV EDYIKEGLEE AIATKQABLE
KT.....P KELDAALNEL G.PDGDEEET PAPEAPAE..QPK
PEK.PAEET.PAPAPK PEKSADQOAE EDYARRSEEE

Bg7561c

YNRLTQQQPP KAEKPAPAPA FKPEQPAPAP K.....

 KKQKVNLENL LSTLDPGGKT QDELDRGAAEAELNKK
 VEALPNPVXB LEEELSPPED NLKDAETNHV EDYIKEGLEE AIATKQAELE
 ET.....P QEVDAALNDL V.PDGGEET PAPAPQPD... ..EPA
 PAPAPNAEQ.PAPAPK PEKSADQQA E DYARRSEGE
 YNRLTQQQPP KAEKPAPAPA FKPEQPAPAP N.....

Ef5668c

.....KEIAR LQSDLKDAEE NNVEDYIKEG LEQAITNKKA ELATTQONID
 KT....QKDL EDAELELEKV LATLDPEGKT QDELDRGAAEAELNEK
 VEALONQVAE LEEELSKLED NLKDAETNNV EDYIKEGLEE AIATKKA E LE
 KT.....Q KELDAALNEL G.PDGDEET PAPAPQPE... ..KPA
 EEP....EN.PAPAPK PEKSADQQA E DYARRSEKE
 YNRLTQQQPP KAEK..PAPA POPEQPAPAP KIE.....

Wu2c

.....
L KEIDSESED YAKEGFRAPL HSKLDAKKAK LSKLEELSDK
 IDELDAEIAK LEDQLKAVEE NNNVE..... .DYSTEGLK TIAAKKTELE
 KTEADLKKAV NEPEKSABEP SQPEKPAAEA PAPEQPTPT
QPEKP AEETPAPKPE K...PAEQPN AEKTDDQQA E DYARRSEEE
 YNRLTQQQPP KAEKPAPA.. POPEQTSSLH

FIG. 13 (Sheet 8)

Complete sequence for *EF5668* *pSP8*
Sequence Range: 1 to 1453

```

      10      20      30      40      50      60      70
      *      *      *      *      *      *      *
TTGACAAATA TTTACGGAGG AGGCTTATGC TTAATATAAG TATAGGCTAA AAATGATTAT CAGAAAAGAG

      80      90     100     110     120     130
      *      *      *      *      *      *      *
GTAAATTTAG ATG AAT AAG AAA AAA ATG ATT TTA ACA AGC CTA GCC AGC GTC GCT ATC TTA GGG
      M  N  K  K  K  M  I  L  T  S  L  A  S  V  A  I  L  G>

     140     150     160     170     180     190
      *      *      *      *      *      *
GCT GGT TTT GTT GCG TCT TCG CCT ACT TTT GTA AGA GCA GAA GAA GCT CCT GTA GCT AAC
      A  G  F  V  A  S  S  F  T  F  V  R  A  E  E  A  P  V  A  N>

     200     210     220     230     240     250
      *      *      *      *      *      *
CAG TCT AAA GCT GAG AAA GAC TAT GAT GCA GCA GTG AAA AAA TCT GAA CCT GCT AAG AAA
      Q  S  K  A  E  K  D  Y  D  A  A  V  K  K  S  E  A  A  K  K>

     260     270     280     290     300     310
      *      *      *      *      *      *
GAT TAC GAA ACG GCT AAA AAG AAA GCA GAA GAC GCT CAG AAG AAA TAT GAT CAG GAT CAG
      D  Y  E  T  A  K  K  K  A  E  D  A  Q  K  K  Y  D  E  D  Q>

     320     330     340     350     360     370
      *      *      *      *      *      *
AAG AAA ACT GAG GCA AAA GCG GAA AAA GAA AGA AAA GCT TCT GAA AAG ATA GCT GAG GCA
      K  K  T  E  A  K  A  E  K  E  R  K  A  S  E  K  I  A  E  A>

     380     390     400     410     420     430
      *      *      *      *      *      *
ACA AAA GAA GTT CAA CAA GCG TAC CTA GCT TAT CTA CAA GCT AGC AAC GAA AGT CAG AGA
      T  K  E  V  Q  Q  A  Y  L  A  Y  L  Q  A  S  N  E  S  Q  R>

     440     450     460     470     480     490
      *      *      *      *      *      *
AAA GAG GCA GAT AAG AAG ATA AAA GAA GCT ACG CAC GCA AAG ATG AGG CGG ACG TGC AAT
      K  E  A  D  K  K  I  K  E  A  T  H  A  K  M  R  R  T  C  N>

     500     510     520     530     540     550
      *      *      *      *      *      *
TTG ACT ATC GAA TTC GAA CAA CAA TTG TAC TTC CTG AAC CAA GTG AGT TAC CTG AGA CTA
      L  T  I  H  F  H  Q  Q  L  Y  F  L  N  Q  V  S  Y  L  R  L>

     560     570     580     590     600     610
      *      *      *      *      *      *
AGA AAA AAG CAG AAG AGG CAA CAA AAG AAG CAG AAG TAT CTA AGA AAA AAT CTG AAG AGG
      R  K  K  Q  K  R  Q  Q  K  K  Q  K  Y  L  R  K  N  L  K  R>

     620     630     640     650     660     670
      *      *      *      *      *      *
CAG CTA AAG ACG TAT AAG TAT AGA AAA ATA AAA TAC TTG AAC AAG ATG CTG AAA ACG AAA
      Q  L  K  R  Y  E  Y  R  K  I  K  Y  L  N  K  M  L  X  T  E>

     680     690     700     710     720     730
      *      *      *      *      *      *
AGA AAA TTG ACG TAC TTC AAA ACA AAG TCG CTG ATT TAT AAA AAG GAA TTG CTC TCC ATC
      R  K  L  T  Y  F  K  T  K  S  L  I  Y  K  K  E  L  L  S  I>

     740     750     760     770     780     790
      *      *      *      *      *      *
AAA ACA GTC GCT GAA TTA AAT AAA GAA ATT GCT AGA CTT CAA AGC GAT TTA AAA GAT GCT
      K  T>
      V  A  E  L  N  I  E  I  A  R  L  Q  S  D  L  K  D  A>

     800     810     820     830     840     850
      *      *      *      *      *      *
GAA GAA AAT AAT GTA GAA GAC TAC ATT AAA GAA GGT TTA GAG CAA GCT ATC ACT AAT AAA
      E  E  N  N  V  E  D  Y  I  K  E  G  L  E  Q  A  I  T  N  K>

     860     870     880     890     900     910
      *      *      *      *      *      *

```

- FIG-13 (Sheet 9)

AAA GCT GAA TTA GCT ACA ACT CAA CAA AAC ATA GAT AAA ACT CAA AAA GAT TTA GAG GAT
K A E L A T T Q Q N I D K T Q K D L E D>
920 930 940 950 960 970
GCT GAA TTA GAA CTT GAA AAA GTA TTA GCT ACA TTA GAC CCT GAA GGT AAA ACT CAA GAT
A E L E L E K V L A T L D P E G K T Q D>
980 990 1000 1010 1020 1030
GAA TTA GAT AAA GAA GCT GCT GAA GCT GAG TTG AAT GAA AAA GTT GAA GCT CTT CAA AAC
E L D K E A A E A E L N E K V E A L Q D>
1040 1050 1060 1070 1080 1090
CAA GTT GCT GAA TTA GAA GAA GAA CTT TCA AAA CTT GAA GAT AAT CTT AAA GAT GCT GAA
Q V A E L E E E L S K L E D N L K D A D>
1100 1110 1120 1130 1140 1150
ACA AAC AAC GTT GAA GAC TAC ATT AAA GAA GGT TTA GAA GAA GCT ATC GCG ACT AAA AAA
T N N V E D Y I K E G L E E A I A T K K>
1160 1170 1180 1190 1200 1210
GCT GAA TTG GAA AAA ACT CAA AAA GAA TTA GAT CCA CCT CTT AAT GAG TTA GCG CCT GAT
A E L E K T Q K E L D A A L N E L G P D>
1220 1230 1240 1250 1260 1270
GGA GAT GAA GAA GAG ACT CCA GCG CCG GCT CCT CAA CCA GAA AAA CCA GCT GAA GAG CCT
G D E E E T P A P A P Q P E K P A E E P>
1280 1290 1300 1310 1320 1330
GAG AAT CCA GCT CCA CCA CCA AAA CCA GAG AAG TCA CCA GAT CAA CAA GCT GAA GAA GAC
E N P A P A P K P E K S A D Q Q A E E D>
1340 1350 1360 1370 1380 1390
TAT GCT COT AGA TCA GAA GAA GAA TAT AAT CGC TTG ACC CAA CAG CAA CCG CCA AAA GCA
Y A R R S E E E Y N R L T Q Q Q P P K A>
1400 1410 1420 1430 1440 1450
GAA AAA CCA GCT CCT GCA CCA CAA CCA GAG CAA CCA GCT CCT GCA CCA AAA ATA GAG GC
E K P A P A P Q P E Q P A P A P K I E A>

FIG-13 - (Sheet 10)

Figure 14





Figure 215



Primer LSM13: gcaagcttatgatatagaaatttgtaac

Primer LSM2: gcgcgtcgacggcttaaaccattcaccattgg

Probe LSMpspA13/2 (from RX1 sequence):

```

aagcttatga tatagaaatt tgtaacaaaa atgtaatatata aaacacttga
caaatattta cggaggaggc ttatacttaa tataagtata gtctgaaaat
gactatcaga aaagaggtaa atttagatga ataagaaaaa aatgatttta
acaagtctag ccagcgtcgc tatcttaggg gctggttttg ttgcgtctca
gcctactgtt gtaagagcag aagaatctcc cgtagccagt cagtctaaag
ctgagaaaaga ctatgatgca gcgaagaaag atgctaagaa tgcgaaaaaa
gcagtagaag atgctcaaaa ggcttttagat gatgcaaaag ctgctcagaa
aaaatatgac gaggatcaga agaaaactga ggagaaagcc gcgctagaaa
aagcagcgtc tgaagagatg gataaggcag tggcagcagt tcaacaagcg
tatctagcct atcaacaagc tacagacaaa gccgcaaaag acgcagcaga
taagatgata gatgaagcta agaaacgcga agaagaggca aaaactaaat
ttaataactgt tcgagcaatg gtagttcctg agccagagca gttggctgag
actaagaaaa aatcagaaga agctaaacaa aaagcaccag aacttactaa
aaaactagaa gaagctaaag caaaattaga agaggctgag aaaaaagcta
ctgaagccaa acaaaaagtg gatgctgaag aagtcgctcc tcaagctaaa
atcgctgaat tggaaaatca agttcataga ctagaacaag agctcaaaga
gattgatgag tctgaatcag aagattatgc taaagaaggt ttccgtgctc
ctcttcaatc taaattggat gccaaaaaag ctaaactatc aaaacttgaa
gagttaagtg ataagattga tgagttagac gctgaaattg caaaacttga
agatcaactt aaagctgctg aagaaaacaa taatgtagaa gactacttta
aagaaggttt agagaaaact attgctgcta aaaaagctga attagaaaaa
actgaagctg accttaagaa agcagttaat gagccagaaa aaccagctcc
agctccagaa actccagccc cagaagcacc agctgaacaa ccaaaaccag
cgccggctcc tcaaccagct cccgcaccaa aaccagagaa gccagctgaa
caaccaaaac cagaaaaaac agatgatcaa caagctgaag aagactatgc
tcgtagatca gaagaagaat ataatcgctt gactcaacag caaccgccaa
aagctgaaaa accagctcct gcaccaaaaa caggctggaa acaagaaaaac
ggtatgtggt acttctacaa tactgatggt tcaatggcga caggatggct
ccaaaacaac gggtcatggt actacctcaa cagcaatggt gctatggcta
caggttggct ccaatacaat gggtcatggt attacctcaa cgctaacggc
gctatggcaa caggttgggc taaagtcaac gggtcatggt actacctcaa
cgctaattggt gctatggcta caggttggct ccaatacaac gggtcatggt
attacctcaa cgctaacggc gctatggcaa caggttgggc taaagtcaac
gggtcatggt actacctcaa cgctaattggt gctatggcta caggttggct
ccaatacaac gggtcatggt actacctcaa cgctaacggc gctatggcta
caggttgggc taaagtcaac gggtcatggt actacctcaa cgctaattggt
gctatggcaa caggttgggt gaaagatgga gatacctggt actatcttga
agcatcaggt gctatgaaag caagccaatg gttcaaagta tcagataaat
ggtactatgt caatggttta ggtgcccttg cagtcaacac aactgtagat
ggctataaag tcaatgccaa tggatgaatg gtttaagccg

```

Figure 2C


Primer LSM12: ccg gatcc agcgtc gctatcttaggggctgggt
Primer LSM6: ctgagtcgactggagtttctggagctggagc

Probe LSMpsA12/6 (from RX1 sequence):

ccagcgtcgc tatcttaggg gctgggttttg ttgcgtctca gcctactgtt
gtaagagcag aagaatctcc cgtagccagt cagtctaaag ctgagaaaga
ctatgatgca gcgaagaaag atgctaagaa tgcgaaaaaa gcagtagaag
atgctcaaaa ggcttttagat gatgcaaaag ctgctcagaa aaaatatgac
gaggatcaga agaaaactga ggagaaagcc gcgctagaaa aagcagcgtc
tgaagagatg gataaggcag tggcagcagt tcaacaagcg tatctagcct
atcaacaagc tacagacaaa gccgcaaaag acgcagcaga taagatgata
gatgaagcta agaaacgcga agaagaggca aaaactaaat ttaatactgt
tcgagcaatg gtagttcctg agccagagca gttggctgag actaagaaaa
aatcagaaga agctaaacaa aaagcaccag aacttactaa aaaactagaa
gaagctaaag caaaattaga agaggctgag aaaaaagcta ctgaagccaa
acaaaaagtg gatgctgaag aagtcgctcc tcaagctaaa atcgctgaat
tggaaaatca agttcataga ctagaacaag agctcaaaga gattgatgag
tctgaatcag aagattatgc taaagaaggt ttccgtgctc ctcttcaatc
taaattggat gccaaaaaag ctaaactatc aaaacttgaa gagttaagtg
ataagattga tgagttagac gctgaaattg caaaacttga agatcaactt
aaagctgctg aagaaaacaa taatgtagaa gactacttta aagaagggtt
agagaaaact attgctgcta aaaaagctga attagaaaaa actgaagctg
accttaagaa agcagttaat gagccagaaa aaccagctcc agctccagaa
actccag

Figure 12D

Figure 3b






Fig. 4




Figure 5 18

| | |
|-------|--|
| SKH2 | 5' CAT ACC gTT TTC TTg TTT CCA gCC -3' |
| LSM13 | 5' gCA AgC TTA TgA TAT AgA AAT TTg TAA C -3' |
| N192 | 5' ggAAggCCATATgCTCAAAGAgATTgATgAgTCT -3' |
| C588 | 5' CCAAggATCCTTAAACCCATTACCAATTggC -3' |

Figure 8.19




Figure 2 Comparison of the structural motifs of PspA and PspC. PspA has a smaller α-helical region, and does not contain the direct repeats within the α-helix (indicated by the dashed lines). The α-helical regions which are homologous between PspA and PspC are indicated by the striped pattern. PCR primers are indicated by the arrows.

AAGCTTATGC TTGTCATATA TCACAAATAT GTAGATCATA TCTGTGTATAG GACAGTAAAA CATCCTAAAT ACCTTTTAAA 80
 TATTTTACCT GAGTTGATG GCTTGACCTT GTTGAGTCAT GCCATATAGA CTTTTGTCTT AGTTTTCCTCA GTTTTATTCAG 160
 TTATTTTGTG TCGACGATA GCTGAAGAGG AAAAGTTAT ACATGAGCTT ATATATCCCA ATGGAAGCAT AAAGAGATAA 240
 ATACAAATTT CGATTTATAT ACAGTTTATA TTGAGTGTAT ATAGTAAGGT TAAAGAAAA ATATagaagg aATTAACAT Met> 320
 GTTTCATCA AAAAGCGAAA GAAAGTACA TTATTCATTT CGTAATTTTA GTATGGAGT AGCTAGTGTA GCTGTTGCCA 400
 PheAlaSer LysSerGlu ArgLysValHis TyrSerIle ArgLysPhe SerIleGlyVal AlaSerVal AlaValAla> -11
 GCTTGTCTT AGGAGGAGTA GTCCATGCAG AAGGGTTAG AAGTGGGAT AACCTCACGG TTACATCTAG TGGGCAAGAT 480
 SerLeuPheLeu GlyGlyVal ValHisAla GluGlyValArg SerGlyAsn AsnLeuThr ValThrSerSer GlyGlnAsp> 17
 ATATCGAAGA AGTATGCTGA TGAAGTCGAG TCCCATCTAG AAGTATATTT GAAGGATGTC AAAAANAATTT TGAANAAGAT 560
 IleSerLys LysTyrAlaAsp GluValGlu SerHisLeu GluSerIleLeu LysIleLeu LysIleVal LysLysAsn LeuLysLysVal> 44
 TCACATACC CAAATGTCTG GCTTATTTAC AAGTTGAGC GAAATTTAAA AGAATATTTT GTATGACTTA AAGTTTATG 640
 GluHisThr GlnAsnVal GlyLeuIleThr LysLeuSer GluIleLys LysLysTyrLeu TyrAspLeu LysValAsn> 70
 TTTTATCGGA AGCTGAGTTG ACGTCAAAA CAAAGAAAC AAAAGNAAG TTAACCCCAA CTTTGTAGCA GTTTAANAANA 720
 ValLeuSerGlu AlaGluLeu ThrSerLys, ThrLysGluThr LysGluLys LeuThrAla ThrPheGluGln PheLysLys> 97
 GATACATTAC CAACAGAAC AGAANAAG GTAGCAGAG CTCAGAGAA GGTGAGAA GCTAAGNAA AAGCCGAGGA 800
 AspThrLeu ProThrGluPro GluLysLys ValAlaGlu AlaGlnLysLys ValGluGlu AlaLysLys LysAlaGluAsp> 124
 TCANAAGAA AAGATCGCC GTAATACCC AACCATTACT TACANAAGC TTGAACCTGA AATTCCTGAG TCCGATGTGG 880
 GlnLysGlu LysAspArg ArgAsnTyrPro ThrIleThr TyrLysThr LeuGluLeuGlu IleAlaGlu SerAspVal> 150
 AAGTTAANA AGCGGAGCTT GAACTAGTAA AAGTGAAAG TAAAGGAACT CAAAGACGAGG AANAATTTAA GCAAGCAGAA 960
 GluValLysLys AlaGluLeu GluLeuVal LysValLysAla LysGluSer GlnAspGlu GluLysIleLys GlnAlaGlu> 177
 GCGGAAGTTG AGAGTAAACA AGCTGAGGCT ACAAGGTTAA AANAATCAA GACAGATCTGT GAAGAAGCTA AACGANAAGC 1040
 AlaGluVal GluSerLysGln AlaGluAla ThrArgLeu LysLysIleLys ThrAspArg GluGluAla LysArgLysAla> 204
 AGATGCTAAG TTGAAGGAAG CTGTTGAAA GATGTAGCG ACTTCAGAGC AAGATTAACC AAAGAGGCGG GCANAACGAG 1120
 AspAlaLys LeuLysGlu AlaValGluLys AsnValAla ThrSerGlu GlnAspLysPro LysArgArg AlaLysArg> 230
 GAGTTTCTGG AGAGCTAGCA ACACCTGATA AAAAGAAAA TGAATCCGAG TCCTCAGATT CTACCGTAGG TGAAGAAACT 1200
 GlyValSerGly GluLeuAla ThrProAsp LysLysGluAsn AspAlaLys SerSerAsp SerSerValGly GluGluThr> 257
 CTTCCAGGCC CATCCCTTAA TATGGCAAT GAAAGTCAGA CAGAACATAG GAAAGATGTC GATGAATATA TAAANAANAAT 1280

LeuProSer ProSerLeuAsn MetAlaAsn GluSerGln ThrGluHisArg LysAspVal AsnIleuTyr IleLysLysMet> 284
GTTGAGTGAG ATCCATTAG ATAGAGGAMA ACATACCCAM AATGTCTACT' TAAACATAMA GTTGAGCGCA ATTAAACGA 1360
LeuSerGlu IleGlnLeu AspArgArgLys HisThrGln AsnValasn LeuAsnIleLys LeuSerAla IleLysThr> 310
AGTATTGTGA TGAATTAGT GTTTTAAAG AGAACTCGA AAGAAGAG TTGACGTCA AACCAMAGC AGAGTTAAC 1440
LysTyrLeuTyr GluLeuSer ValLeuLys GluAsnSerLys LysGluGlu LeuThrSer LysThrLysAla GluLeuThr> 337
GCAGCTTTTG AGCAGTTTAA AAGATACA TTGAACCAG AAAAAAGGT AGCAGAGGT GAGAAGAGG TTGAAGAGC 1520
AlaAlaPhe GluGlnPheLys LysAspThr LeuLysPro GluLysLysVal AlaGluAla GluLysLys ValGluGluAla> 364
TAAGAAAMA GCCAAGGATC AAAAAAGA AGATCGCCGT AACTACCCAA CCATACCTTA CAAACGCTT GAACCTGMAA 1600
LysLysLys AlaLysAsp GlnLysGluGlu AspArgArg AsnTyrPro ThrAsnThrTyr LysThrLeu GluLeuGlu> 390
TTCTGTAGTC CGATGTGAA GTTAAGAG CCGAGCTTGA ACTAGTAAA GAGGAAGCTA ACGATCTCG AACGAGGAA 1680
IleAlaGluSer AspValLys ValLysGlu AlaGluLeuGlu LeuValLys GluGluAla AsnGluSerArg AsnGluGlu> 417
AAATTNAGC AAGCAAGA GAACTTGAG AGTAANAAG CTGAGGCTAC AGGTTAGAA AATATCAGA CAGATCGTAA 1760
LysIleLys GlnAlaLysGlu LysValGlu SerLysLys AlaGluAlaThr ArgLeuGlu LysIleLys ThrAspArgLys> 444
AAAGCAGAA GAAGAAGCTA ACGAAAAGC AGAAGATCT' GAGAAAMAG CTGCTGAGC CAAACAAAAMA GTGGATGCTG 1840
LysAlaGlu GluGluAla LysArgLysAla GluGluSer GluLysLys AlaAlaGluAla LysGlnLys ValaspAla> 470
AAGATATGC TCTTGAGCT AATATCGCTG AGTTGGATTA TGAACTTCAG AGACTAGAA AAGAGCTCA AGAGATTGAT 1920
GluGluTyrAla LeuGluAla, LysIleAla GluLeuGluTyr GluValGln ArgLeuGlu LysGluLeuLys GluIleasp> 497
GAGTCTGACT CAGAGATTTA TCTTAAGAA GGCCTCCG'G CTCCTCTTCA ATCTAATTG GATACCNAAA AAGCTAAMCT 200
GluSerAsp SerGluAspTyr LeuLysGlu GlyLeuArg AlaProLeuGln SerLysLeu AspThrLys LysAlaLysLeu> 524
ATCAAAACTT GAAGATTGA GTGATAAGT TGATAGTTA GACGC'GAAA TTGCNAACT TGAAGTTCA CTTAAGATG 2080
SerLysLeu GluGluLeu SerAspLysIle AspGluLeu AspAlaGlu IleAlaLysLeu GluValGln LeuLysasp> 550
CTGAAGGAA CATAATGTA GAAGCTACT TTAAGAGAG TTTAGAGAA ACTACTGCTG AGNAAAAGC TGAATTAGAA 2160
AlaGluGlyAsn AsnAsnVal GluAlaTyr PheLysGluGly LeuGluLys ThrThrAla GluLysLysAla GluLeuGlu> 577
AAGCTGAAG CTGACCTTAA GAAGCAGTT GATGAGCCAG AACTCCAGC TCCGCTCCT CAACCAAGTC CAGCTCCAGA 2240
LysAlaGlu AlaAspLeuLys LysAlaVal AspGluPro GluThrProAla ProAlaPro GlnProAla ProAlaProGlu> 604
AAACCAGCT GAAAACCAG CTCCAGCTCC AGAAMACCA GCTCCAGCTC CAGNAAACC AGCTCCAGCT CCAGNAAAC 2320
LysProAla GluLysPro AlaProAlaPro GluLysPro AlaProAla ProGluLysPro AlaProAla ProGluLys> 630
CAGCTCCAGC TCCAGNAAA CCAGCTCCAG CTCCAGNAAA ACCAGCTCCA ACTCCAGAAA CTCCNAAAC AGGCTGMAA 2400
ProAlaProAla ProGluLys ProAlaPro AlaProGluLys ProAlaPro ThrProGlu ThrProLysThr GlyTrpLys> 657

CAAGAAACG GTATGTGGTA CTTCTACAAT ACTGATGGTT CANTGGCAAC AGGCTGGCTC CAAACAATG GCTCATGGTA 2480
 GlnGluAsn GlyMetTrpTyr PheTyrAsn ThrAspGly SerMetAlaThr GlyTrpLeu GlnAsnAsn GlySerTrpTyr> 684

 CTACCTCAAC AGCAATGGCG CTATGGCGAC AGGATGGCTC CAAACAATG GCTCATGGTA CTACCTCAAC AGCAATGGCG 2560
 TyrLeuAsn SerAsnGly AlaMetAlaThr GlyTrpLeu GlnAsnAsn GlySerTrpTyr TyrLeuAsn SerAsnGly> 710

 CTATGGCGAC AGGATGGCTC CAATACAATG GTTCATGGTA CTACCTCAAC GCTNAATGGTG ATATGGCGAC AGGATGGCTC 2640
 AlaMetAlaThr GlyTrpLeu GlnTyrAsn GlySerTrpTyr TyrLeuAsn AlaAsnGly AspMetAlaThr GlyTrpLeu> 737

 CAATACAATG GTTCATGGTA CTACCTCAAC GCTNAATGGTG ATATGGCGAC AGGATGGTTC CAATACAATG GTTCATGGTA 2720
 GlnTyrAsn GlySerTrpTyr TyrLeuAsn AlaAsnGly AspMetAlaThr GlyTrpPhe GlnTyrAsn GlySerTrpTyr> 764

 CTACCTCAAC GCTNAATGGTG ATATGGCGAC AGGATGGTTC CAATACAATG GTTCATGGTA CTACCTCAAC GCTNAATGGTG 2800
 TyrLeuAsn AlaAsnGly AspMetAlaThr GlyTrpPhe GlnTyrAsn GlySerTrpTyr TyrLeuAsn AlaAsnGly> 790

 ATATGGCGAC AGGATGGCTC CAATACAATG GTTCATGGTA CTACCTCAAC AGCAATGGTG CTATGGTAAC AGGATGGCTC 2880
 AspMetAlaThr GlyTrpLeu GlnTyrAsn GlySerTrpTyr TyrLeuAsn SerAsnGly AlaMetValThr GlyTrpLeu> 817

 CAAACAATG GCTCATGGTA CTACCTCAAC GCTNAATGGTG ATATGGCGAC AGGATGGTG AATGCTGGTA 2960
 GlnAsnAsn GlySerTrpTyr TyrLeuAsn AlaAsnGly SerMetAlaThr AspTrpVal LysAspGly AspThrTrpTyr> 844

 CTATCTTGAA GCATCAGGTG CTATGAAAGC AGCCCAATGG TTCNAAGTAT CAGATAAATG GTACTATGTC AATGGCTCAG 3040
 TyrLeuGlu AlaSerGly AlaMetLysAla SerGlnTrp PheLysVal SerAspLysTrp TyrTrpVal AsnGlySer> 870

 GTGCCCTTGC AGTCAACACA .ACTGTAGAT GCTATAGAGT CAATGCCAAT GGTGAATGGG TAAACTAAAC TTAATATAAC 892
 GlyAlaLeuAla ValAsnThr ThrValAsp SerTyrArgVal AsnAlaAsn GlyGluTrp ValAsn>

 TAGTTAATAC TGACTTCCTG TMAGAACTCT TTAAAGTATT CCCTACAAAT ACCATATCCT TTCAGTAGAT AATATACCCCT 3200

 TGTAGGAAGT TTAGATTAAA AATAACTCT GTAATCCTA GCCGGATTTA TAGCGCTAGA GACTACGGAG TTTTITTGAT 3280

 GAGGAAAGAA TGGCGGCATT CAAGAGACTC TTTAAGAGAG TTACGGGCTTT TAAACTATTIA AGCTTCTCC AATTGCAAGA 3360

 GGGCTTCAAT CTCTGCTAGG TGCTAGCTTG CGAAATGGCT CCCACGGAGT TTGGCRGGGC CAGATGTTCC ACGGAGGTAG 3440

 TGAGGAGCGA GGCCCGGGA TTC

Figure 2 Amino acid and nucleotide sequence of PspC.
 A putative -10 and -35 regions are underlined. A ribosomal binding site is in lower case.

Figure 9²². Bestfit analysis of PspA and PspC. Percent identity is 69% and percent similarity is 77%. Amino acids of PspA are on the bottom line (1-588) and amino acids of PspC are on the top line (249-891). A dashed line indicates identity.

| | a | b | c | d | e | f | g | |
|-----|-----|-----|-----|-----|-----|-----|-----|---|
| 1 | | | | | | | | Glu Gly Val Arg Ser Gly Asn Asn Leu Thr |
| 11 | | | | | | | | Val Thr Ser Ser Gly |
| 16 | | Gln | Asp | Ile | Ser | Lys | Lys | |
| 22 | Tyr | Ala | Asp | Glu | Val | Glu | Ser | |
| 29 | | | His | Leu | Glu | Ser | Ile | |
| 34 | Leu | Lys | Asp | Val | Lys | Lys | Asn | |
| 41 | Leu | Lys | Lys | | | | | |
| 44 | Val | Gln | His | Thr | Gln | Asn | Val | |
| 51 | | | Gly | Leu | Ile | Thr | Lys | |
| 56 | Leu | Ser | Glu | Ile | Lys | Lys | Lys | |
| 63 | | | | | | | Tyr | |
| 64 | Leu | Tyr | Asp | Leu | Lys | | | |
| 69 | Val | Asn | Val | Leu | Ser | Glu | Ala | |
| 76 | | | Glu | Leu | Thr | Ser | Lys | |
| 81 | | | | Thr | Lys | Glu | Thr | |
| 85 | Lys | Glu | Lys | Leu | Thr | Ala | Thr | |
| 92 | Phe | Glu | Gln | Phe | Lys | Lys | Asp | |
| 99 | | | | | | | | Thr Leu Pro Thr Glu Pro |
| 105 | | | | | Glu | Lys | Lys | |
| 108 | Val | Ala | Glu | Ala | Gln | Lys | Lys | |
| 115 | Val | Glu | Glu | Ala | Lys | Lys | Lys | |
| 122 | | | | Ala | Glu | Asp | Gln | |
| 126 | Lys | Glu | Lys | Asp | Arg | Arg | Asn | |
| 133 | Tyr | Pro | Thr | Ile | Thr | | | |
| 138 | Tyr | Lys | Thr | Leu | Glu | Leu | Glu | |
| 145 | Ile | Ala | Glu | Ser | Asp | Val | Glu | |
| 152 | Val | Lys | Lys | Ala | Glu | Leu | Glu | |
| 159 | Leu | Val | Lys | Val | Lys | Ala | Lys | |
| 166 | Glu | Ser | Gln | Asp | Glu | Glu | Lys | |
| 173 | Ile | Lys | Gln | Ala | Glu | Ala | Glu | |
| 180 | Val | Glu | Ser | Lys | Gln | Ala | Glu | |
| 187 | | | | Ala | Thr | Arg | | |
| 190 | Leu | Lys | Lys | Ile | Lys | Thr | Asp | |
| 197 | Arg | Glu | Glu | Ala | Lys | Arg | Lys | |
| 204 | | Ala | Asp | Ala | Lys | Leu | Lys | |
| 210 | | Glu | Ala | Val | Glu | Lys | Asn | |
| 216 | Val | Ala | Thr | Ser | Glu | Gln | Asp | |
| 223 | Lys | | | | | | | |
| 224 | | | | | | | | Pro Lys Arg Arg Ala Lys Arg Gly Val Ser |
| 234 | | | | | | | | Gly Glu Leu Ala Thr Pro Asp Lys Lys Glu |
| 244 | | | | | | | | Asn Asp Ala Lys Ser Ser Asp Ser Ser Val |
| 254 | | | | | | | | Gly Glu Glu Thr Leu Pro Ser Pro Ser Leu |
| 264 | | | | | | | | Asn Met Ala Asn |
| 268 | | | | | Glu | Ser | Gln | |
| 271 | | Thr | Glu | His | Arg | Lys | Asp | |
| 277 | Val | Asp | Glu | Tyr | | | | |
| 281 | Ile | Lys | Lys | Met | Leu | Ser | Glu | |
| 288 | Ile | Gln | Leu | Asp | Arg | Arg | Lys | |
| 295 | | | His | Thr | Gln | Asn | Val | |
| 300 | | | Asn | Leu | Asn | Ile | Lys | |
| 305 | Leu | Ser | Ala | Ile | Lys | Thr | Lys | |
| 312 | | | | Tyr | Leu | Tyr | Glu | |
| 316 | Leu | Ser | Val | Leu | Lys | Glu | Asn | |
| 323 | | | | | Ser | Lys | | |
| 325 | Lys | Glu | Glu | Leu | Thr | Ser | Lys | |
| 332 | | | | Thr | Lys | Ala | Glu | |
| 336 | Leu | Thr | Ala | Ala | Phe | Glu | Gln | |

| | | |
|-----|-----------------------------|---------------------|
| 343 | Phe Lys Lys | |
| 346 | | Asp Thr Leu Lys Pro |
| 351 | | Glu Lys Lys |
| 354 | Val Ala Glu Ala Glu Lys Lys | |
| 361 | Val Glu Glu Ala Lys Lys Lys | |
| 368 | | Ala Lys Asp Gln |
| 372 | Lys Glu Glu Asp Arg Arg Asn | |
| 379 | | Tyr |
| 380 | | Pro Thr Asn Thr |
| 384 | Tyr Lys Thr Leu Glu Leu Glu | |
| 391 | Ile Ala Glu Ser Asp Val Lys | |
| 398 | Val Lys Glu Ala Glu | |
| 403 | Leu Glu Leu Val Lys Glu Glu | |
| 410 | Ala Asn Glu Ser Arg Asn Glu | |
| 417 | Glu Lys Ile Lys Gln Ala | |
| 423 | Lys Glu Lys Val Glu Ser Lys | |
| 430 | Lys Ala Glu Ala Thr Arg | |
| 436 | Leu Glu Lys Ile Lys Thr Asp | |
| 443 | Arg Lys Lys Ala Glu Glu Glu | |
| 450 | | Ala Lys Arg Lys |
| 454 | Ala Glu Glu Ser Glu Lys Lys | |
| 461 | Ala Ala Glu Ala Lys Gln Lys | |
| 468 | Val Asp Ala Glu Glu Tyr Ala | |
| 475 | | Leu Glu Ala Lys |
| 479 | Ile Ala Glu Leu Glu Tyr Glu | |
| 486 | Val Gln Arg Leu Glu Lys Glu | |
| 493 | Leu Lys Glu | |
| 496 | Ile Asp Glu Ser Asp Ser Glu | |
| 503 | Asp Tyr Leu Lys Glu Gly | |
| 509 | Leu Arg Ala | |
| 512 | | Pro Leu Gln Ser Lys |
| 517 | Leu Asp Thr Lys Lys Ala Lys | |
| 524 | Leu Ser Lys | |
| 527 | Leu Glu Glu Leu Ser Asp Lys | |
| 534 | Ile Asp Glu Leu Asp Ala Glu | |
| 541 | Ile Ala Lys Leu Glu Val Gln | |
| 548 | Leu Lys Asp Ala Glu Gly Asn | |
| 555 | | Asn Asn |
| 557 | Val Glu Ala Tyr Phe Lys Glu | |
| 564 | | Gly Leu Glu Lys Thr |
| 569 | | Thr Ala Glu Lys Lys |
| 574 | Ala Glu Leu Glu Lys Ala | |
| 580 | Glu Ala Asp Leu Lys Lys Ala | |
| 587 | Val Asp Glu | |

Figure 2. The coiled-coil motif of the α helix of PspC.
Amino acids that are not in the coiled-coil motif are in the right column.

LXS200 by LXS200 Mat... Plot
Tuesday, August 20, 1996 3:38 PM

Window Size = 30
Min. % Score = 65
Hash Value = 6

Strand = Both
Jump = 1

Scoring Matrix: DNA database matrix




Figure 24

Sequence of the alpha helical and proline region of LXS532 (PspC.D39). Nucleotides 1-516 include the upstream region, noncoding region. The alpha helical region is underlined (bp 517-1112) and the proline region is italicized (1113-1326).

```
1  CCAAGCTATT AGGTGACACT ATAGAATACT CAAGCTATGC ATCAAGCTTA
51  TGCTTGTCAA TAATCACAAA TATGTAGATC ATATCTTGTT TAGGACAGTA
101 AAACATCCTA ATTACTTTTT AAATATTCTT CCTGAGTTGA TTGGCTTGAC
151 CTTGTTGAGT CATGCTTATG TGACTTTTGT TTTAGTTTTT CCAGTTTATG
201 CAGTTATTTT GTATCGACGA ATAGCTGAAG AGGAAAAGCT ATTACATGAA
251 GTTATAATCC CAAATGGAAG CATAAAGAGA TAAATACAAA ATTCGATTTA
301 TATACAGTTC ATATTGAAGT AATATAGTAA GGTAAAGAA AAAATATAGA
351 AGGAAATAAA CATGTTTGCA TCAAAAAGCG AAAGAAAAGT ACATTATTCA
401 ATTCGTAAAT TTAGTATTGG AGTANCTAGT GTAGCTGTTG CCAGTCTTGT
451 TATGGGAAGT GTGGTTCATG CSACCAGARA AACGARGGAA GTACCCAAGC
501 AGCCMCTTCT TCTAATATGG CAAAGACAGA ACATAGGAAA GCYGCTAAAC
551 MAGTCGTCGA TGAATATATA GAAAAAATGT TGAGGGAGAT TCAACTAGAT
601 AGAAGAAAAC ATACCCAAAA TGTCCGCTTA AACATAAAGT TGAGCGCAAT
651 TANAACGAAG TATTTGCGTG AATTAANTGT TNTAGAAGAG AAGTCGAANN
701 ATGAGTTGCC GTCAGAAATA AAAGCGAAGT TAGACGCCGC TTTTGANAAG
751 TTTAAAAAAG ATACATTGAA ACCAGGAGAA AAGGTAGCNG AAGCTAAGAA
801 GAANGTTGAA GAAGCTAAGA AWAAAGCCRA GGATCAAAAA GAAGAAGATC
851 GYCGTAACTA CCAACCAAT ACTTRCAAAA CGCTTGACCT TGAAATTGCT
901 GAGTYCGATG TGAAAGTTAA AGAAGCGGAG CTTGAACTAG TAAARGAGGA
951 AGCTMMRGAA YCTCGAGACG AGGAAAAAAT TAAGCAAGCA AAAGCGAAAG
1001 TTGAGAGTAA AAAAGCTGAG GCTACAAGGT TAGAAAACAT CAAGACAGAT
1051 NGTAAAAAAG CAGAAGAAGA AGNTAAACGA AAAGCAGCAG AAGAAGATAA
1101 AGTTAAAGAA AAACCAGCTG AACAACCACA ACCAGCGCCG GNTACTCAAC
1151 CAGAAAAACC AGCTCCAAAA CCAGAGAAGC CAGCTGAACA ACCAAAAGCA
1201 GAAAAAACAG ATGATCAACA AGCTGAAGAA GACTATGCTC GTAGATCAGA
1251 AGAAGAATAT AATCGCTTGA NTCAACAGCA ACCGCCAAAA ACTGAAAAAC
```

Fig. 25

1301 CAGCACAACC ATNTACTCCA AAAACA

Fig. 25 (continued)

Comparison of nucleotides of pspA.Rx1 to pspC.D39. PspA is the top line (bp 875- 1322) and pspC is the bottom line (bp 877- 1326). The region which is most homologous includes the nucleotides which encode the proline region (bp 1113 pspC, bp 1128 pspA).

Percent Similarity: 77.083 Percent Identity: 74.769

```

875 AAAAAGCTAAACTATCAAACTTGAAGAGTTAAGTGATAAGATTGATGAG 924
    |||| ||| || | ||| ||| |||: || || || ||
877 AAAACGCTTGACCTTGAAA..TTGCTGAGTYCGATGTGAAAGTTAAAGAA 924

925 TTAGACGCTGAAATTGCAAACTTGAAGATCAACTTAAAGCTGCTGAAGA 974
    || |||| | | |||: |||| |::: | || ||
925 GCGGAGCTTGAAGTAGTAAARGAGGAAGCTMMRGAAYCTCGAGACGAGGA 974

975 AAACAATAATGTAGAAGACTACTTTAAAGAAGGTTTAGAGAAAAGTATTG 1024
    ||| | ||| || | | | ||| ||| |||
975 AAAAATTAAGCAAGCAAA.....AGCGAAAGTTGAGAG..... 1007

1025 CTGCTAAAAAAGCTGA.....ATTAGAAAAAGCTGAAGCTGACCTT 1065
    ||||| ||||| ||||| ||| | | | : |
1008 ....TAAAAAGCTGAGGCTACAAGGTTAGAAAACATCAAGACAGATNGT 1053

1066 AAGAAAGCAGTTAATGAGCCAGAAAAACCAGCTCCAGCTCCAGAACTCC 1115
    || ||||| || | || ||: ||| | ||| ||| |
1054 AAAAAAGCAG...AAGAAGAAGNTAAACGAAAAGCAGCAGAAGAAGATAA 1100

1116 AGCCCCAGAAGCACCAGCTGAACAACCAAAACCAGCGCCGGCTCCTCAAC 1165
    || |||| ||||| ||||| ||||| |||||: |||||
1101 AGTTAAAGAAAAACCAGCTGAACAACCACAACCAGCGCCGNTACTCAAC 1150

1166 CAG...CTCCGCACCAAAACCAGAGAAGCCAGCTGAACAACCAAAACCA 1212
    ||| || || ||||| ||||| ||||| ||||| ||||| ||
1151 CAGAAAAACCAGCTCCAAAACCAGAGAAGCCAGCTGAACAACCAAAAGCA 1200

1213 GAAAAAACAGATGATCAACAAGCTGAAGAAGACTATGCTCGTAGATCAGA 1262
    ||||| ||||| ||||| ||||| ||||| ||||| |||||
1201 GAAAAAACAGATGATCAACAAGCTGAAGAAGACTATGCTCGTAGATCAGA 1250

1263 AGAAGAATATAATCGCTTGACTCAACAGCAACCGCCAAAAGCTGAAAAAC 1312
    ||||| ||||| ||||| ||||| ||||| ||||| |||||
1251 AGAAGAATATAATCGCTTGANTCAACAGCAACCGCCAAAAGCTGAAAAAC 1300

1313 CAGCTC.....CTGCACCAAAAACA 1332
    |||| | : | |||||
1301 CAGCACAACCATNTACTCCAAAACA 1326

```

Fig. 2b

BESTFIT pspC.EF6796 and pspC.D39. pspC of EF6796 is the top line (bp 1-bp 1042) and pspC of D39 is the bottom line (bp 44 - bp 1087). The sequences are highly homologous in the upstream noncoding region and the DNA encoding the proline region.

Percent Similarity: 88.322 Percent Identity: 86.065

```

1  AAGCTTATGCTTGTCAATAATCACAAATATGTAGATCATATCTTGTTTAG 50
   ||||||||||||||||||||||||||||||||||||||||||||||||
44 AAGCTTATGCTTGTCAATAATCACAAATATGTAGATCATATCTTGTTTAG 93

51  GACAGTAAAACATCCTAATTACTTTTTAAATATTTTACCTGAGTTGATTG 100
   ||||||||||||||||||||||||||||||||||||| ||||||||
94  GACAGTAAAACATCCTAATTACTTTTTAAATATTCTTCCTGAGTTGATTG 143

101 GCTTGACCTTGTTGAGTCATGCCTATATGACTTTTGTTTTAGTTTTTCCA 150
   |||||||||||||||||||| || ||||||||||||||||||||
144 GCTTGACCTTGTTGAGTCATGCTTATGTGACTTTTGTTTTAGTTTTTCCA 193

151 GTTTATGCAGTTATTTTGTATCGACGAATAGCTGAAGAGGAAAAGTTATT 200
   ||||||||||||||||||||||||||||||||||||||||||||
194 GTTTATGCAGTTATTTTGTATCGACGAATAGCTGAAGAGGAAAAGCTATT 243

201 ACATGAAGTTATAATCCCAAATGGAAGCATAAAGAGATAAATACAAAATT 250
   ||||||||||||||||||||||||||||||||||||||||||||
244 ACATGAAGTTATAATCCCAAATGGAAGCATAAAGAGATAAATACAAAATT 293

251 CGATTTATATACAGTTCATATTGAAGTGATATAGTAAGGTAAAGAAAAA 300
   ||||||||||||||||||||||||||||||||||||||||||||
294 CGATTTATATACAGTTCATATTGAAGTAATATAGTAAGGTAAAGAAAAA 343

301 ATATAGAAGGAAATAAACATGTTTGCATCAAAAAGCGAAAGAAAAGTACA 350
   ||||||||||||||||||||||||||||||||||||||||||||
344 ATATAGAAGGAAATAAACATGTTTGCATCAAAAAGCGAAAGAAAAGTACA 393

351 TTATTCAATTCGTAAATTTAGTATTGGAGTAGCTAGTGTAGCTGTTGCCA 400
   ||||||||||||||||||||||||||||:||||||||||||||
394 TTATTCAATTCGTAAATTTAGTATTGGAGTANCTAGTGTAGCTGTTGCCA 443

401 GCTTGTTCTTAGGAGGAGTAGTCCATGCAGAAGGGGTTAGAAGTGGAAT 450
   | | | | | | | | | | | | | | | | | | | | : | | |
444 GTCTTGTTATGGGAAGTGTGGTTCATGC..SACCAGARAAACGARGGAAG 491

451 AACCTCA....CGGTTACATCTAGTGGGCAAGATATATCGAAGAAGTATG 496
   ||| | | | | | | | | | | | | | | | | | |
492 TACCCAAGCAGCCMCTTCTTCTAATATGGCAAAGACAGAACATAGGAAAG 541

497 CTGATGAA.....GTCGAGTCGCATCTAGAAAGTATATTGAAGGATGTC 540
   |:| | | | | | | | | | | | | | | | | | |
542 CYGCTAAACMAGTCGTCGATGAATATATAGAAAAAATGTTGAGGGAGATT 591

541 AAAAAAATTGAAAAAAGTTCAACATACCCAAAATGTCGGCTTAATTAC 590
   || | | | | | | | | | | | | | | | | | |
592 CAACTAGATAGAAGAA.....AACATACCCAAAATGTCGCCCTTAACAT 635

591 AAAGTTGAGCGAAATTAAAAAGAAGTATTTGTATGACTTAAAAGTTA... 637
   ||||||||| ||||:| ||||||||| ||| ||||: |||:

```

Fig. 27

Percent Similarity: 71.212 Percent Identity: 67.803

| | | |
|-----|--|-----|
| 306 | SQTEHRKD...VDEYIKKMLSEIQLDRRKHTQNVLNLIKLSAIKTKYLY | 351 |
| | | |
| 2 | AKTEHRKAAKXVVDEYIEKMLREIQLDRRKHTQNVALNIKLSAIXTKYLR | 51 |
| 352 | ELSVLKENSKEELTSKTKAELTAAFEQFKDTLKPEKKVAEAEKKVEEA | 401 |
| | | |
| 52 | ELXVXEEKS.XXELPSEIKAKLDAAFXKFKDTLKPGEKVAEAKKXVEEA | 100 |
| 402 | KKKAKDQKEEDRRNYPTNTYKTLLELEIAESDVKVKEAELELVKEEANESR | 451 |
| | : | |
| 101 | KXKAXDQKEEDRRNYPTNTXKTLDLLEIAEXDVKVKEAELELVKEEAXEXR | 150 |
| 452 | NEEKIKQAKEKVESKKAETRLEKIKTDRKKAEEEEAKRKAEESEKKAAEA | 501 |
| | : : | |
| 151 | DEEKIKQAKAKVESKKAETRLENIKTDXKKAEEXKRKAAEEDK..... | 195 |
| | . | |
| | . | |
| | . | |
| 552 | SKLDTKKAKLSKLEELSDKIDELDAEIAKLEVQLKDAEGNNNVEAYFKEG | 601 |
| | . . | |
| 196 |VKEKPAEQ..... | 203 |
| 602 | LEKTTAEKKAELEKAEADLKKAVDEPETPAPAPQPAPAPEKPAEKPAAPAP | 651 |
| | : : . . . : . . : . | |
| 204 |PQPAPXTQPEKPAPKPEKPAEQPKAEK | 230 |
| 652 | EKPAPAPEKPAPEKPAPEKP.APAPEKPAPEKPAPTPETPKT | 691 |
| | : . : . . | |
| 231 | TDDQQAEDDYARRSEEEYNRLXQOOPPKTEKPAO.PXTPKT | 270 |

Fig. 28

Percent Similarity: 56.767 Percent Identity: 42.857

Fig. 29

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/14819

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : 530/300, 324, 326, 350; 536/22.1, 23.1, 22.7; 435/91

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 530/300, 324, 326, 350; 536/22.1, 23.1, 22.7; 435/91

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, DIALOG

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|---|--------------------------------|
| X | Microbial Pathogenesis, Volume 17, issued 1994, McDaniel et al, "Localization of protection-eliciting epitopes on PspA of <i>Streptococcus pneumoniae</i> between amino acid residues 192 and 260", pages 323-337, see pages 326, 327, 328, Table 2. | 1-11, 22, 23 ----- 24-42 |
| Y | | |
| X | Microbial Pathogenesis, Volume 13, issued 1992, McDaniel et al, "Molecular localization of variable and conserved regions of <i>pspA</i> and identification of additional <i>pspA</i> homologous sequences in <i>Streptococcus pneumoniae</i> ", pages 261-269, see page 261. | 12, 13 |

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

| | | |
|---|-----|--|
| * Special categories of cited documents: | "T" | later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention |
| "A" document defining the general state of the art which is not considered to be of particular relevance | "X" | document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone |
| "E" earlier document published on or after the international filing date | "Y" | document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art |
| "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) | "Z" | document member of the same patent family |
| "O" document referring to an oral disclosure, use, exhibition or other means | | |
| "P" document published prior to the international filing date but later than the priority date claimed | | |

Date of the actual completion of the international search

16 DECEMBER 1996

Date of mailing of the international search report

17 JAN 1997

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

H. F. SIDBERRY

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US96/14819

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|---------------|--|--------------------------|
| X --- Y | Abstracts of the 93rd General Meeting of the American Society for Microbiology, issued May 1993, Crain et al., "Simultaneous Expression of Two Apparent PspA Molecules by Some Capsular group 6 Isolates of Streptococcus pneumoniae", Abstract No. D-71, page 107. See entire abstract, | 20, 21 ----- 24-42 |
| X | WO, A, 92/14488 (UAB RESEARCH FOUNDATION) 03 September 1992, see pages 11, 12 and 17. | 14-18 |

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/14819

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐

The additional search fees were accompanied by the applicant's protest.

☐

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/14819

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

A61K 38/00, 39/00, 39/02; C07H 19/00, 21/02; C07K 2/00, 4/04, 14/20; C12P 19/34

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Group I, Claims 1-11, 13, 22, and 23 drawn to an amino acid molecule, immunogens, DNA encoding the amino acid molecule, PCR, primers and probes, classified in Classes 530, 536 and 435; subclasses, 300, 324; 324, 326 350; 22.1, 91.1, 91.2.

Group II, claim 12, drawn to PCR probe which distinguishes between PspA and PspA-like molecules, classified in Class 435 and 536, subclasses 91.2 and 27.1.

Group III, claims 14-18, drawn to PspA extract obtained by a process, classified in Class 530, subclass 350.

Group IV, claims 19-21, drawn to a method of enhancing the immunogenicity of a PspA composition, classified in Class 435, subclass 69.7.

Group V, claims 24-42, drawn to amino acid molecule comprising surface protein C, composition, DNA, primers, probes, classified in Class 530, 536, 435, subclasses 300, 324, 325, 326, 350; 22.1; 91.1, 91.2.

The inventions listed as Groups I, II, III, IV, and V do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

The technical feature(s) of Group I, peptides, compositions, DNA, primers and probes of PspA, are not the technical feature of Group II, which is a probe which distinguishes between PspA and PspA-like molecules. The technical feature of Group III, is a PspA extract which is obtained by a process, PspA, proteins and extracts may be obtained by another process, and the extracted PspA of Group III, is not the technical feature which links Groups I, and II, IV or V.

The technical feature (elements) of Group IV is not the technical feature of Groups I, II, III, or V. The technical feature of Group V is PspC, which is not required in the Groupings of I, II, III or V. Further, the technical feature of Group I, the PspA protein, was known in the art.

Accordingly, the claims are not so linked by a special technical within the meaning of PCT Rule 13.2 so as to form a single inventive concept.